

EXPONENTIAL SUMS AND RANK OF TRIPLE PERSYMMETRIC MATRICES OVER $\mathbb{F}_2$

JORGEN CHERLY

RÉSUMÉ. Notre travail concerne une généralisation des résultats obtenus dans :
Exponential sums and rank of double persymmetric matrices over $\mathbf{F}_2$
arXiv : 0711.1937.

Soit $\mathbb{K}^3$ le $\mathbb{K}$ - espace vectoriel de dimension 3 où $\mathbb{K}$ dénote le corps des séries de Laurent formelles $\mathbb{F}_2((T^{-1}))$. Nous calculons en particulier des sommes exponentielles (dans $\mathbb{K}^3$) de la forme

$$\sum_{\deg Y \leq k-1} \sum_{\deg Z \leq s-1} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V \leq s+m+l-1} E(\xi YV)$$

où (t, η, ξ) est dans la boule unité de $\mathbb{K}^3$.

Nous démontrons qu'elles dépendent uniquement du rang de matrices triples persymétriques avec des entrées dans $\mathbb{F}_2$, c'est-à-dire des matrices de la forme $\begin{bmatrix} A \\ B \\ C \end{bmatrix}$

où A est une matrice $s \times k$ persymétrique, B une matrice $(s+m) \times k$ persymétrique et C une matrice $(s+m+l) \times k$ persymétrique (une matrice $[\alpha_{i,j}]$ est persymétrique si $\alpha_{i,j} = \alpha_{r,s}$ pour $i+j = r+s$). En outre, nous établissons plusieurs formules concernant des propriétés de rang de partitions de matrices triples persymétriques, ce qui nous conduit à une formule récurrente du nombre

$\Gamma_i \begin{bmatrix} s \\ s+m+l \end{bmatrix} \times k$ des matrices de rang i de la forme $\begin{bmatrix} A \\ B \\ C \end{bmatrix}$ Nous déduisons de cette

formule récurrente que si $0 \leq i \leq \inf(s-1, k-1)$, le nombre $\Gamma_i \begin{bmatrix} s \\ s+m+l \end{bmatrix} \times k$ dépend

uniquement de i. D'autre part, si $i \geq 2s+m+1, k \geq i$, $\Gamma_i \begin{bmatrix} s \\ s+m+l \end{bmatrix} \times k$ peut être

calculé à partir du nombre $\Gamma_{2s'+m'+1} \begin{bmatrix} s' \\ s'+m'+l' \end{bmatrix} \times k'$ de matrices de rang $(2s'+m'+1)$ de

la forme $\begin{bmatrix} A' \\ B' \\ C' \end{bmatrix}$ où A' est une matrice $s' \times k'$ persymétrique, B' une matrice

$(s'+m') \times k'$ persymétrique et C' une matrice $(s'+m'+l') \times k'$ où s', m', l'

et k' dépendent de i, s, m, l et k. La preuve de ce résultat est basée sur une formule du nombre de matrices de rang i de la forme $\begin{bmatrix} A \\ b_- \end{bmatrix}$ où A est une matrice

double persymétrique et b_- une matrice ligne avec entrées dans $\mathbb{F}_2$. Nous montrons également que le nombre R de représentations dans $\mathbb{F}_2[T]$ des équations

$$\begin{cases} YZ + Y_1 Z_1 + \dots + Y_{q-1} Z_{q-1} = 0 \\ YU + Y_1 U_1 + \dots + Y_{q-1} U_{q-1} = 0 \\ YV + Y_1 V_1 + \dots + Y_{q-1} V_{q-1} = 0 \end{cases}$$

associées aux sommes exponentielles

$\sum_{\deg Y \leq k-1} \sum_{\deg Z \leq s-1} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V \leq s+m+l-1} E(\xi YV)$
est donné par une intégrale sur la boule unité de $\mathbb{K}^3$ et est une combinaison linéaire

de $\Gamma_i \begin{bmatrix} s \\ s+m+l \end{bmatrix} \times k$ pour $i \geq 0$. Nous pouvons alors calculer explicitement le nombre R. Notre article est, pour des raisons de longueur, limité au cas $m \geq 0, l = 0$.

ABSTRACT. Our work concerns a generalization of the results obtained in : Exponential sums and rank of double persymmetric matrices over $\mathbb{F}_2$
arXiv : 0711.1937.

Let $\mathbb{K}^3$ be the 3-dimensional vectorspace over $\mathbb{K}$ where $\mathbb{K}$ denotes the field of Laurent Series $\mathbb{F}_2((T^{-1}))$. We compute in particular exponential sums, (in $\mathbb{K}^3$) of the form

$$\sum_{\deg Y \leq k-1} \sum_{\deg Z \leq s-1} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V \leq s+m+l-1} E(\xi YV)$$

where (t, η, ξ) is in the unit interval of $\mathbb{K}^3$. We show that they only depend on the rank of some associated triple persymmetric matrices with entries in $\mathbb{F}_2$, that is matrices of the form $\begin{bmatrix} A \\ B \\ C \end{bmatrix}$ where A is a $s \times k$ persymmetric matrix, B a $(s+m) \times k$ persymmetric matrix and C is a $(s+m+l) \times k$ persymmetric matrix (A matrix $[\alpha_{i,j}]$ is persymmetric if $\alpha_{i,j} = \alpha_{r,s}$ for $i+j = r+s$). Besides, we establish several formulas concerning rank properties of partitions of triple persymmetric

matrices, which leads to a recurrent formula for the number $\Gamma_i \begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k$ of rank i matrices of the form $\begin{bmatrix} A \\ B \\ C \end{bmatrix}$. We deduce from the recurrent formula that if

$0 \leq i \leq \inf(s-1, k-1)$ then $\Gamma_i \begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k$ depends only on i. On the other

hand, if $i \geq 2s+m+1, k \geq i$, $\Gamma_i \begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k$ can be computed from the number

$\Gamma_{2s'+m'+1} \begin{bmatrix} s'+m' \\ s'+m'+l' \end{bmatrix} \times k'$ of rank $(2s'+m'+1)$ matrices of the form $\begin{bmatrix} A' \\ B' \\ C' \end{bmatrix}$ where A' is a

$s' \times k'$ persymmetric matrix, B' a $(s'+m') \times k'$ persymmetric matrix and C' a $(s'+m'+l') \times k'$ persymmetric matrix, where s', m', l' and k' depend on i, s, m, l and k. The proof of this result is based on a formula of the number of rank i matrices of the form $\begin{bmatrix} A \\ b_- \end{bmatrix}$ where A is double persymmetric and b_- a one-row matrix with entries in $\mathbb{F}_2$. We also prove that the number R of representations in $\mathbb{F}_2[T]$ of the polynomial equations

$$\begin{cases} YZ + Y_1 Z_1 + \dots + Y_{q-1} Z_{q-1} = 0 \\ YU + Y_1 U_1 + \dots + Y_{q-1} U_{q-1} = 0 \\ YV + Y_1 V_1 + \dots + Y_{q-1} V_{q-1} = 0 \end{cases}$$

associated to the exponential sums

$$\sum_{\deg Y \leq k-1} \sum_{\deg Z \leq s-1} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V \leq s+m+l-1} E(\xi YV)$$

is given by an integral over the unit interval of $\mathbb{K}^3$, and is a linear combination of

the $\Gamma_i \begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k$ pour $i \geq 0$. We can then compute explicitly the number R. Our article is for reasons of length limited to the case $m \geq 0, l = 0$

CONTENTS

1. Exponential sums and rank of matrices of the form $\begin{bmatrix} A \\ B \end{bmatrix}$, where B is a double persymmetric matrix with entries in $\mathbb{F}_2$	8
1.1. Notation	8
1.2. Introduction	13
1.3. Statement of main results	13
1.4. Some Lemmas	17
1.5. Proofs of Theorems 1.4,1.5,1.6,1.7 and 1.8.	32
2. A recurrent formula for the number $\Gamma_i \begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k$ of rank i matrices of the form $\begin{bmatrix} A \\ B \\ C \end{bmatrix}$ such that A is a $s \times k$ persymmetric matrix over	

$\mathbb{F}_2$, B a $(s+m) \times k$ persymmetric matrix and C a $(s+m+l) \times k$ persymmetric matrix	32
2.1. Notation	32
2.2. Introduction	42
2.3. Computation of exponential sums in $\mathbb{K}^3$	42
2.4. Rank properties of partitions on triple persymmetric matrices	48
2.5. Integration formulas for exponential sums over the unit interval of $\mathbb{K}^3$	52
2.6. Some rank formulas for partitions of triple persymmetric matrices	58
2.7. A recurrent formula for the number of rank i matrices of the form $\begin{bmatrix} A \\ B \\ C \end{bmatrix}$, where A, B and C are persymmetric matrices over $\mathbb{F}_2$,	67
3. Rank properties of a partition of triple persymmetric matrices	73
3.1. Notation	73
3.2. Introduction	76
3.3. Computation of exponential sums in $\mathbb{K}^3$	77
3.4. Some rank formulas for partitions of triple persymmetric matrices related to $\sigma_{i,i,i,i}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \\ \delta \end{bmatrix} \times k}$	84
4. Study of the remainder $\Delta_i^{\begin{bmatrix} s \\ s+m \\ s+m+l \end{bmatrix} \times k}$ in the recurrent formula	93
4.1. Notation	94
4.2. Introduction	94
4.3. A rank formula for $\sigma_{i,i,i,i}^{\begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{bmatrix} \times k}$	94
4.4. A rank formula for $\Delta_i^{\begin{bmatrix} s \\ s+m \\ s+m+l \end{bmatrix} \times k}$	99
5. A recurrent formula for $\omega_i(s, m, l, k)$	101
5.1. Notation	101
5.2. Introduction	101
5.3. The recursive formula for $\omega_i(s, m, l, k)$	101
6. COMPUTATION OF $\Gamma_i^{\begin{bmatrix} s \\ s+m \\ s+m+l \end{bmatrix} \times k}$ for $i \leq s-1, i \leq k, m \geq 0, l \geq 0$	102
6.1. Notation	103
6.2. Introduction	103
6.3. Computation of $\Gamma_i^{\begin{bmatrix} s \\ s+m \\ s+m+l \end{bmatrix} \times k}$ for $i \leq s-1, i \leq k, m \geq 0, l \geq 0$	103
7. Computation of $\omega_i(s, m, l, k+1) - \omega_i(s, m, l, k)$ in the case $s \leq i \leq \inf(3s+2m+l, k), k > i$	112
7.1. Notation	112
7.2. Introduction	112
7.3. An equivalent to the formula for $\omega_i(s, m, l, k)$ in the case $s \leq i \leq \inf(3s+2m+l, k)$	112
7.4. Computation of $A_i^{[s,m,l,k]}$	113
7.5. Computation of $\omega_i(s, m, l, k+1) - \omega_i(s, m, l, k)$	122
8. Computation of $\Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}}$ for $0 \leq j \leq 2s$	131
8.1. Notation	131
8.2. Introduction	131

8.3.	A reduction formula for $\Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}}$	for $0 \leq j \leq 2s$	131
8.4.	An explicit formula for the difference		
	$\Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}}$	for $0 \leq j \leq 2s$	134
9.	An explicit formula for $\Delta_k \Gamma_i^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}}$, for $0 \leq i \leq 3s$		
9.1.	Notation		
9.2.	Introduction		
9.3.	Computation of $\Delta_k \Gamma_{s+j}^{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix}} = \Gamma_{s+j}^{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times (k+1)} - \Gamma_{s+j}^{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k}$	for $j=0,1,2$	142
9.4.	Computation of $\Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{s+j-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}}$,	for $0 \leq j \leq s-1$	146
9.5.	Computation of $\Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}}$	for $0 \leq j \leq s-1$	148
9.6.	Computation of $\Delta_k \Gamma_{2s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{2s+j-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}}$,	for $0 \leq j \leq s$	149
9.7.	Computation of $\Delta_k \Gamma_{2s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}}$	for $0 \leq j \leq s$	152
10.	An explicit formula for $\Gamma_i^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times k}$, for $s \leq i \leq 3s$, $k \geq i$		
10.1.	Notation		
10.2.	Introduction		
10.3.	Computation of $\Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times k}$	for $j=0,1,2$, $j \leq s-1$, $k \geq s+j$	156
10.4.	An explicit formula for $\Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times k}$	for $1 \leq j \leq s-1$, $k \geq s+j$	160
10.5.	Explicit formulas for $\Gamma_{2s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times k}$	where $k \geq 2s+1$ and for $\Gamma_{2s+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+1)}$	163
10.6.	Explicit formulas for $\Gamma_{2s+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times k}$	where $k \geq 2s+2$ and for $\Gamma_{2s+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+2)}$	164
10.7.	Explicit formulas for $\Gamma_{2s+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times k}$	where $k \geq 2s+3$ and for $\Gamma_{2s+3}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+3)}$	167
10.8.	Un explicit reduction formula for $\Gamma_{2s+1+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times k}$	for $0 \leq j \leq s-1$, $k \geq 2s+1+j$	169
11.	Computation of $\Delta_k \Gamma_i^{\begin{bmatrix} s+m \\ s+m \end{bmatrix}}$ for $0 \leq i \leq 3s+2m$		
11.1.	Notation		
11.2.	Introduction		
11.3.	Computation of $\Delta_k \Gamma_{s+j}^{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix}}$	for $0 \leq j \leq m-1$, $l \geq 0$	178
11.4.	Computation of $\Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}}$	for $0 \leq j \leq s-1$, $l \geq 0$	180
11.5.	Computation of $\Delta_k \Gamma_{s+m+j}^{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+m+j-1}^{\begin{bmatrix} s+m \\ s+m+l-1 \end{bmatrix}}$	for $0 \leq j \leq s-1$, $m \geq 1$, $l \geq 0$	183
11.6.	Computation of $\Delta_k \Gamma_{s+m+1}^{\begin{bmatrix} s+m \\ s+m \end{bmatrix}}$	for $m \geq 0$	185
11.7.	Computation of $\Delta_k \Gamma_{2s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+j-1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}}$	for $0 \leq j \leq l$	186

11.8.	Computation of $\Delta_k \Gamma_{2s+l+1+j}^{\begin{bmatrix} s \\ s+l \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+l+j}^{\begin{bmatrix} s \\ s+l-1 \end{bmatrix}}$	for $0 \leq j \leq s-1$	189
11.9.	Computation of $\Delta_k \Gamma_{2s+m+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+m+j-1}^{\begin{bmatrix} s \\ s+m-1 \end{bmatrix}}$	for $0 \leq j \leq m$	194
11.10.	Computation of $\Delta_k \Gamma_{2s+2m+j+1}^{\begin{bmatrix} s \\ s+m \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+2m+j}^{\begin{bmatrix} s \\ s+m-1 \end{bmatrix}}$	for $0 \leq j \leq s-1$	197
11.11.	Formulas summary		198
11.12.	Computation of $\Delta_k \Gamma_{s+m+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix}}$	for $m \geq 1, 1 \leq j \leq s-1$	199
11.13.	Computation of $\Delta_k \Gamma_{2s+m}^{\begin{bmatrix} s \\ s+m \end{bmatrix}}$	for $m \geq 1$	200
11.14.	Computation of $\Delta_k \Gamma_{2s+m+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix}}$	for $m \geq 2, 1 \leq j \leq m$	201
11.15.	Computation of $\Delta_k \Gamma_{2s+2m+1+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix}}$	for $m \geq 2, 0 \leq j \leq s-1$	204
11.16.	Summary of principal results in section 11		207
12.	An explicit formula for $\Gamma_i^{\begin{bmatrix} s \\ s+m \end{bmatrix} \times k}$,	for $s \leq i \leq 3s+2m, k \geq i$	208
12.1.	Notation		208
12.2.	Introduction		208
12.3.	Computation of $\Gamma_{s+i}^{\begin{bmatrix} s \\ s+m \end{bmatrix} \times k} - \Gamma_{s+i}^{\begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+i+1)}$,	for $0 \leq i \leq 2s+2m, k \geq s+i+1$	208
12.4.	Computation of $\Gamma_{s+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix} \times k}$,	for $m \geq 2, 0 \leq j \leq m, k \geq s+m$ and $j = m+1, k = s+m+1$	210
12.5.	Computation of $\Gamma_{s+m+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix} \times k}$,	for $1 \leq j \leq s+2, k \geq s+m+j, m \geq 2, j = s+3, k = 2s+m+3$.	216
12.6.	An explicit formula for $\Gamma_i^{\begin{bmatrix} s \\ s+1 \end{bmatrix} \times k}$,	for $s \leq i \leq 3s+2, k \geq i$	225
12.7.	Computation of $\Gamma_i^{\begin{bmatrix} s \\ s+m \end{bmatrix} \times k}$	for $2s+m+1 \leq i \leq 3s+2m, k \geq i, m \geq 2$	227
12.8.	An explicit formula for $\Gamma_i^{\begin{bmatrix} s \\ s+m \end{bmatrix} \times k}$,	for $1 \leq i \leq 3s+2m, k \geq i$	229
	References		233

CROQUIS DE LA PREUVE

- (1) Nous avons calculé explicitement le nombre de matrices de rang i de la forme $\begin{bmatrix} A \\ B \end{bmatrix}$ où B est une matrice double persymétrique.
- (2) Nous avons obtenu une formule récurrente du nombre $\Gamma_i^{\begin{bmatrix} s & m \\ s+m+l \end{bmatrix} \times k}$ des matrices de rang i de la forme $\begin{bmatrix} A \\ B \\ C \end{bmatrix}$ où A est une matrice $s \times k$ persymétrique, B une matrice $(s+m) \times k$ persymétrique et C une matrice $(s+m+l) \times k$ persymétrique
- (3) Nous ne disposons au début que d'un système de 2 équations en $3s+2m+l$ inconnues :

$$\sum_{i=0}^{\inf(3s+2m+l, k)} \Gamma_i^{\begin{bmatrix} s & m \\ s+m+l \end{bmatrix} \times k} = 2^{3k+3s+2m+l-3}$$

et

$$\sum_{i=0}^{\inf(3s+2m+l, k)} \Gamma_i^{\begin{bmatrix} s & m \\ s+m+l \end{bmatrix} \times k} \cdot 2^{-i} = 2^{2k+3s+2m+l-3} + 2^{3k-3} - 2^{2k-3}.$$

- (4) On commencera par calculer explicitement le nombre $\Gamma_i^{\begin{bmatrix} s & m \\ s+m+l \end{bmatrix} \times k}$ pour $i \leq s-1$, $i \leq k$, $m \geq 0$, $l \geq 0$, en utilisant successivement les 2 équations dans (3).
- (5) Nous calculerons ensuite explicitement le nombre $\Gamma_i^{\begin{bmatrix} s & m \\ s+m+l \end{bmatrix} \times k}$ pour $s \leq i \leq \inf(2s+m, k)$ en utilisant successivement les résultats obtenus dans (4) et (3).
- (6) A partir de ces résultats nous calculerons le nombre $\Gamma_{2s+m+1}^{\begin{bmatrix} s & m \\ s+m+l \end{bmatrix} \times k}$.
- (7) Guidé par des résultats obtenus dans (2), j'émet les hypothèses suivantes :

$$\Gamma_{2s+1+m+j}^{\begin{bmatrix} s & m \\ s+m+l \end{bmatrix} \times k} = 16^j \Gamma_{2s+1+m}^{\begin{bmatrix} s & m \\ s+m+(l-j) \end{bmatrix} \times (k-j)} \quad \text{si } 0 \leq j \leq l,$$

$$\Gamma_{2s+1+m+l+j}^{\begin{bmatrix} s & m \\ s+m+l \end{bmatrix} \times k} = 16^{2j+l} \Gamma_{2s+1+m-j}^{\begin{bmatrix} s & m-j \\ s+(m-j) \end{bmatrix} \times (k-(2j+l))} \quad \text{si } 0 \leq j \leq m,$$

$$\Gamma_{2s+1+2m+l+j}^{\begin{bmatrix} s & m \\ s+m+l \end{bmatrix} \times k} = 16^{2m+l+3j} \Gamma_{2(s-j)+1}^{\begin{bmatrix} s-j & s-j \\ s-j \end{bmatrix} \times (k-(2m+l)-3j)} \quad \text{si } 0 \leq j \leq s-1.$$

- (8) Une fois ces hypothèses vérifiées, nous pouvons utiliser (6) et (7) pour obtenir

$$\text{le nombre } \Gamma_i^{\begin{bmatrix} s & m \\ s+m+l \end{bmatrix} \times k} \text{ pour } 2s+m+2 \leq i \leq \inf(3s+2m+l, k).$$

SKETCH OF THE PROOF

- (1) We have explicitly calculated the number of rank i matrices of the form $\begin{bmatrix} A \\ B \end{bmatrix}$ where B is a double persymmetric matrix.
- (2) We have obtained a recurrent formula for the number $\Gamma_i \begin{bmatrix} s & m \\ s+m+l \end{bmatrix} \times k$ of rank i matrices of the form $\begin{bmatrix} A \\ B \\ C \end{bmatrix}$ where A is a $s \times k$ persymmetric matrix, B a $(s+m) \times k$ persymmetric matrix and C is a $(s+m+l) \times k$ persymmetric matrix.
- (3) In the beginning we only have a system of two equations in $3s+2m+l$ unknowns :

$$\sum_{i=0}^{\inf(3s+2m+l, k)} \Gamma_i \begin{bmatrix} s & m \\ s+m+l \end{bmatrix} \times k = 2^{3k+3s+2m+l-3}$$

et

$$\sum_{i=0}^{\inf(3s+2m+l, k)} \Gamma_i \begin{bmatrix} s & m \\ s+m+l \end{bmatrix} \times k \cdot 2^{-i} = 2^{2k+3s+2m+l-3} + 2^{3k-3} - 2^{2k-3}.$$

- (4) We start by calculating explicitly the number $\Gamma_i \begin{bmatrix} s & m \\ s+m+l \end{bmatrix} \times k$ pour $i \leq s-1$, $i \leq k$, $m \geq 0$, $l \geq 0$ by using successively the two equations in (3).
- (5) We then compute explicitly the number $\Gamma_i \begin{bmatrix} s & m \\ s+m+l \end{bmatrix} \times k$ for $s \leq i \leq \inf(2s+m, k)$ using successively the results obtained in (4) and (3).
- (6) From these results we calculate the number $\Gamma_{2s+m+1} \begin{bmatrix} s & m \\ s+m+l \end{bmatrix} \times k$ of triple persymmetric $(3s+2m+l) \times k$ rank $(2s+m+1)$ matrices.
- (7) Leaded by the results obtained in (2), I suggest the following hypotheses :

$$\Gamma_{2s+1+m+j} \begin{bmatrix} s & m \\ s+m+l \end{bmatrix} \times k = 16^j \Gamma_{2s+1+m} \begin{bmatrix} s & m \\ s+m+(l-j) \end{bmatrix} \times (k-j) \quad \text{si } 0 \leq j \leq l,$$

$$\Gamma_{2s+1+m+l+j} \begin{bmatrix} s & m \\ s+m+l \end{bmatrix} \times k = 16^{2j+l} \Gamma_{2s+1+m-j} \begin{bmatrix} s & m \\ s+(m-j) \end{bmatrix} \times (k-(2j+l)) \quad \text{si } 0 \leq j \leq m,$$

$$\Gamma_{2s+1+2m+l+j} \begin{bmatrix} s & m \\ s+m+l \end{bmatrix} \times k = 16^{2m+l+3j} \Gamma_{2(s-j)+1} \begin{bmatrix} s-j \\ s-j \end{bmatrix} \times (k-(2m+l)-3j) \quad \text{si } 0 \leq j \leq s-1.$$

- (8) Having proved these hypotheses, we can use (6) and (7) to obtain the number

$$\Gamma_i \begin{bmatrix} s & m \\ s+m+l \end{bmatrix} \times k \quad \text{for } 2s+m+2 \leq i \leq \inf(3s+2m+l, k).$$

1. Exponential sums and rank of matrices of the form $\begin{bmatrix} A \\ B \end{bmatrix}$, where B is a double persymmetric matrix with entries in $\mathbb{F}_2$

1.1. Notation.

1.1.1. *Analyses on $\mathbb{K}$.* We denote by $\mathbb{F}_2((\frac{1}{T})) = \mathbb{K}$ the completion of the field $\mathbb{F}_2(T)$, the field of rational functions over the finite field $\mathbb{F}_2$, for the infinity valuation $\mathbf{v} = \mathbf{v}_\infty$ defined by $\mathbf{v}(\frac{A}{B}) = \deg B - \deg A$ for each pair (A, B) of non-zero polynomials. Then every element non-zero t in $\mathbb{F}_2((\frac{1}{T}))$ can be expanded in a unique way in a convergent Laurent series $t = \sum_{j=-\infty}^{-\mathbf{v}(t)} t_j T^j$ where $t_j \in \mathbb{F}_2$.

We associate to the infinity valuation $\mathbf{v} = \mathbf{v}_\infty$ the absolute value $|\cdot|_\infty$ defined by

$$|t|_\infty = |t| = 2^{-\mathbf{v}(t)}.$$

We denote E the Character of the additive locally compact group $\mathbb{F}_2((\frac{1}{T}))$ defined by

$$E\left(\sum_{j=-\infty}^{-\mathbf{v}(t)} t_j T^j\right) = \begin{cases} 1 & \text{if } t_{-1} = 0, \\ -1 & \text{if } t_{-1} = 1. \end{cases}$$

We denote $\mathbb{P}$ the valuation ideal in $\mathbb{K}$, also denoted the unit interval of $\mathbb{K}$, i.e. the open ball of radius 1 about 0 or, alternatively, the set of all Laurent series

$$\sum_{i \geq 1} \alpha_i T^{-i} \quad (\alpha_i \in \mathbb{F}_2)$$

and, for every rational integer j , we denote by $\mathbb{P}_j$ the ideal $\{t \in \mathbb{K} \mid \mathbf{v}(t) > j\}$. The sets $\mathbb{P}_j$ are compact subgroups of the additive locally compact group $\mathbb{K}$.

All $t \in \mathbb{F}_2((\frac{1}{T}))$ may be written in a unique way as $t = [t] + \{t\}$, $[t] \in \mathbb{F}_2[T]$, $\{t\} \in \mathbb{P} (= \mathbb{P}_0)$.

We denote by dt the Haar measure on $\mathbb{K}$ chosen so that

$$\int_{\mathbb{P}} dt = 1.$$

Definition 1.1. We introduce the following definitions in $\mathbb{K}$:

- Let s, m and k denote rational integers such that $s \geq 2$, $m \geq 0$ and $k \geq 1$.
- A matrix $D = [\alpha_{i,j}]$ is said to be persymmetric if $\alpha_{i,j} = \alpha_{r,s}$ whenever $i+j = r+s$.
- Set $t = \sum_{i \geq 1} \alpha_i T^{-i} \in \mathbb{P}$, we denote by $D_{s \times k}(t)$ the following $s \times k$ persymmetric matrix

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \dots & \alpha_{s+k-2} & \alpha_{s+k-1} \end{pmatrix}.$$

- Set $\eta = \sum_{i \geq 1} \beta_i T^{-i} \in \mathbb{P}$, we denote by $D_{(s+m) \times k}(\eta)$ the following $(s+m) \times k$ persymmetric matrix

$$\begin{pmatrix} \beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \dots & \beta_{k+m-1} & \beta_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \end{pmatrix}.$$

1.1.2. *Analyses on the two-dimensional $\mathbb{K}$ -vectorspace.* Let $\mathbb{K} \times \mathbb{K} = \mathbb{K}^2$ be the 2-dimensional vector space over $\mathbb{K}$. Let $(t, \eta) \in \mathbb{K}^2$ and $|(t, \eta)| = \sup\{|t|, |\eta|\} = 2^{-\inf(\mathfrak{v}(t), \mathfrak{v}(\eta))}$.

It is easy to see that $(t, \eta) \longrightarrow |(t, \eta)|$ is an ultrametric valuation on $\mathbb{K}^2$, that is, $(t, \eta) \longrightarrow |(t, \eta)|$ is a norm and $|((t, \eta) + (t', \eta'))| \leq \max\{|(t, \eta)|, |(t', \eta')|\}$.

We denote by $d(t, \eta) = dt d\eta$ the Haar measure on $\mathbb{K}^2$ chosen so that the measure on the unit interval of $\mathbb{K}^2$ is equal to one, thus

$$\iint_{\mathbb{P} \times \mathbb{P}} d(t, \eta) = \int_{\mathbb{P}} dt \int_{\mathbb{P}} d\eta = 1 \cdot 1 = 1.$$

Let $(t, \eta) = (\sum_{i=-\infty}^{-\mathfrak{v}(t)} t_i T^i, \sum_{i=-\infty}^{-\mathfrak{v}(\eta)} \eta_i T^i) \in \mathbb{K}^2$, we denote χ the Character on $(\mathbb{K}^2, +)$ defined by

$$\chi\left(\sum_{i=-\infty}^{-\mathfrak{v}(t)} t_i T^i, \sum_{i=-\infty}^{-\mathfrak{v}(\eta)} \eta_i T^i\right) = E\left(\sum_{i=-\infty}^{-\mathfrak{v}(t)} t_i T^i\right) \cdot E\left(\sum_{i=-\infty}^{-\mathfrak{v}(\eta)} \eta_i T^i\right) = \begin{cases} 1 & \text{if } t_{-1} + \eta_{-1} = 0, \\ -1 & \text{if } t_{-1} + \eta_{-1} = 1. \end{cases}$$

Definition 1.2. We introduce the following definitions in the two-dimensional $\mathbb{K}$ -vectorspace.

- Let k, s and m denote rational integers such that $k \geq 1$, $s \geq 2$ and $m \geq 0$.
- We denote by $\mathbb{P}/\mathbb{P}_i \times \mathbb{P}/\mathbb{P}_j$ a complete set of coset representatives of $\mathbb{P}_i \times \mathbb{P}_j$ in $\mathbb{P} \times \mathbb{P}$, for instance $\mathbb{P}/\mathbb{P}_{s+k-1} \times \mathbb{P}/\mathbb{P}_{s+m+k-1}$ denotes a complete set of coset representatives of $\mathbb{P}_{s+k-1} \times \mathbb{P}_{s+m+k-1}$ in $\mathbb{P} \times \mathbb{P}$.
- Set $(t, \eta) = (\sum_{i \geq 1} \alpha_i T^{-i}, \sum_{i \geq 1} \beta_i T^{-i}) \in \mathbb{P} \times \mathbb{P}$.

We denote by $D_{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right] \times k}(t, \eta)$ any $(2s+m) \times k$ matrix, such that after a rearrangement of the rows, if necessary, we can obtain the following double persymmetric matrix $\left[\frac{D_{s \times k}(t)}{D_{(s+m) \times k}(\eta)}\right]$

$$\left(\begin{array}{cccccc} \alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \dots & \alpha_{s+k-2} & \alpha_{s+k-1} \\ \hline \beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \dots & \beta_{k+m-1} & \beta_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \end{array} \right).$$

- Set $(t, \eta) = (\sum_{i \geq 1} \alpha_i T^{-i}, \sum_{i \geq 1} \beta_i T^{-i}) \in \mathbb{P} \times \mathbb{P}$.

Let $\Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k}$ denote the number of double persymmetric $(2s+m) \times k$ rank i matrices of the form $\left[\frac{D_{s \times k}(t)}{D_{(s+m) \times k}(\eta)} \right]$, that is

$$\Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k} = \text{Card} \left\{ (t, \eta) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \mid r(D^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k}(t, \eta)) = i \right\}.$$

1.1.3. Analyses on the three-dimensional $\mathbb{K}$ -vectorspace. Let $\mathbb{K} \times \mathbb{K} \times \mathbb{K} = \mathbb{K}^3$ be the 3-dimensional vector space over $\mathbb{K}$. Let $(t, \eta, \xi) \in \mathbb{K}^3$ and $|(t, \eta, \xi)| = \sup \{|t|, |\eta|, |\xi|\} = 2^{-\inf(\mathfrak{v}(t), \mathfrak{v}(\eta), \mathfrak{v}(\xi))}$.

It is easy to see that $(t, \eta, \xi) \longrightarrow |(t, \eta, \xi)|$ is an ultrametric valuation on $\mathbb{K}^3$.

That is, $(t, \eta, \xi) \longrightarrow |(t, \eta, \xi)|$ is a norm and $|(t, \eta, \xi) + (t', \eta', \xi')| \leq \max \{|(t, \eta, \xi)|, |(t', \eta', \xi')|\}$.

We denote by $d(t, \eta, \xi) = dt d\eta d\xi$ the Haar measure on $\mathbb{K}^3$ chosen so that the measure on the unit interval of $\mathbb{K}^3$ is equal to one, that is

$$\iiint_{\mathbb{P} \times \mathbb{P} \times \mathbb{P}} d(t, \eta, \xi) = \int_{\mathbb{P}} dt \int_{\mathbb{P}} d\eta \int_{\mathbb{P}} d\xi = 1 \cdot 1 \cdot 1 = 1.$$

Let $(t, \eta, \xi) = (\sum_{i=-\infty}^{-\mathfrak{v}(t)} t_i T^i, \sum_{i=-\infty}^{-\mathfrak{v}(\eta)} \eta_i T^i, \sum_{i=-\infty}^{-\mathfrak{v}(\xi)} \xi_i T^i) \in \mathbb{K}^3$, we denote ψ the Character on $(\mathbb{K}^3, +)$ defined by

$$\begin{aligned} \psi \left(\sum_{i=-\infty}^{-\mathfrak{v}(t)} t_i T^i, \sum_{i=-\infty}^{-\mathfrak{v}(\eta)} \eta_i T^i, \sum_{i=-\infty}^{-\mathfrak{v}(\xi)} \xi_i T^i \right) &= E \left(\sum_{i=-\infty}^{-\mathfrak{v}(t)} t_i T^i \right) \cdot E \left(\sum_{i=-\infty}^{-\mathfrak{v}(\eta)} \eta_i T^i \right) \cdot E \left(\sum_{i=-\infty}^{-\mathfrak{v}(\xi)} \xi_i T^i \right) \\ &= \begin{cases} 1 & \text{if } t_{-1} + \eta_{-1} + \xi_{-1} = 0 \\ -1 & \text{if } t_{-1} + \eta_{-1} + \xi_{-1} = 1 \end{cases} \end{aligned}$$

Definition 1.3. We introduce the following definitions in the three-dimensional $\mathbb{K}$ -vectorspace.

- We denote by $\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_k$ a complete set of coset representatives of $\mathbb{P}_{k+s-1} \times \mathbb{P}_{k+s+m-1} \times \mathbb{P}_k$ in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$.

- Set $(t, \eta, \xi) = \left(\sum_{i \geq 1} \alpha_i T^{-i}, \sum_{i \geq 1} \beta_i T^{-i}, \sum_{i \geq 1} \gamma_i T^{-i} \right) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P}$ We denote by $D \left[\begin{smallmatrix} s \\ 1+m \end{smallmatrix} \right] \times k (t, \eta, \xi)$ any $(2s + m + 1) \times k$ matrix, such that after a rearrangement of the rows, if necessary, we can obtain the following triple persymmetric matrix $\left[\begin{smallmatrix} D_{s \times k}(t) \\ D_{(s+m) \times k}(\eta) \\ D_{1 \times k}(\xi) \end{smallmatrix} \right]$

$$\left(\begin{array}{cccccc} \alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \dots & \alpha_{s+k-2} & \alpha_{s+k-1} \\ \hline \beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \dots & \beta_{k+m-1} & \beta_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \\ \hline \gamma_1 & \gamma_2 & \gamma_3 & \dots & \gamma_{k-1} & \gamma_k \end{array} \right).$$

- Set

$$(\xi_1, \xi_2, \dots, \xi_n, t, \eta) = \left(\sum_{i \geq 1} \gamma_{1i} T^{-i}, \sum_{i \geq 1} \gamma_{2i} T^{-i}, \dots, \sum_{i \geq 1} \gamma_{ni} T^{-i}, \sum_{i \geq 1} \alpha_i T^{-i}, \sum_{i \geq 1} \beta_i T^{-i} \right) \in \mathbb{P}^{n+2}$$

We denote by $D \left[\begin{smallmatrix} n \\ 1+m+l \end{smallmatrix} \right] \times k (\xi, t, \eta)$ the following $(n + 2m + l + 2) \times k$ matrix, where the n first rows form a $n \times k$ matrix over the finite field $\mathbb{F}_2$, the following $(1+m)$ rows form a $(1+m) \times k$ persymmetric matrix and the $(1+m+l)$ last rows form a $(1+m+l) \times k$ persymmetric matrix.

$$\left(\begin{array}{cccccc} \gamma_{11} & \gamma_{12} & \gamma_{13} & \dots & \gamma_{1,k-1} & \gamma_{1,k} \\ \gamma_{21} & \gamma_{22} & \gamma_{23} & \dots & \gamma_{2,k-1} & \gamma_{2,k} \\ \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ \gamma_{n1} & \gamma_{n2} & \gamma_{n3} & \dots & \gamma_{n,k-1} & \gamma_{n,k} \\ \hline \alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ \alpha_{1+m} & \alpha_{2+m} & \alpha_{3+m} & \dots & \alpha_{k+m-1} & \alpha_{k+m} \\ \hline \beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{1+m+l} & \beta_{2+m+l} & \beta_{3+m+l} & \dots & \beta_{k+m+l-1} & \beta_{k+m+l} \end{array} \right)$$

- Consider the following partition of the matrix $D \begin{bmatrix} s \\ s+m \\ 1 \end{bmatrix} \times k (t, \eta, \xi)$

$$\left(\begin{array}{cccccc} \alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \dots & \alpha_{s+k-2} & \alpha_{s+k-1} \\ \beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \dots & \beta_{k+m-1} & \beta_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \\ \hline \gamma_1 & \gamma_2 & \gamma_3 & \dots & \gamma_{k-1} & \gamma_k \end{array} \right).$$

- We define

$$\sigma_{i,i}^{\left[\begin{smallmatrix} s \\ s+m \\ \gamma_1- \end{smallmatrix} \right] \times k}$$

to be the cardinality of the following set

$$\left\{ (t, \eta, \xi) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_k \mid r(D \begin{bmatrix} s \\ s+m \\ 1 \end{bmatrix} \times k (t, \eta)) = r(D \begin{bmatrix} s \\ s+m \\ 1 \end{bmatrix} \times k (t, \eta, \xi)) = i \right\}.$$

- We define

$$\sigma_{i-1,i}^{\left[\begin{smallmatrix} s \\ s+m \\ \gamma_1- \end{smallmatrix} \right] \times k}$$

to be the cardinality of the following set

$$\left\{ (t, \eta, \xi) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_k \mid r(D \begin{bmatrix} s \\ s+m \\ 1 \end{bmatrix} \times k (t, \eta)) = i-1, \quad r(D \begin{bmatrix} s \\ s+m \\ 1 \end{bmatrix} \times k (t, \eta, \xi)) = i \right\}.$$

- Set $(t, \eta, \xi) = (\sum_{i \geq 1} \alpha_i T^{-i}, \sum_{i \geq 1} \beta_i T^{-i}, \sum_{i \geq 1} \gamma_i T^{-i}) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P}$ Let $\Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \\ 1 \end{smallmatrix} \right] \times k}$ denote the number of triple persymmetric $(2s+m+1) \times k$ rank i matrices of

the form $\left[\frac{D_{s \times k}(t)}{\frac{D_{(s+m) \times k}(\eta)}{D_{1 \times k}(\xi)}} \right]$, that is

$$\Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \\ 1 \end{smallmatrix} \right] \times k} = \text{Card} \left\{ (t, \eta, \xi) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_k \mid r(D \begin{bmatrix} s \\ s+m \\ 1 \end{bmatrix} \times k (t, \eta, \xi)) = i \right\}.$$

- Set $(t, \eta, \xi) = (\sum_{i \geq 1} \alpha_i T^{-i}, \sum_{i \geq 1} \beta_i T^{-i}, \sum_{i \geq 1} \gamma_i T^{-i}) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P}$

Let $\Gamma_i^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k}$ denote the number of triple persymmetric $(2m+l+3) \times k$ rank i matrices of the form $\left[\frac{D_{1 \times k}(t)}{\frac{D_{(1+m) \times k}(\eta)}{D_{(1+m+l) \times k}(\xi)}} \right]$, that is

$$\Gamma_i^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right]^{\times k}} = \text{Card} \left\{ (t, \eta, \xi) \in \mathbb{P}/\mathbb{P}_k \times \mathbb{P}/\mathbb{P}_{k+m} \times \mathbb{P}/\mathbb{P}_{k+m+l} \mid r(D^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right]^{\times k}}(t, \eta, \xi)) = i \right\}.$$

1.2. Introduction. We generalize the results obtained in [1]

1.3. Statement of main results.

Theorem 1.4. For all $1 \leq i \leq \inf(k, 2s + m + 1)$ we have

$$\Gamma_i^{\left[\begin{smallmatrix} s+m \\ s+1 \end{smallmatrix} \right]^{\times k}} = (2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{\left[\begin{smallmatrix} s+m \\ s+1 \end{smallmatrix} \right]^{\times k}} + 2^i \cdot \Gamma_i^{\left[\begin{smallmatrix} s+m \\ s+1 \end{smallmatrix} \right]^{\times k}}$$

Theorem 1.5. We have the following reduction formulas

$$\begin{aligned} \Gamma_{m+3+j}^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right]^{\times k}} &= 2^{4j} \Gamma_{m+3}^{\left[\begin{smallmatrix} 1+m \\ 1+m+(l-j) \end{smallmatrix} \right]^{\times (k-j)}} && \text{if } 0 \leq j \leq l \\ \Gamma_{m+l+3+j}^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right]^{\times k}} &= 2^{8j+4l} \Gamma_{m+3-j}^{\left[\begin{smallmatrix} 1+m \\ 1+(m-j) \end{smallmatrix} \right]^{\times (k-(2j+l))}} && \text{if } 0 \leq j \leq m \end{aligned}$$

Theorem 1.6. We have for $m \geq 2$, $l \geq 3$ and $k \geq 2m + l$

$$\Gamma_i^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right]^{\times k}} \text{ is equal to } \begin{cases} 1 & \text{if } i = 0 \\ 2^k + 17 & \text{if } i = 1 \\ (21 \cdot 2^{3i-7} - 3 \cdot 2^{2i-5}) \cdot 2^k + 315 \cdot 2^{4i-8} - 21 \cdot 2^{3i-6} & \text{if } 2 \leq i \leq m \\ 21 \cdot 2^{k+3m-4} + 13 \cdot 2^{k+2m-3} + 315 \cdot 2^{4m-4} - 85 \cdot 2^{3m-3} & \text{if } i = m+1 \\ 2^{2k+m} + 21 \cdot 2^{k+3m-1} + 9 \cdot 2^{k+2m-1} + 315 \cdot 2^{4m} - 157 \cdot 2^{3m} & \text{if } i = m+2 \\ 3 \cdot 2^{2k-m+2i-6} + 21 \cdot 2^{k+3i-7} + 21 \cdot 2^{k-m+3i-8} + 315 \cdot 2^{4i-8} - 315 \cdot 2^{4i-m-9} & \text{if } m+3 \leq i \leq m+l \\ 3 \cdot 2^{2k+m+2l-4} + 21 \cdot 2^{k+3m+3l-4} + 149 \cdot 2^{k+2m+3l-5} + 315 \cdot 2^{4m+4l-4} - 827 \cdot 2^{3m+4l-5} & \text{if } i = m+l+1 \\ 11 \cdot 2^{2k+m+2l-2} + 21 \cdot 2^{k+3m+3l-1} + 261 \cdot 2^{k+2m+3l-2} + 315 \cdot 2^{4m+4l} - 1627 \cdot 2^{3m+4l-1} & \text{if } i = m+l+2 \\ 21 \cdot 2^{2k-2m-l+3i-9} + 21 \cdot 2^{k+3i-7} + 525 \cdot 2^{k-2m-l+4i-11} + 315 \cdot 2^{4i-8} - 3255 \cdot 2^{5i-2m-l-12} & \text{if } m+l+3 \leq i \leq 2m+l+1 \\ 53 \cdot 2^{2k+4m+2l-3} - 159 \cdot 2^{k+6m+3l-3} + 53 \cdot 2^{8m+4l-2} & \text{if } i = 2m+l+2 \\ 2^{3k+2m+l} - 7 \cdot 2^{2k+4m+2l} + 7 \cdot 2^{k+6m+3l+1} - 2^{8m+4l+3} & \text{if } i = 2m+l+3 \end{cases}$$

Theorem 1.7. Let $\Gamma_i^{\left[\begin{smallmatrix} (n) \\ 1+m \\ 1+m+l \end{smallmatrix} \right]^{\times k}}$ denote the number of matrices of the form $\begin{bmatrix} A \\ B \\ C \end{bmatrix}$ of rank i such that A is a $n \times k$ matrix over $\mathbb{F}_2$, B a $(1+m) \times k$ persymmetric matrix and C a $(1+m+l) \times k$ persymmetric matrix and where $\Gamma_i^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right]^{\times k}}$ denote the number of double persymmetric $(2+2m+l) \times k$ rank i matrices of the form $\begin{bmatrix} B \\ C \end{bmatrix}$

Then $\Gamma_i^{\left[\begin{smallmatrix} (n) \\ 1+m \\ 1+m+l \end{smallmatrix} \right]^{\times k}}$ expressed as a linear combination of the $\Gamma_{i-j}^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right]^{\times k}}$ is equal to

$$\sum_{j=0}^n 2^{(n-j) \cdot (i-j)} a_j^{(n)} \prod_{l=1}^j (2^k - 2^{i-l}) \cdot \Gamma_{i-j}^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right]^{\times k}} \quad \text{for } 0 \leq i \leq \inf(k, n+2m+l+2),$$

where $a_j^{(n)}$ satisfies the linear recurrence relation

$$a_j^{(n)} = 2^j \cdot a_j^{(n-1)} + a_{j-1}^{(n-1)}, \quad n = 2, 3, 4, \dots \quad \text{for } 1 \leq j \leq n-1.$$

The explicit value of $a_j^{(n)}$ is given by the following formula

$$a_j^{(n)} = (-1)^j \cdot 2^{jn - \frac{j(j-1)}{2}} + \sum_{s=0}^{j-1} (-1)^s \prod_{l=0}^{j-(s+1)} \frac{2^{n+1} - 2^l}{2^{j-s} - 2^l} \cdot 2^{s(n-j) + \frac{s(s+1)}{2}} \quad \text{for } 1 \leq j \leq n-1.$$

We set

$$a_0^{(n)} = a_n^{(n)} = 1$$

and $\Gamma_{i-j}^{\left[\frac{1+m}{1+m+l} \right] \times k} = 0$ if $i-j \notin \{0, 1, 2, \dots, \inf(k, 2m+l+2)\}$.

Theorem 1.8. Let $f_{m,l,k}(t, \eta, \xi_1, \xi_2, \dots, \xi_n)$ be the exponential sum in $\mathbb{P}^{n+2}$ defined by

$$(t, \eta, \xi_1, \xi_2, \dots, \xi_n) \in \mathbb{P}^{n+2} \longrightarrow \sum_{\deg Y \leq k-1} \sum_{\deg Z \leq 1+m} E(tYZ) \sum_{\deg U \leq 1+m+l} E(\eta YU) \sum_{\deg V_1 \leq 0} E(\xi_1 YV_1) \sum_{\deg V_2 \leq 0} E(\xi_2 YV_2) \dots \sum_{\deg V_n \leq 0} E(\xi_n YV_n).$$

Set

$$(t, \eta, \xi_1, \xi_2, \dots, \xi_n) = \left(\sum_{i \geq 1} \alpha_i T^{-i}, \sum_{i \geq 1} \beta_i T^{-i}, \sum_{i \geq 1} \gamma_{1i} T^{-i}, \dots, \sum_{i \geq 1} \gamma_{ni} T^{-i} \right) \in \mathbb{P}^{n+2}.$$

Then

$$f_{m,l,k}(t, \eta, \xi_1, \xi_2, \dots, \xi_n) = 2^{k+2m+l+n+4-r(D \left[\frac{\binom{n}{1+m}}{1+m+l} \right] \times k} (\xi_1, \xi_2, \dots, \xi_n, t, \eta)$$

where

$$D \left[\frac{\binom{n}{1+m}}{1+m+l} \right] \times k (\xi_1, \xi_2, \dots, \xi_n, t, \eta)$$

denotes the following $(2m+l+n+2) \times k$ matrix

$$\begin{pmatrix} \gamma_{11} & \gamma_{12} & \gamma_{13} & \dots & \gamma_{1,k-1} & \gamma_{1,k} \\ \gamma_{21} & \gamma_{22} & \gamma_{23} & \dots & \gamma_{2,k-1} & \gamma_{2,k} \\ \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ \hline \gamma_{n1} & \gamma_{n2} & \gamma_{n3} & \dots & \gamma_{n,k-1} & \gamma_{n,k} \\ \hline \alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ \hline \alpha_{1+m} & \alpha_{2+m} & \alpha_{3+m} & \dots & \alpha_{k+m-1} & \alpha_{k+m} \\ \hline \beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{1+m+l} & \beta_{2+m+l} & \beta_{3+m+l} & \dots & \beta_{k+m+l-1} & \beta_{k+m+l} \end{pmatrix}$$

Then the number of solutions

$$(Y_1, Z_1, U_1, V_1^{(1)}, V_2^{(1)}, \dots, V_n^{(1)}, Y_2, Z_2, U_2, V_1^{(2)}, V_2^{(2)}, \dots, V_n^{(2)}, \dots, Y_q, Z_q, U_q, V_1^{(q)}, V_2^{(q)}, \dots, V_n^{(q)})$$

of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0 \\ Y_1 U_1 + Y_2 U_2 + \dots + Y_q U_q = 0 \\ Y_1 V_1^{(1)} + Y_2 V_1^{(2)} + \dots + Y_q V_1^{(q)} = 0 \\ Y_1 V_2^{(1)} + Y_2 V_2^{(2)} + \dots + Y_q V_2^{(q)} = 0 \\ \vdots \\ Y_1 V_n^{(1)} + Y_2 V_n^{(2)} + \dots + Y_q V_n^{(q)} = 0 \end{cases}$$

satisfying the degree conditions

$$\deg Y_i \leq k-1, \quad \deg Z_i \leq m+1, \quad \deg U_i \leq m+l+1, \quad \deg V_j^i \leq 0, \quad \text{for } 1 \leq j \leq n \quad 1 \leq i \leq q$$

is equal to the following integral over the unit interval in $\mathbb{K}^{n+2}$

$$\int_{\mathbb{P}^{n+2}} f_{m,l,k}^q(t, \eta, \xi_1, \xi_2, \dots, \xi_n) dt d\eta d\xi_1 d\xi_2 \dots d\xi_n.$$

Observing that $f_{m,l,k}^q(t, \eta, \xi_1, \xi_2, \dots, \xi_n)$ is constant on cosets of $\mathbb{P}_{k+m} \times \mathbb{P}_{k+m+l} \times \mathbb{P}_k^n$, the above integral is equal to

$$2^{q(k+2m+l+n+4)-(2m+l+k(n+2))} \sum_{i=0}^{\inf(k, n+2m+l+2)} \Gamma_i^{\left[\begin{smallmatrix} (n) \\ 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} 2^{-iq}.$$

Example. $n=2, m=1, l=3, k \geq 5, q=3$

$$\begin{aligned} & \Gamma_i^{\left[\begin{smallmatrix} (2) \\ 2 \\ 5 \end{smallmatrix} \right] \times k} \\ &= 2^{2i} \Gamma_i^{\left[\begin{smallmatrix} 2 \\ 5 \end{smallmatrix} \right] \times k} + 3 \cdot 2^{i-1} (2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{\left[\begin{smallmatrix} 2 \\ 5 \end{smallmatrix} \right] \times k} + (2^k - 2^{i-1}) (2^k - 2^{i-2}) \cdot \Gamma_{i-2}^{\left[\begin{smallmatrix} 2 \\ 5 \end{smallmatrix} \right] \times k} \quad \text{for } 0 \leq i \leq \inf(k, 9) \end{aligned}$$

$$\Gamma_i^{\left[\begin{smallmatrix} (2) \\ 1+1 \\ 1+1+3 \end{smallmatrix} \right] \times k} = \begin{cases} 1 & \text{if } i = 0 \\ 3 \cdot 2^k + 33 & \text{if } i = 1 \\ 2^{2k} + 83 \cdot 2^k + 886 & \text{if } i = 2 \\ 33 \cdot 2^{2k} + 978 \cdot 2^k + 29352 & \text{if } i = 3 \\ 2^{3k+1} + 182 \cdot 2^{2k} + 16408 \cdot 2^k + 937408 & \text{if } i = 4 \\ 3 \cdot 2^{3k+1} + 189 \cdot 2^{2k+3} + 8191 \cdot 2^{k+6} + 12911 \cdot 2^{10} & \text{if } i = 5 \\ 3 \cdot 2^{3k+3} + 36672 \cdot 2^{2k} + 11271168 \cdot 2^k - 399015936 & \text{if } i = 6 \\ 11 \cdot 2^{3k+5} + 1022464 \cdot 2^{2k} - 100679680 \cdot 2^k + 2163212288 & \text{if } i = 7 \\ 117 \cdot 2^{3k+7} - 3354624 \cdot 2^{2k} + 214695936 \cdot 2^k - 3925868544 & \text{if } i = 8 \\ 2^{4k+5} - 480 \cdot 2^{3k+5} + 2240 \cdot 2^{2k+10} - 1920 \cdot 2^{k+16} + 2^{31} & \text{if } i = 9 \end{cases}$$

If $k = 5$ we obtain

$$\Gamma_i^{\left[\begin{smallmatrix} (2) \\ 1+1 \\ 1+1+3 \end{smallmatrix} \right] \times 5} = \begin{cases} 1 & \text{if } i = 0 \\ 129 & \text{if } i = 1 \\ 4566 & \text{if } i = 2 \\ 94440 & \text{if } i = 3 \\ 1714368 & \text{if } i = 4 \\ 31740928 & \text{if } i = 5 \end{cases}$$

Then the number of solutions

$$(Y_1, Z_1, U_1, V_1^{(1)}, V_2^{(1)}, Y_2, Z_2, U_2, V_1^{(2)}, V_2^{(2)}, Y_3, Z_3, U_3, V_1^{(3)}, V_2^{(3)})$$

of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + Y_3 Z_3 = 0 \\ Y_1 U_1 + Y_2 U_2 + Y_3 U_3 = 0 \\ Y_1 V_1^{(1)} + Y_2 V_1^{(2)} + Y_3 V_1^{(3)} = 0 \\ Y_1 V_2^{(1)} + Y_2 V_2^{(2)} + Y_3 V_2^{(3)} = 0 \end{cases}$$

satisfying the degree conditions

$$\deg Y_i \leq 4, \quad \deg Z_i \leq 2, \quad \deg U_i \leq 5, \quad \deg V_j^i \leq 0, \quad \text{for } 1 \leq j \leq 2 \quad 1 \leq i \leq 3$$

is equal to

$$\int_{\mathbb{P}^4} f_{1,3,5}^3(t, \eta, \xi_1, \xi_2) dt d\eta d\xi_1 d\xi_2 = 2^{23} \sum_{i=0}^5 \Gamma_i^{\left[\begin{smallmatrix} (2) \\ 2 \\ 5 \end{smallmatrix} \right] \times k} 2^{-3i} = 13281 \cdot 2^{20}.$$

$$\text{where for } (t, \eta, \xi_1, \xi_2) = \left(\sum_{i \geq 1} \alpha_i T^{-i}, \sum_{i \geq 1} \beta_i T^{-i}, \sum_{i \geq 1} \gamma_{1i} T^{-i}, \sum_{i \geq 1} \gamma_{2i} T^{-i} \right) \in \mathbb{P}^4.$$

$$\begin{aligned} f_{1,3,5}(t, \eta, \xi_1, \xi_2) &= \sum_{\deg Y \leq 4} \sum_{\deg Z \leq 2} E(tYZ) \sum_{\deg U \leq 5} E(\eta YU) \sum_{\deg V_1 \leq 0} E(\xi_1 YV_1) \sum_{\deg V_2 \leq 0} E(\xi_2 YV_2) \\ &= 2^{16-r(D \left[\begin{smallmatrix} (2) \\ 2 \\ 5 \end{smallmatrix} \right] \times k}(\xi_1, \xi_2 t, \eta)) \end{aligned}$$

and

$$D^{\left[\begin{smallmatrix} (2) \\ 2 \\ 5 \end{smallmatrix} \right] \times k}(\xi_1, \xi_2, t, \eta) = \begin{pmatrix} \begin{array}{ccccc} \gamma_{11} & \gamma_{12} & \gamma_{13} & \gamma_{1,4} & \gamma_{1,5} \\ \gamma_{21} & \gamma_{22} & \gamma_{23} & \gamma_{2,4} & \gamma_{2,5} \\ \hline \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 \\ \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 \\ \hline \beta_1 & \beta_2 & \beta_3 & \beta_4 & \beta_5 \\ \beta_2 & \beta_3 & \beta_4 & \beta_5 & \beta_6 \\ \beta_3 & \beta_4 & \beta_5 & \beta_6 & \beta_7 \\ \beta_4 & \beta_5 & \beta_6 & \beta_7 & \beta_8 \\ \beta_5 & \beta_6 & \beta_7 & \beta_8 & \beta_9 \end{array} \end{pmatrix}$$

1.4. Some Lemmas.

Lemma 1.9. *Let $(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P}$ and*

$$g_{s,m,k}(t, \eta, \xi) = g(t, \eta, \xi) = \sum_{\deg Y \leq k-1} \sum_{\deg Z \leq s-1} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V=0} E(\xi YV).$$

Then

$$g(t, \eta, \xi) = \begin{cases} 2^{k+2s+m-r(D \begin{bmatrix} s & \\ & s+m \end{bmatrix} \times k)(t, \eta)} & \text{if } r(D \begin{bmatrix} s & \\ & s+m \end{bmatrix} \times k)(t, \eta) = r(D \begin{bmatrix} s & \\ & 1 \end{bmatrix} \times k)(t, \eta, \xi) \\ 0 & \text{otherwise.} \end{cases}$$

Proof. We generalize the argument given in the proof of Lemma 6.2 in [1]
Consider the matrix

$$D \begin{bmatrix} s & \\ & s+m \end{bmatrix} \times k(t, \eta, \xi) = \begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \dots & \alpha_{s+k-2} & \alpha_{s+k-1} \\ \beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \dots & \beta_{k+m-1} & \beta_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \\ \hline \gamma_1 & \gamma_2 & \gamma_3 & \dots & \gamma_{k-1} & \gamma_k \end{pmatrix}.$$

It follows

$$\begin{aligned} g(t, \eta, \xi) &= 2^{2s+m} \sum_{\substack{\deg Y \leq k-1 \\ Y \in \ker D \begin{bmatrix} s & \\ & s+m \end{bmatrix} \times k(t, \eta)}} E(\xi Y) \\ &= 2^{2s+m} \cdot \left[\sum_{\substack{\deg Y \leq k-1 \\ Y \in \ker D \begin{bmatrix} s & \\ & s+m \end{bmatrix} \times k(t, \eta, \xi)}} 1 - \sum_{\substack{\deg Y \leq k-1 \\ Y \in \ker D \begin{bmatrix} s & \\ & s+m \end{bmatrix} \times k(t, \eta) \\ \sum_{j=0}^{k-1} \gamma_j \delta_j = 1}} 1 \right] \\ &= 2^{2s+m} \cdot \left[2 \cdot 2^{k-r(D \begin{bmatrix} s & \\ & s+m \end{bmatrix} \times k)(t, \eta, \xi)} - 2^{k-r(D \begin{bmatrix} s & \\ & s+m \end{bmatrix} \times k)(t, \eta)} \right]. \end{aligned}$$

□

Lemma 1.10. *Set $h(t, \eta) = \sum_{\deg \leq k-1} \sum_{\deg Z \leq s-1} E(tYZ) \sum_{\deg U \leq s+m-1} E(tYU)$ and let $q \geq 2$ be an integer, then*

$$g^q(t, \eta, \xi) = g(t, \eta, \xi) \cdot h^{q-1}(t, \eta).$$

Proof. Obviously we get

$$g^2(t, \eta, \xi) = \left[2^{2s+m} \sum_{\substack{\deg Y_1 \leq k-1 \\ Y_1 \in \ker D \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k}} E(\xi Y_1) \right] \cdot \left[2^{2s+m} \sum_{\substack{\deg Y_2 \leq k-1 \\ Y_2 \in \ker D \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k}} E(\xi Y_2) \right].$$

Define

$$\begin{cases} Y_1 + Y_2 = Y_3 & \deg Y_3 \leq k-1 \\ Y_1 = Y_4 & \deg Y_4 \leq k-1. \end{cases}$$

Then we obtain

$$\begin{aligned} g^2(t, \eta, \xi) &= 2^{2(2s+m)} \sum_{\substack{\deg Y_1 \leq k-1 \\ Y_1 \in \ker D \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k}} \sum_{\substack{\deg Y_2 \leq k-1 \\ Y_2 \in \ker D \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k}} E(\xi(Y_1 + Y_2)) \\ &= \left[2^{2s+m} \sum_{\substack{\deg Y_4 \leq k-1 \\ Y_4 \in \ker D \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k}} 1 \right] \cdot \left[2^{1+m} \sum_{\substack{\deg Y_3 \leq k-1 \\ Y_3 \in \ker D \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k}} E(\xi Y_3) \right] \\ &= h(t, \eta) \cdot g(t, \eta, \xi). \end{aligned}$$

By recurrence on q we get $g^q(t, \eta, \xi) = g(t, \eta, \xi) \cdot h^{q-1}(t, \eta)$. $\square$

Lemma 1.11. *We have*

$$\int_{\mathbb{P}} \int_{\mathbb{P}} \int_{\mathbb{P}} g^q(t, \eta, \xi) dt d\eta d\xi = 2^{q(k+2s+m)-(3k+2s+m-2)} \sum_{i=0}^{\inf(k, 2s+m)} \sigma_{i,i}^{\left[\begin{smallmatrix} s+m \\ \gamma_1- \end{smallmatrix} \right] \times k} 2^{-iq}$$

Proof. Recall Definition 1.3, then Lemma 1.9 gives, by observing that $g(t, \eta, \xi)$ is constant on cosets of $\mathbb{P}_{k+s-1} \times \mathbb{P}_{k+s+m-1} \times \mathbb{P}_k$

$$\begin{aligned} \int_{\mathbb{P}} \int_{\mathbb{P}} \int_{\mathbb{P}} g^q(t, \eta, \xi) dt d\eta d\xi &= \sum_{\substack{(t, \eta, \xi) \in \mathbb{P}_{k+s-1} \times \mathbb{P}_{k+s+m-1} \times \mathbb{P}_k \\ r(D \left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times k (t, \eta)) = r(D \left[\begin{smallmatrix} s+m \\ 1 \end{smallmatrix} \right] \times k (t, \eta, \xi))}} 2^{k+2s+m-r(D \left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times k (t, \eta))} \int_{\mathbb{P}_{k+s-1}} dt \int_{\mathbb{P}_{k+s+m-1}} d\eta \int_{\mathbb{P}_k} d\xi \\ &= \sum_{i=0}^{\inf(k, 2s+m)} \sigma_{i,i}^{\left[\begin{smallmatrix} s+m \\ \gamma_1- \end{smallmatrix} \right] \times k} 2^{q(k+2s+m-i)} 2^{-(k+s-1)} 2^{-(k+s+m-1)} 2^{-k}. \end{aligned}$$

$\square$

Lemma 1.12. *Equally*

$$\int_{\mathbb{P}} \int_{\mathbb{P}} \int_{\mathbb{P}} g^q(t, \eta, \xi) dt d\eta d\xi = 2^{q(k+2s+m)-(3k+2s+m-2)} \sum_{i=0}^{\inf(k, 2s+m)} 2^i \cdot \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k} 2^{-iq}$$

Proof. From Lemma 1.10 recalling Definition 1.2 we get by Fubini's theorem

$$\begin{aligned} \int_{\mathbb{P}} \int_{\mathbb{P}} \int_{\mathbb{P}} g^q(t, \eta, \xi) dt d\eta d\xi &= \int_{\mathbb{P}} \int_{\mathbb{P}} \int_{\mathbb{P}} g(t, \eta, \xi) h^{q-1}(t, \eta) dt d\eta d\xi \\ &= \int_{\mathbb{P}} \int_{\mathbb{P}} h^{q-1}(t, \eta) \left(\int_{\mathbb{P}} g(t, \eta, \xi) d\xi \right) dt d\eta = 2^{2s+m} \int_{\mathbb{P}} \int_{\mathbb{P}} h^{q-1}(t, \eta) dt d\eta \end{aligned}$$

$$\begin{aligned}
&= 2^{2s+m} \cdot \sum_{(t,\eta) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1}} 2^{(q-1)(2s+m+k-r(D \begin{bmatrix} s+m \\ s+m \end{bmatrix} \times k)(t,\eta))} \int_{\mathbb{P}_{k+s-1}} dt \int_{\mathbb{P}_{k+s+m-1}} dtd\eta \\
&= 2^{2s+m} \cdot \sum_{i=0}^{\inf(2s+m,k)} \sum_{(t,\eta) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1}} 2^{(q-1)(2s+m+k-i)} \int_{\mathbb{P}_{k+s-1}} dt \int_{\mathbb{P}_{k+s+m-1}} dtd\eta \\
&= 2^{2s+m} \cdot \sum_{i=0}^{\inf(2s+m,k)} \Gamma_i \begin{bmatrix} s+m \\ s+m \end{bmatrix} \times k \cdot 2^{-(q-1)i} \cdot 2^{(q-1)(2s+m+k)} \cdot 2^{-(k+s-1)} \cdot 2^{-(k+s+m-1)}.
\end{aligned}$$

□

Lemma 1.13. For all $0 \leq i \leq \inf(k, 2s+m)$ we have $\sigma_{i,i}^{\begin{bmatrix} s+m \\ \gamma_{1-} \end{bmatrix} \times k} = 2^i \cdot \Gamma_i \begin{bmatrix} s+m \\ s+m \end{bmatrix} \times k$

Proof. Lemma 1.11 and Lemma 1.12 give

$$\sum_{i=0}^{\inf(k, 2s+m)} \left(\sigma_{i,i}^{\begin{bmatrix} s+m \\ \gamma_{1-} \end{bmatrix} \times k} - 2^i \cdot \Gamma_i \begin{bmatrix} s+m \\ s+m \end{bmatrix} \times k \right) \cdot 2^{-iq} = 0 \quad \text{for all } q \geq 2.$$

□

Lemma 1.14. For all $1 \leq i \leq \inf(k, 2s+m+1)$ we have

$$\sigma_{i-1,i}^{\begin{bmatrix} s+m \\ \gamma_{1-} \end{bmatrix} \times k} = (2^k - 2^{i-1}) \cdot \Gamma_{i-1} \begin{bmatrix} s+m \\ s+m \end{bmatrix} \times k$$

Proof. Consider the matrix

$$\left[\frac{D_{s \times k}(t)}{D_{(s+m) \times k}(\eta)} \right] = \begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \dots & \alpha_{s+k-2} & \alpha_{s+k-1} \\ \hline \beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \dots & \beta_{k+m-1} & \beta_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \\ \hline \gamma_1 & \gamma_2 & \gamma_3 & \dots & \gamma_{k-1} & \gamma_k \end{pmatrix}.$$

Obviously by Lemma 1.13

$$2^k \cdot \Gamma_{i-1} \begin{bmatrix} s+m \\ s+m \end{bmatrix} \times k = \sigma_{i-1,i-1}^{\begin{bmatrix} s+m \\ \gamma_{1-} \end{bmatrix} \times k} + \sigma_{i-1,i}^{\begin{bmatrix} s+m \\ \gamma_{1-} \end{bmatrix} \times k}$$

$$\begin{aligned}
\Leftrightarrow \sigma_{i-1,i}^{\left[\frac{s+m}{\gamma_{1-}}\right] \times k} &= 2^k \cdot \Gamma_{i-1}^{\left[\frac{s+m}{\gamma_{1-}}\right] \times k} - 2^{i-1} \cdot \Gamma_{i-1}^{\left[\frac{s+m}{\gamma_{1-}}\right] \times k} \\
\Leftrightarrow \sigma_{i-1,i}^{\left[\frac{s+m}{\gamma_{1-}}\right] \times k} &= (2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{\left[\frac{s+m}{\gamma_{1-}}\right] \times k} \quad \text{if } 1 \leq i \leq \inf(k, 2s+m+1).
\end{aligned}$$

□

Lemma 1.15. *For all $1 \leq i \leq \inf(k, 2s+m+1)$ we have*

$$(1.1) \quad \Gamma_i^{\left[\frac{s+m}{\gamma_{1-}}\right] \times k} = (2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{\left[\frac{s+m}{\gamma_{1-}}\right] \times k} + 2^i \cdot \Gamma_i^{\left[\frac{s+m}{\gamma_{1-}}\right] \times k}$$

Proof. From Lemmas 1.13, 1.14 we get

$$\Gamma_i^{\left[\frac{s+m}{\gamma_{1-}}\right] \times k} = \sigma_{i,i}^{\left[\frac{s+m}{\gamma_{1-}}\right] \times k} + \sigma_{i-1,i}^{\left[\frac{s+m}{\gamma_{1-}}\right] \times k} = 2^i \cdot \Gamma_i^{\left[\frac{s+m}{\gamma_{1-}}\right] \times k} + (2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{\left[\frac{s+m}{\gamma_{1-}}\right] \times k}.$$

□

Lemma 1.16. *We have the following reduction formulas*

$$(1.2) \quad \Gamma_{m+3+j}^{\left[\frac{1+m}{1+m+l}\right] \times k} = 2^{4j} \Gamma_{m+3}^{\left[\frac{1+m}{1+m+(l-j)}\right] \times (k-j)} \quad \text{if } 0 \leq j \leq l$$

$$(1.3) \quad \Gamma_{m+l+3+j}^{\left[\frac{1+m}{1+m+l}\right] \times k} = 2^{8j+4l} \Gamma_{m+3-j}^{\left[\frac{1+m}{1+(m-j)}\right] \times (k-(2j+l))} \quad \text{if } 0 \leq j \leq m$$

Proof. From (1.1) we get with $s \rightarrow 1+m$, $m \rightarrow l$

$$(1.4) \quad \Gamma_i^{\left[\frac{1+m}{1+m+l}\right] \times k} = (2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{\left[\frac{1+m}{1+m+l}\right] \times k} + 2^i \cdot \Gamma_i^{\left[\frac{1+m}{1+m+l}\right] \times k}$$

Using the reduction formulas proved in [2, section 3] we get from (1.4) with $i \rightarrow m+3+j$.

$$\begin{aligned}
\Gamma_{m+3+j}^{\left[\frac{1+m}{1+m+l}\right] \times k} &= (2^k - 2^{m+2+j}) \cdot \Gamma_{m+2+j}^{\left[\frac{1+m}{1+m+l}\right] \times k} + 2^{m+3+j} \cdot \Gamma_{m+3+j}^{\left[\frac{1+m}{1+m+l}\right] \times k} \\
&= (2^k - 2^{m+2+j}) \cdot 2^{3j} \cdot \Gamma_{m+2}^{\left[\frac{1+m}{1+m+(l-j)}\right] \times (k-j)} + 2^{m+3+j} \cdot 2^{3j} \cdot \Gamma_{m+3}^{\left[\frac{1+m}{1+m+(l-j)}\right] \times (k-j)} \\
&= (2^{k+3j} - 2^{m+2+4j}) \cdot \Gamma_{m+2}^{\left[\frac{1+m}{1+m+(l-j)}\right] \times (k-j)} + 2^{m+3+4j} \cdot \Gamma_{m+3}^{\left[\frac{1+m}{1+m+(l-j)}\right] \times (k-j)}
\end{aligned}$$

From (1.4) with $i \rightarrow m+3$, $l \rightarrow l-j$, $k \rightarrow k-j$ we obtain

$$\Gamma_{m+3}^{\left[\frac{1+m}{1+m+(l-j)}\right] \times (k-j)} = (2^{k-j} - 2^{m+2}) \cdot \Gamma_{m+2}^{\left[\frac{1+m}{1+m+(l-j)}\right] \times (k-j)} + 2^{m+3} \cdot \Gamma_{m+3}^{\left[\frac{1+m}{1+m+(l-j)}\right] \times (k-j)}$$

which proves (1.2)

We prove (1.3) in the same way, that is

$$\begin{aligned}
\Gamma_{m+l+3+j}^{\left[\begin{smallmatrix} 1 \\ 1+m+l \end{smallmatrix} \right] \times k} &= (2^k - 2^{m+l+2+j}) \cdot \Gamma_{m+l+2+j}^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} + 2^{m+l+3+j} \cdot \Gamma_{m+l+3+j}^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} \\
&= (2^k - 2^{m+l+2+j}) \cdot 2^{6j+3l} \cdot \Gamma_{m+2-j}^{\left[\begin{smallmatrix} m+1-j \\ m+1-j \end{smallmatrix} \right] \times (k-l-2j)} + 2^{m+l+3+j} \cdot 2^{6j+3l} \cdot \Gamma_{m+3-j}^{\left[\begin{smallmatrix} m+1-j \\ m+1-j \end{smallmatrix} \right] \times (k-l-2j)} \\
&= (2^{k+6j+3l+3} - 2^{m+4l+7j+2}) \cdot \Gamma_{m+2}^{\left[\begin{smallmatrix} m+1-j \\ m+1-j \end{smallmatrix} \right] \times (k-l-2j)} + 2^{m+4l+7j+3} \cdot \Gamma_{m+3}^{\left[\begin{smallmatrix} m+1-j \\ m+1-j \end{smallmatrix} \right] \times (k-l-2j)}
\end{aligned}$$

From (1.4) with $i \rightarrow m+3-j$, $m \rightarrow m-j$, $l \rightarrow 0$, $k \rightarrow k-(2j+l)$ we obtain

$$\Gamma_{m+3-j}^{\left[\begin{smallmatrix} 1 \\ 1+(m-j) \end{smallmatrix} \right] \times (k-(2j+l))} = (2^{k-(2j+l)} - 2^{m+2-j}) \cdot \Gamma_{m+2-j}^{\left[\begin{smallmatrix} 1+(m-j) \\ 1+(m-j) \end{smallmatrix} \right] \times (k-(2j+l))} + 2^{m+3-j} \cdot \Gamma_{m+3-j}^{\left[\begin{smallmatrix} 1+(m-j) \\ 1+(m-j) \end{smallmatrix} \right] \times (k-(2j+l))}$$

which proves (1.3)

□

Lemma 1.17. *We have for $m \geq 2$, $k > m+3$*

$$\Gamma_{m+3}^{\left[\begin{smallmatrix} 1 \\ 1+m+l \end{smallmatrix} \right] \times k} = \begin{cases} 3 \cdot 2^{2k+m} + 21 \cdot 2^{k+3m+2} + 21 \cdot 2^{k+2m+1} + 315 \cdot 2^{4m+4} - 315 \cdot 2^{3m+3} & \text{if } l \geq 3 \\ 3 \cdot 2^{2k+m} + 21 \cdot 2^{k+3m+2} + 149 \cdot 2^{k+2m+1} + 315 \cdot 2^{4m+4} - 827 \cdot 2^{3m+3} & \text{if } l = 2 \\ 11 \cdot 2^{2k+m} + 21 \cdot 2^{k+3m+2} + 261 \cdot 2^{k+2m+1} + 315 \cdot 2^{4m+4} - 1627 \cdot 2^{3m+3} & \text{if } l = 1 \\ 21 \cdot 2^{2k+m} + 21 \cdot 2^{k+3m+2} + 525 \cdot 2^{k+2m+1} + 315 \cdot 2^{4m+4} - 3255 \cdot 2^{3m+3} & \text{if } l = 0 \end{cases}$$

Proof. The case $l \geq 3$

From (1.4) we get with $i \rightarrow m+3$

$$\Gamma_{m+3}^{\left[\begin{smallmatrix} 1 \\ 1+m+l \end{smallmatrix} \right] \times k} = (2^k - 2^{m+2}) \cdot \Gamma_{m+2}^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} + 2^{m+3} \cdot \Gamma_{m+3}^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k}$$

Applying Theorem 3.13 in [1, see section 3] with $s \rightarrow 1+m$, $m \rightarrow l$ we obtain

$$\begin{aligned}
\Gamma_{m+3}^{\left[\begin{smallmatrix} 1 \\ 1+m+l \end{smallmatrix} \right] \times k} &= (2^k - 2^{m+2}) \cdot [3 \cdot 2^{k+m} + 21 \cdot (2^{3m+2} - 2^{2m+1})] + 2^{m+3} \cdot [3 \cdot 2^{k+m+2} + 21 \cdot (2^{3m+5} - 2^{2m+4})] \\
&= 3 \cdot 2^{2k+m} + 21 \cdot 2^{k+3m+2} + 21 \cdot 2^{k+2m+1} + 315 \cdot 2^{4m+4} - 315 \cdot 2^{3m+3}
\end{aligned}$$

The other cases are proved in a similar way by applying results proved in [2, see section 3]

□

Lemma 1.18. *We have for $m \geq 2$ and $l \geq 3$ and $k \geq 2m+l$*

$$\Gamma_i^{\left[\begin{smallmatrix} 1 \\ 1+m+l \end{smallmatrix} \right] \times k} \text{ is equal to}$$

$$\begin{aligned}
& \begin{cases} 1 & \text{if } i = 0 \\ 2^k + 17 & \text{if } i = 1 \\ (21 \cdot 2^{3i-7} - 3 \cdot 2^{2i-5}) \cdot 2^k + 315 \cdot 2^{4i-8} - 21 \cdot 2^{3i-6} & \text{if } 2 \leq i \leq m \\ 21 \cdot 2^{k+3m-4} + 13 \cdot 2^{k+2m-3} + 315 \cdot 2^{4m-4} - 85 \cdot 2^{3m-3} & \text{if } i = m+1 \\ 2^{2k+m} + 21 \cdot 2^{k+3m-1} + 9 \cdot 2^{k+2m-1} + 315 \cdot 2^{4m} - 157 \cdot 2^{3m} & \text{if } i = m+2 \\ 3 \cdot 2^{2k-m+2i-6} + 21 \cdot 2^{k+3i-7} + 21 \cdot 2^{k-m+3i-8} + 315 \cdot 2^{4i-8} - 315 \cdot 2^{4i-m-9} & \text{if } m+3 \leq i \leq m+l \\ 3 \cdot 2^{2k+m+2l-4} + 21 \cdot 2^{k+3m+3l-4} + 149 \cdot 2^{k+2m+3l-5} + 315 \cdot 2^{4m+4l-4} - 827 \cdot 2^{3m+4l-5} & \text{if } i = m+l+1 \\ 11 \cdot 2^{2k+m+2l-2} + 21 \cdot 2^{k+3m+3l-1} + 261 \cdot 2^{k+2m+3l-2} + 315 \cdot 2^{4m+4l} - 1627 \cdot 2^{3m+4l-1} & \text{if } i = m+l+2 \\ 21 \cdot 2^{2k-2m-l+3i-9} + 21 \cdot 2^{k+3i-7} + 525 \cdot 2^{k-2m-l+4i-11} + 315 \cdot 2^{4i-8} - 3255 \cdot 2^{5i-2m-l-12} & \text{if } m+l+3 \leq i \leq 2m+l+1 \\ 53 \cdot 2^{2k+4m+2l-3} - 159 \cdot 2^{k+6m+3l-3} + 53 \cdot 2^{8m+4l-2} & \text{if } i = 2m+l+2 \\ 2^{3k+2m+l} - 7 \cdot 2^{2k+4m+2l} + 7 \cdot 2^{k+6m+3l+1} - 2^{8m+4l+3} & \text{if } i = 2m+l+3 \end{cases} \\
& = \begin{cases} 1 & \text{if } i = 0 \\ 2^k + 17 & \text{if } i = 1 \\ (21 \cdot 2^{3i-7} - 3 \cdot 2^{2i-5}) \cdot 2^k + 315 \cdot 2^{4i-8} - 21 \cdot 2^{3i-6} & \text{if } 2 \leq i \leq m \\ 21 \cdot 2^{k+3m-4} + 13 \cdot 2^{k+2m-3} + 315 \cdot 2^{4m-4} - 85 \cdot 2^{3m-3} & \text{if } i = m+1 \\ 2^{2k+m} + 21 \cdot 2^{k+3m-1} + 9 \cdot 2^{k+2m-1} + 315 \cdot 2^{4m} - 157 \cdot 2^{3m} & \text{if } i = m+2 \\ 3 \cdot 2^{2k-m+2i-6} + 21 \cdot 2^{k+3i-7} + 21 \cdot 2^{k-m+3i-8} + 315 \cdot 2^{4i-8} - 315 \cdot 2^{4i-m-9} & \text{if } m+3 \leq i \leq m+l \\ 3 \cdot 2^{2k+m+2l-4} + 21 \cdot 2^{k+3m+3l-4} + 149 \cdot 2^{k+2m+3l-5} + 315 \cdot 2^{4m+4l-4} - 827 \cdot 2^{3m+4l-5} & \text{if } i = m+l+1 \\ 11 \cdot 2^{2k+m+2l-2} + 21 \cdot 2^{k+3m+3l-1} + 261 \cdot 2^{k+2m+3l-2} + 315 \cdot 2^{4m+4l} - 1627 \cdot 2^{3m+4l-1} & \text{if } i = m+l+2 \\ 21 \cdot 2^{2k-2m-l+3i-9} + 21 \cdot 2^{k+3i-7} + 525 \cdot 2^{k-2m-l+4i-11} + 315 \cdot 2^{4i-8} - 3255 \cdot 2^{5i-2m-l-12} & \text{if } m+l+3 \leq i \leq 2m+l+1 \\ 53 \cdot 2^{2k+4m+2l-3} - 159 \cdot 2^{k+6m+3l-3} + 53 \cdot 2^{8m+4l-2} & \text{if } i = 2m+l+2 \\ 2^{3k+2m+l} - 7 \cdot 2^{2k+4m+2l} + 7 \cdot 2^{k+6m+3l+1} - 2^{8m+4l+3} & \text{if } i = 2m+l+3 \end{cases}
\end{aligned}$$

Proof. Using (1.4) and the results obtained in [2, see section 3] together with the

Lemmas 1.16 and 1.17 we can easily compute $\Gamma_i^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k}$ for $0 \leq i \leq 2m+l+3$ $\square$

Example. We have for $m = 2, l = 3, k \geq 7$

$$\Gamma_i^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} = \begin{cases} 1 & \text{if } i = 0 \\ 2^k + 17 & \text{if } i = 1 \\ 9 \cdot 2^k + 294 & \text{if } i = 2 \\ 55 \cdot 2^{k+1} + 4360 & \text{if } i = 3 \\ 2^{2k+2} + 93 \cdot 2^{k+3} + 70592 & \text{if } i = 4 \\ 3 \cdot 2^{2k+2} + 189 \cdot 2^{k+5} + 1128960 & \text{if } i = 5 \\ 3 \cdot 2^{2k+4} + 317 \cdot 2^{k+5} + 13869056 & \text{if } i = 6 \\ 11 \cdot 2^{2k+6} + 429 \cdot 2^{k+11} + 893 \cdot 2^{17} & \text{if } i = 7 \\ 21 \cdot 2^{2k+8} + 693 \cdot 2^{k+14} - 1541406720 & \text{if } i = 8 \\ 53 \cdot 2^{2k+11} - 159 \cdot 2^{k+18} + 53 \cdot 2^{26} & \text{if } i = 9 \\ 2^{3k+7} - 7 \cdot 2^{2k+14} + 7 \cdot 2^{k+22} - 2^{31} & \text{if } i = 10 \end{cases}$$

Lemma 1.19. We have for $m \geq 2, l = 2$ and $k \geq 2m+2$

$$\Gamma_i^{\left[\begin{smallmatrix} 1+m \\ 1+m+2 \end{smallmatrix} \right] \times k} = \begin{cases} 1 & \text{if } i = 0 \\ 2^k + 17 & \text{if } i = 1 \\ (21 \cdot 2^{3i-7} - 3 \cdot 2^{2i-5}) \cdot 2^k + 315 \cdot 2^{4i-8} - 21 \cdot 2^{3i-6} & \text{if } 2 \leq i \leq m \\ 21 \cdot 2^{k+3m-4} + 13 \cdot 2^{k+2m-3} + 315 \cdot 2^{4m-4} - 85 \cdot 2^{3m-3} & \text{if } i = m+1 \\ 2^{2k+m} + 21 \cdot 2^{k+3m-1} + 9 \cdot 2^{k+2m-1} + 315 \cdot 2^{4m} - 157 \cdot 2^{3m} & \text{if } i = m+2 \\ 3 \cdot 2^{2k+m} + 21 \cdot 2^{k+3m+2} + 149 \cdot 2^{k+2m+1} + 315 \cdot 2^{4m+4} - 827 \cdot 2^{3m+3} & \text{if } i = m+3 \\ 11 \cdot 2^{2k+m+2} + 21 \cdot 2^{k+3m+5} + 261 \cdot 2^{k+2m+4} + 315 \cdot 2^{4m+8} - 1627 \cdot 2^{3m+7} & \text{if } i = m+4 \\ 21 \cdot 2^{2k-2m+3i-11} + 21 \cdot 2^{k+3i-7} + 525 \cdot 2^{k-2m+4i-13} + 315 \cdot 2^{4i-8} - 3255 \cdot 2^{5i-2m-14} & \text{if } m+5 \leq i \leq 2m+3 \\ 53 \cdot 2^{2k+4m+1} - 159 \cdot 2^{k+6m+3} + 53 \cdot 2^{8m+6} & \text{if } i = 2m+4 \\ 2^{3k+2m+2} - 7 \cdot 2^{2k+4m+4} + 7 \cdot 2^{k+6m+7} - 2^{8m+11} & \text{if } i = 2m+5 \end{cases}$$

Example. We have for $m = 2, l = 2, k \geq 6$

$$\Gamma_i^{\left[\begin{smallmatrix} 1 \\ 1+2 \\ 1+2+2 \end{smallmatrix} \right] \times k} = \begin{cases} 1 & \text{if } i = 0 \\ 2^k + 17 & \text{if } i = 1 \\ 9 \cdot 2^k + 294 & \text{if } i = 2 \\ 55 \cdot 2^{k+1} + 4360 & \text{if } i = 3 \\ 2^{2k+2} + 93 \cdot 2^{k+3} + 70592 & \text{if } i = 4 \\ 3 \cdot 2^{2k+2} + 317 \cdot 2^{k+5} + 866816 & \text{if } i = 5 \\ 11 \cdot 2^{2k+4} + 429 \cdot 2^{k+8} + 7315456 & \text{if } i = 6 \\ 21 \cdot 2^{2k+6} + 693 \cdot 2^{k+11} - 735 \cdot 2^{17} & \text{if } i = 7 \\ 53 \cdot 2^{2k+9} - 159 \cdot 2^{k+15} + 53 \cdot 2^{22} & \text{if } i = 8 \\ 2^{3k+4} - 7 \cdot 2^{2k+12} + 7 \cdot 2^{k+19} - 2^{27} & \text{if } i = 9 \end{cases}$$

Lemma 1.20. We have for $m \geq 2$ and $l = 1$ and $k \geq 2m + 1$

$\Gamma_i^{\left[\begin{smallmatrix} 1 \\ 1+m \\ 1+m+1 \end{smallmatrix} \right] \times k}$ is equal to

$$= \begin{cases} 1 & \text{if } i = 0 \\ 2^k + 17 & \text{if } i = 1 \\ (21 \cdot 2^{3i-7} - 3 \cdot 2^{2i-5}) \cdot 2^k + 315 \cdot 2^{4i-8} - 21 \cdot 2^{3i-6} & \text{if } 2 \leq i \leq m \\ 21 \cdot 2^{k+3m-4} + 13 \cdot 2^{k+2m-3} + 315 \cdot 2^{4m-4} - 85 \cdot 2^{3m-3} & \text{if } i = m + 1 \\ 2^{2k+m} + 21 \cdot 2^{k+3m-1} + 73 \cdot 2^{k+2m-1} + 315 \cdot 2^{4m} - 413 \cdot 2^{3m} & \text{if } i = m + 2 \\ 11 \cdot 2^{2k+m} + 21 \cdot 2^{k+3m+2} + 261 \cdot 2^{k+2m+1} + 315 \cdot 2^{4m+4} - 1627 \cdot 2^{3m+3} & \text{if } i = m + 3 \\ 21 \cdot 2^{2k-2m+3i-10} + 21 \cdot 2^{k+3i-7} + 525 \cdot 2^{k-2m+4i-12} + 315 \cdot 2^{4i-8} - 3255 \cdot 2^{5i-2m-13} & \text{if } m + 4 \leq i \leq 2m + 2 \\ 53 \cdot 2^{2k+4m-1} - 159 \cdot 2^{k+6m} + 53 \cdot 2^{8m+2} & \text{if } i = 2m + 3 \\ 2^{3k+2m+1} - 7 \cdot 2^{2k+4m+2} + 7 \cdot 2^{k+6m+4} - 2^{8m+7} & \text{if } i = 2m + 4 \end{cases}$$

Example. We have for $m = 2$, $l = 1$, $k \geq 5$

$$\Gamma_i^{\left[\begin{smallmatrix} 1 \\ 1+2 \\ 1+2+1 \end{smallmatrix} \right] \times k} = \begin{cases} 1 & \text{if } i = 0 \\ 2^k + 17 & \text{if } i = 1 \\ 9 \cdot 2^k + 294 & \text{if } i = 2 \\ 55 \cdot 2^{k+1} + 4360 & \text{if } i = 3 \\ 2^{2k+2} + 1256 \cdot 2^k + 54208 & \text{if } i = 4 \\ 11 \cdot 2^{2k+2} + 13728 \cdot 2^k + 457216 & \text{if } i = 5 \\ 21 \cdot 2^{2k+4} + 693 \cdot 2^{k+8} - 735 \cdot 2^{13} & \text{if } i = 6 \\ 53 \cdot 2^{2k+7} - 159 \cdot 2^{k+12} + 53 \cdot 2^{18} & \text{if } i = 7 \\ 2^{3k+5} - 7 \cdot 2^{2k+10} + 7 \cdot 2^{k+16} - 2^{23} & \text{if } i = 8 \end{cases}$$

Lemma 1.21. We have for $m \geq 2$ and $l = 0$ and $k \geq 2m$

$\Gamma_i^{\left[\begin{smallmatrix} 1 \\ 1+m \\ 1+m \end{smallmatrix} \right] \times k}$ is equal to

$$= \begin{cases} 1 & \text{if } i = 0 \\ 2^k + 17 & \text{if } i = 1 \\ (21 \cdot 2^{3i-7} - 3 \cdot 2^{2i-5}) \cdot 2^k + 315 \cdot 2^{4i-8} - 21 \cdot 2^{3i-6} & \text{if } 2 \leq i \leq m \\ 21 \cdot 2^{k+3m-4} + 45 \cdot 2^{k+2m-3} + 315 \cdot 2^{4m-4} - 213 \cdot 2^{3m-3} & \text{if } i = m+1 \\ 3 \cdot 2^{2k+m} + 21 \cdot 2^{k+3m-1} + 129 \cdot 2^{k+2m-1} + 315 \cdot 2^{4m} - 813 \cdot 2^{3m} & \text{if } i = m+2 \\ 21 \cdot 2^{2k-2m+3i-9} + 21 \cdot 2^{k+3i-7} + 525 \cdot 2^{k-2m+4i-11} + 315 \cdot 2^{4i-8} - 3255 \cdot 2^{5i-2m-12} & \text{if } m+3 \leq i \leq 2m+1 \\ 53 \cdot 2^{2k+4m-3} - 159 \cdot 2^{k+6m-3} + 53 \cdot 2^{8m-2} & \text{if } i = 2m+2 \\ 2^{3k+2m} - 7 \cdot 2^{2k+4m} + 7 \cdot 2^{k+6m+1} - 2^{8m+3} & \text{if } i = 2m+3 \end{cases}$$

Example. We have for $m = 2$, $l = 0$, $k \geq 4$

$$\Gamma_i^{\begin{bmatrix} 1 & 1 \\ 1+2 & 1+2 \end{bmatrix} \times k} = \begin{cases} 1 & \text{if } i = 0 \\ 2^k + 17 & \text{if } i = 1 \\ 9 \cdot 2^k + 294 & \text{if } i = 2 \\ 87 \cdot 2^{k+1} + 3336 & \text{if } i = 3 \\ 3 \cdot 2^{2k+2} + 213 \cdot 2^{k+3} + 28608 & \text{if } i = 4 \\ 21 \cdot 2^{2k+2} + 693 \cdot 2^{k+5} - 735 \cdot 2^9 & \text{if } i = 5 \\ 53 \cdot 2^{2k+5} - 159 \cdot 2^{k+9} + 53 \cdot 2^{14} & \text{if } i = 6 \\ 2^{3k+4} - 7 \cdot 2^{2k+8} + 7 \cdot 2^{k+13} - 2^{19} & \text{if } i = 7 \end{cases}$$

Lemma 1.22. We have for $m = 1$ and $l \geq 3$ and $k \geq 2 + l$

$\Gamma_i^{\begin{bmatrix} 1 & 1 \\ 1+1 & 1+1+l \end{bmatrix} \times k}$ is equal to

$$= \begin{cases} 1 & \text{if } i = 0 \\ 2^k + 17 & \text{if } i = 1 \\ 17 \cdot 2^k + 230 & \text{if } i = 2 \\ 2^{2k+1} + 51 \cdot 2^{k+1} + 3784 & \text{if } i = 3 \\ 3 \cdot 2^{2k+2i-7} + 105 \cdot 2^{k+3i-9} + 945 \cdot 2^{4i-10} & \text{if } 4 \leq i \leq l+1 \\ 3 \cdot 2^{2k+2l-3} + 233 \cdot 2^{k+3l-3} + 433 \cdot 2^{4l-2} & \text{if } i = l+2 \\ 11 \cdot 2^{2k+2l-1} + 345 \cdot 2^{k+3l} - 367 \cdot 2^{4l+2} & \text{if } i = l+3 \\ 53 \cdot 2^{2k+2l+1} - 159 \cdot 2^{k+3l+3} + 3392 \cdot 2^{4l} & \text{if } i = l+4 \\ 2^{3k+l+2} - 7 \cdot 2^{2k+2l+4} + 7 \cdot 2^{k+3l+7} - 2^{4l+11} & \text{if } i = l+5 \end{cases}$$

$$\Gamma_i^{\begin{bmatrix} 2 \\ 5 \end{bmatrix} \times k} = \begin{cases} 1 & \text{if } i = 0 \\ 9 & \text{if } i = 1 \\ 2^{k+1} + 62 & \text{if } i = 2 \\ 3 \cdot 2^{k+1} + 504 & \text{if } i = 3 \\ 3 \cdot 2^{k+3} + 4032 & \text{if } i = 4 \\ 11 \cdot 2^{k+5} + 15872 & \text{if } i = 5 \\ 21 \cdot 2^{k+7} - 86016 & \text{if } i = 6 \\ 2^{2k+5} - 3 \cdot 2^{k+10} + 2^{16} & \text{if } i = 7 \end{cases}$$

$$\Gamma_i \begin{bmatrix} 1 \\ 1+1 \\ 1+1+3 \end{bmatrix} \times k = \begin{cases} 1 & \text{if } i = 0 \\ 2^k + 17 & \text{if } i = 1 \\ 17 \cdot 2^k + 230 & \text{if } i = 2 \\ 2^{2k+1} + 51 \cdot 2^{k+1} + 3784 & \text{if } i = 3 \\ 3 \cdot 2^{2k+1} + 105 \cdot 2^{k+3} + 60480 & \text{if } i = 4 \\ 3 \cdot 2^{2k+3} + 233 \cdot 2^{k+6} + 433 \cdot 2^{10} & \text{if } i = 5 \\ 11 \cdot 2^{2k+5} + 345 \cdot 2^{k+9} - 367 \cdot 2^{14} & \text{if } i = 6 \\ 53 \cdot 2^{2k+7} - 651264 \cdot 2^k + 13893632 & \text{if } i = 7 \\ 2^{3k+5} - 7 \cdot 2^{2k+10} + 7 \cdot 2^{k+16} - 2^{23} & \text{if } i = 8 \end{cases}$$

$$\Gamma_i \begin{bmatrix} 1 \\ 1+1 \\ 1+1+2 \end{bmatrix} \times k = \begin{cases} 1 & \text{if } i = 0 \\ 2^k + 17 & \text{if } i = 1 \\ 17 \cdot 2^k + 230 & \text{if } i = 2 \\ 2^{2k+1} + 51 \cdot 2^{k+1} + 3784 & \text{if } i = 3 \\ 3 \cdot 2^{2k+1} + 233 \cdot 2^{k+3} + 433 \cdot 2^6 & \text{if } i = 4 \\ 11 \cdot 2^{2k+3} + 345 \cdot 2^{k+6} - 367 \cdot 2^{10} & \text{if } i = 5 \\ 53 \cdot 2^{2k+5} - 159 \cdot 2^{k+9} + 3392 \cdot 2^8 & \text{if } i = 6 \\ 2^{3k+4} - 7 \cdot 2^{2k+8} + 7 \cdot 2^{k+13} - 2^{19} & \text{if } i = 7 \end{cases}$$

$$\Gamma_i \begin{bmatrix} 1 \\ 1+1 \\ 1+1+1 \end{bmatrix} \times k = \begin{cases} 1 & \text{if } i = 0 \\ 2^k + 17 & \text{if } i = 1 \\ 17 \cdot 2^k + 230 & \text{if } i = 2 \\ 2^{2k+1} + 115 \cdot 2^{k+1} + 1736 & \text{if } i = 3 \\ 11 \cdot 2^{2k+1} + 345 \cdot 2^{k+3} - 367 \cdot 2^6 & \text{if } i = 4 \\ 53 \cdot 2^{2k+3} - 159 \cdot 2^{k+6} + 3392 \cdot 2^4 & \text{if } i = 5 \\ 2^{3k+3} - 7 \cdot 2^{2k+6} + 7 \cdot 2^{k+10} - 2^{15} & \text{if } i = 6 \end{cases}$$

$$\Gamma_i \begin{bmatrix} 1 \\ 1+1 \\ 1+1 \end{bmatrix} \times k = \begin{cases} 1 & \text{if } i = 0 \\ 2^k + 17 & \text{if } i = 1 \\ 33 \cdot 2^k + 102 & \text{if } i = 2 \\ 3 \cdot 2^{2k+1} + 171 \cdot 2^{k+1} - 1464 & \text{if } i = 3 \\ 53 \cdot 2^{2k+1} - 159 \cdot 2^{k+3} + 3392 & \text{if } i = 4 \\ 2^{3k+2} - 7 \cdot 2^{2k+4} + 7 \cdot 2^{k+7} - 2^{11} & \text{if } i = 5 \end{cases}$$

Lemma 1.23. The case $m = 0$, $k \geq 3$

$$\Gamma_i \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \times k = \begin{cases} 1 & \text{if } i = 0 \\ 7(2^k - 1) & \text{if } i = 1 \\ 7 \cdot 2^{2k} - 21 \cdot 2^k + 14 & \text{if } i = 2 \\ 2^{3k} - 7 \cdot 2^{2k} + 7 \cdot 2^{k+1} - 8 & \text{if } i = 3 \end{cases}$$

The case $m = 1$, $k \geq 4$

$$\Gamma_i \begin{bmatrix} 1 \\ 1+1 \end{bmatrix} \times k = \begin{cases} 1 & \text{if } i = 0 \\ 3 \cdot 2^k + 9 & \text{if } i = 1 \\ 2^{2k} + 47 \cdot 2^k - 54 & \text{if } i = 2 \\ 27 \cdot 2^{2k} - 151 \cdot 2^k + 172 & \text{if } i = 3 \\ 2^{3k+1} - 7 \cdot 2^{2k+2} + 7 \cdot 2^{k+4} - 128 & \text{if } i = 4 \end{cases}$$

The case $m = 2, k \geq 5$

$$\Gamma_i \begin{bmatrix} 1 \\ 1+2 \end{bmatrix} \times k = \begin{cases} 1 & \text{if } i = 0 \\ 3 \cdot 2^k + 9 & \text{if } i = 1 \\ 2^{2k} + 15 \cdot 2^k + 158 & \text{if } i = 2 \\ 3 \cdot 2^{2k} + 191 \cdot 2^{k+1} - 1576 & \text{if } i = 3 \\ 27 \cdot 2^{2k} - 81 \cdot 2^{k+4} + 3456 & \text{if } i = 4 \\ 2^{3k+2} - 7 \cdot 2^{2k+4} + 7 \cdot 2^{k+7} - 2048 & \text{if } i = 5 \end{cases}$$

The case $4 \leq k \leq 2 + m$

$$\Gamma_i \begin{bmatrix} 1 \\ 1+m \end{bmatrix} \times k = \begin{cases} 1 & \text{if } i = 0 \\ 3 \cdot 2^k + 9 & \text{if } i = 1 \\ 2^{2k} + 15 \cdot 2^k + 158 & \text{if } i = 2 \\ 3 \cdot 2^{2k+2i-6} + 63 \cdot [2^{k+3i-8} + 5 \cdot 2^{4i-9}] & \text{if } 3 \leq i \leq k-1 \\ 2^{3k+m} - 47 \cdot 2^{4k-9} & \text{if } i = k \end{cases}$$

The case $3 \leq m \leq k-3$

$$\Gamma_i \begin{bmatrix} 1 \\ 1+m \end{bmatrix} \times k = \begin{cases} 1 & \text{if } i = 0 \\ 3 \cdot 2^k + 9 & \text{if } i = 1 \\ 2^{2k} + 15 \cdot 2^k + 158 & \text{if } i = 2 \\ 3 \cdot 2^{2k+2i-6} + 63 \cdot [2^{k+3i-8} + 5 \cdot 2^{4i-9}] & \text{if } 3 \leq i \leq m \\ 3 \cdot 2^{2k+2m-4} + 191 \cdot 2^{k+3m-5} - 197 \cdot 2^{4m-5} & \text{if } i = m+1 \\ 27 \cdot 2^{2k+2m-2} - 81 \cdot 2^{k+3m-2} + 27 \cdot 2^{4m-1} & \text{if } i = m+2 \\ 2^{3k+m} - 7 \cdot 2^{2k+2m} + 7 \cdot 2^{k+3m+1} - 2^{4m+3} & \text{if } i = m+3 \end{cases}$$

Proof. The proofs of Lemma 1.19, Lemma 1.20, Lemma 1.21, Lemma 1.22 and Lemma 1.23 are similar to the proof of Lemma 1.18. $\square$

Lemma 1.24. Let $\Gamma_i \begin{bmatrix} (n) \\ 1+m+l \end{bmatrix} \times k$ denote the number of matrices of the form $\begin{bmatrix} A \\ C \end{bmatrix}$ of rank i such that A is a $n \times k$ matrix over $\mathbb{F}_2$, B a $(1+m) \times k$ persymmetric matrix and C a $(1+m+l) \times k$ persymmetric matrix and where $\Gamma_i \begin{bmatrix} 1+m \\ 1+m+l \end{bmatrix} \times k$ denotes the number of double persymmetric $(2+2m+l) \times k$ rank i matrices of the form $\begin{bmatrix} B \\ C \end{bmatrix}$

Then $\Gamma_i \begin{bmatrix} (n) \\ 1+m+l \end{bmatrix} \times k$ expressed as a linear combination of the $\Gamma_{i-j} \begin{bmatrix} 1+m \\ 1+m+l \end{bmatrix} \times k$ is equal to

$$\sum_{j=0}^n 2^{(n-j) \cdot (i-j)} a_j^{(n)} \prod_{l=1}^j (2^k - 2^{i-l}) \cdot \Gamma_{i-j} \begin{bmatrix} 1+m \\ 1+m+l \end{bmatrix} \times k \quad \text{for } 0 \leq i \leq \inf(k, n+2m+l+2),$$

where $a_j^{(n)}$ satisfies the linear recurrence relation

$$(1.6) \quad a_j^{(n)} = 2^j \cdot a_j^{(n-1)} + a_{j-1}^{(n-1)}, \quad n = 2, 3, 4, \dots \quad \text{for } 1 \leq j \leq n-1.$$

The explicit value of $a_j^{(n)}$ is given by the following formula

$$(1.7) \quad a_j^{(n)} = (-1)^j \cdot 2^{jn - \frac{j(j-1)}{2}} + \sum_{s=0}^{j-1} (-1)^s \prod_{l=0}^{j-(s+1)} \frac{2^{n+1} - 2^l}{2^{j-s} - 2^l} \cdot 2^{s(n-j) + \frac{s(s+1)}{2}} \quad \text{for } 1 \leq j \leq n-1.$$

We set

$$a_0^{(n)} = a_n^{(n)} = 1$$

$$\text{and } \Gamma_{i-j}^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} = 0 \quad \text{if } i-j \notin \{0, 1, 2, \dots, \inf(k, 2m+l+2)\}.$$

Proof. The proof of Lemma 1.23 is somewhat similar to the proof of Lemma 11.1 in [1], indeed a analogous proof of Lemma 1.15 gives the following generalization

$$(1.8) \quad \Gamma_i^{\left[\begin{smallmatrix} (n) \\ 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} = 2^i \cdot \Gamma_i^{\left[\begin{smallmatrix} (n-1) \\ 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} + (2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{\left[\begin{smallmatrix} (n-1) \\ 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} \quad \text{for } 0 \leq i \leq \inf(k, n+2m+l+2).$$

We prove (1.5) by induction on n .

From (1.8) we get respectively for $n=1$ and $n=2$

$$(1.9) \quad \Gamma_i^{\left[\begin{smallmatrix} 1 \\ 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} = (2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} + 2^i \cdot \Gamma_i^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} \quad \text{for } 0 \leq i \leq \inf(k, 2m+l+3)$$

$$(1.10) \quad \Gamma_i^{\left[\begin{smallmatrix} (2) \\ 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} = (2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} + 2^i \cdot \Gamma_i^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} \quad \text{for } 0 \leq i \leq \inf(k, 2m+l+4)$$

From (1.9) and (1.10) we deduce

$$(1.11) \quad \Gamma_i^{\left[\begin{smallmatrix} (2) \\ 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} = 2^{2i} \Gamma_i^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} + 3 \cdot 2^{i-1} (2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} \\ + (2^k - 2^{i-1}) (2^k - 2^{i-2}) \cdot \Gamma_{i-2}^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} \quad \text{for } 0 \leq i \leq \inf(k, 2m+l+4).$$

Hence by (1.9) and (1.11) the formula (1.5) holds for $n=1$ and $n=2$.

Assume now that (1.5) holds for the number $n-1$, that is

$$(1.12) \quad 2^i \cdot \Gamma_i^{\left[\begin{smallmatrix} (n-1) \\ 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} = \sum_{j=0}^{n-1} 2^{(n-1-j) \cdot (i-j)} 2^i a_j^{(n-1)} \prod_{l=1}^j (2^k - 2^{i-l}) \cdot \Gamma_{i-j}^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k}$$

$$(1.13) \quad (2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{\left[\begin{smallmatrix} (n-1) \\ 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} = (2^k - 2^{i-1}) \left[\sum_{j=0}^{n-1} 2^{(n-1-j) \cdot (i-1-j)} a_j^{(n-1)} \prod_{l=1}^j (2^k - 2^{i-1-l}) \cdot \Gamma_{i-1-j}^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} \right] \\ = \sum_{j=0}^{n-1} 2^{(n-1-j) \cdot (i-1-j)} a_j^{(n-1)} \prod_{l=1}^{j+1} (2^k - 2^{i-l}) \cdot \Gamma_{i-(j+1)}^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} \\ = \sum_{j=1}^n 2^{(n-j) \cdot (i-j)} a_{j-1}^{(n-1)} \prod_{l=1}^j (2^k - 2^{i-l}) \cdot \Gamma_{i-j}^{\left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k}.$$

From (1.12), (1.13) and (1.7) it follows

$$\begin{aligned}
\Gamma_i \left[\begin{smallmatrix} (n) \\ 1+m+l \end{smallmatrix} \right] \times k &= 2^{ni} \Gamma_i \left[\begin{smallmatrix} 1+m+l \\ 1+m+l \end{smallmatrix} \right] \times k + \sum_{j=1}^{n-1} 2^{(n-j) \cdot (i-j)} (2^j a_j^{(n-1)} + a_{j-1}^{(n-1)}) \prod_{l=1}^j (2^k - 2^{i-l}) \cdot \Gamma_{i-j} \left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k \\
&+ a_{n-1}^{(n-1)} \prod_{l=1}^n (2^k - 2^{i-l}) \Gamma_{i-n} \left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k \\
&= \sum_{j=0}^n 2^{(n-j) \cdot (i-j)} a_j^{(n)} \prod_{l=1}^j (2^k - 2^{i-l}) \cdot \Gamma_{i-j} \left[\begin{smallmatrix} 1+m \\ 1+m+l \end{smallmatrix} \right] \times k
\end{aligned}$$

which proves (1.5).

By formula (1.5) with $m = l = 0$, we obtain

$$(1.14) \quad \Gamma_i \left[\begin{smallmatrix} (n) \\ 1 \end{smallmatrix} \right] \times k = \sum_{j=0}^n 2^{(n-j) \cdot (i-j)} a_j^{(n)} \prod_{l=1}^j (2^k - 2^{i-l}) \cdot \Gamma_{i-j} \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right] \times k \quad \text{for } 0 \leq i \leq \inf(k, n+2)$$

We have obviously

$$(1.15) \quad \Gamma_{i-j} \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right] \times k = \begin{cases} 1 & \text{if } i-j = 0, \\ 3 \cdot (2^k - 1) & \text{if } i-j = 1, \\ 2^{2k} - 3 \cdot 2^k + 2 & \text{if } i-j = 2. \end{cases}$$

From (1.14) and (1.15) we deduce

$$\begin{aligned}
(1.16) \quad \Gamma_i \left[\begin{smallmatrix} (n) \\ 1 \end{smallmatrix} \right] \times k &= \sum_{j=i-2}^i 2^{(n-j) \cdot (i-j)} a_j^{(n)} \prod_{l=1}^j (2^k - 2^{i-l}) \cdot \Gamma_{i-j} \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right] \times k \\
&= a_{i-2}^{(n)} \cdot 2^{2n-2i+4} \prod_{l=1}^{i-2} (2^k - 2^{i-l}) \cdot (2^{2k} - 3 \cdot 2^k + 2) + a_{i-1}^{(n)} \cdot 2^{n-i+1} \cdot \prod_{l=1}^{i-1} (2^k - 2^{i-l}) \cdot 3 \cdot (2^k - 1) \\
&+ a_i^{(n)} \cdot \prod_{l=1}^i (2^k - 2^{i-l}) \\
&= [2^{2n-2i+4} \cdot a_{i-2}^{(n)} + 3 \cdot 2^{n-i+1} \cdot a_{i-1}^{(n)} + a_i^{(n)}] \cdot \prod_{l=1}^i (2^k - 2^{i-l}) \\
&= [2^{2n-2i+4} \cdot a_{i-2}^{(n)} + 3 \cdot 2^{n-i+1} \cdot a_{i-1}^{(n)} + a_i^{(n)}] \cdot \prod_{l=0}^{i-1} (2^k - 2^l)
\end{aligned}$$

On the other hand by George Landsberg [3] we have

$$(1.17) \quad \Gamma_i \left[\begin{smallmatrix} (n) \\ 1 \end{smallmatrix} \right] \times k = \prod_{l=0}^{i-1} \frac{(2^{n+2} - 2^l)(2^k - 2^l)}{(2^i - 2^l)}.$$

Hence by (1.16) and (1.17) we have the formula

$$(1.18) \quad 2^{2n-2i+4} \cdot a_{i-2}^{(n)} + 3 \cdot 2^{n-i+1} \cdot a_{i-1}^{(n)} + a_i^{(n)} = \prod_{l=0}^{i-1} \frac{2^{n+2} - 2^l}{2^i - 2^l}.$$

We can then from (1.18) compute successively $a_i^{(n)}$ $i = 1, 2, \dots$ for $n \geq i$
We get

$$a_1^{(n)} = 2^n - 1 \quad \text{for } n \geq 1$$

$$a_2^{(n)} = \frac{2^{2n-1} - 3 \cdot 2^{n-1} + 1}{3} \quad \text{for } n \geq 2$$

To establish that $a_j^{(n)}$ is given by the formula (1.8) we compute the sum

$$2^{2n-2i+4} \cdot a_{i-2}^{(n)} + 3 \cdot 2^{n-i+1} \cdot a_{i-1}^{(n)} + a_i^{(n)}.$$

We get:

$$\begin{aligned} & 2^{2n-2i+4} \cdot \left[(-1)^{i-2} \cdot 2^{(i-2)n - \frac{(i-2)(i-3)}{2}} + \sum_{s=0}^{i-3} (-1)^s \prod_{l=0}^{i-s-3} \frac{2^{n+1} - 2^l}{2^{i-s-2} - 2^l} \cdot 2^{s(n-i+2) + \frac{s(s+1)}{2}} \right] \\ & + 3 \cdot 2^{n-i+1} \cdot \left[(-1)^{i-1} \cdot 2^{(i-1)n - \frac{(i-1)(i-2)}{2}} + \sum_{s=0}^{i-2} (-1)^s \prod_{l=0}^{i-s-2} \frac{2^{n+1} - 2^l}{2^{i-s-1} - 2^l} \cdot 2^{s(n-i+1) + \frac{s(s+1)}{2}} \right] \\ & + (-1)^i \cdot 2^{in - \frac{i(i-1)}{2}} + \sum_{s=0}^{i-1} (-1)^s \prod_{l=0}^{i-(s+1)} \frac{2^{n+1} - 2^l}{2^{i-s} - 2^l} \cdot 2^{s(n-i) + \frac{s(s+1)}{2}} = \prod_{l=0}^{i-1} \frac{2^{n+2} - 2^l}{2^i - 2^l}. \end{aligned}$$

□

Lemma 1.25. Let $f_{m,l,k}(t, \eta, \xi_1, \xi_2, \dots, \xi_n)$ be the exponential sum in $\mathbb{P}^{n+2}$ defined by

$$(t, \eta, \xi_1, \xi_2, \dots, \xi_n) \in \mathbb{P}^{n+2} \longrightarrow \sum_{\deg Y \leq k-1} \sum_{\deg Z \leq 1+m} E(tYZ) \sum_{\deg U \leq 1+m+l} E(\eta YU) \sum_{\deg V_1 \leq 0} E(\xi_1 YV_1) \sum_{\deg V_2 \leq 0} E(\xi_2 YV_2) \dots \sum_{\deg V_n \leq 0} E(\xi_n YV_n).$$

Set

$$(t, \eta, \xi_1, \xi_2, \dots, \xi_n) = \left(\sum_{i \geq 1} \alpha_i T^{-i}, \sum_{i \geq 1} \beta_i T^{-i}, \sum_{i \geq 1} \gamma_{1i} T^{-i}, \dots, \sum_{i \geq 1} \gamma_{ni} T^{-i} \right) \in \mathbb{P}^{n+2}.$$

Then

$$f_{m,l,k}(t, \eta, \xi_1, \xi_2, \dots, \xi_n) = 2^{k+2m+l+n+4-r(D \left[\begin{smallmatrix} (n) \\ 1+m \\ 1+m+l \end{smallmatrix} \right] \times k} (\xi_1, \xi_2, \dots, \xi_n, t, \eta)$$

where

$$D \left[\begin{smallmatrix} (n) \\ 1+m \\ 1+m+l \end{smallmatrix} \right] \times k (\xi_1, \xi_2, \dots, \xi_n, t, \eta)$$

denotes the following $(2m + l + n + 2) \times k$ matrix

$$\begin{pmatrix} \gamma_{11} & \gamma_{12} & \gamma_{13} & \cdots & \gamma_{1,k-1} & \gamma_{1,k} \\ \gamma_{21} & \gamma_{22} & \gamma_{23} & \cdots & \gamma_{2,k-1} & \gamma_{2,k} \\ \vdots & \vdots & \vdots & \cdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \cdots & \vdots & \vdots \\ \hline \gamma_{n1} & \gamma_{n2} & \gamma_{n3} & \cdots & \gamma_{n,k-1} & \gamma_{n,k} \\ \hline \alpha_1 & \alpha_2 & \alpha_3 & \cdots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \cdots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \cdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \cdots & \vdots & \vdots \\ \hline \alpha_{1+m} & \alpha_{2+m} & \alpha_{3+m} & \cdots & \alpha_{k+m-1} & \alpha_{k+m} \\ \hline \beta_1 & \beta_2 & \beta_3 & \cdots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \beta_4 & \cdots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{1+m+l} & \beta_{2+m+l} & \beta_{3+m+l} & \cdots & \beta_{k+m+l-1} & \beta_{k+m+l} \end{pmatrix}$$

Then the number of solutions

$$(Y_1, Z_1, U_1, V_1^{(1)}, V_2^{(1)}, \dots, V_n^{(1)}, Y_2, Z_2, U_2, V_1^{(2)}, V_2^{(2)}, \dots, V_n^{(2)}, \dots, Y_q, Z_q, U_q, V_1^{(q)}, V_2^{(q)}, \dots, V_n^{(q)})$$

of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0 \\ Y_1 U_1 + Y_2 U_2 + \dots + Y_q U_q = 0 \\ Y_1 V_1^{(1)} + Y_2 V_1^{(2)} + \dots + Y_q V_1^{(q)} = 0 \\ Y_1 V_2^{(1)} + Y_2 V_2^{(2)} + \dots + Y_q V_2^{(q)} = 0 \\ \vdots \\ Y_1 V_n^{(1)} + Y_2 V_n^{(2)} + \dots + Y_q V_n^{(q)} = 0 \end{cases}$$

satisfying the degree conditions

$$\deg Y_i \leq k-1, \quad \deg Z_i \leq m+1, \quad \deg U_i \leq m+l+1, \quad \deg V_j^i \leq 0, \quad \text{for } 1 \leq j \leq n \quad 1 \leq i \leq q$$

is equal to the following integral over the unit interval in $\mathbb{K}^{n+2}$

$$\int_{\mathbb{P}^{n+2}} f_{m,l,k}^q(t, \eta, \xi_1, \xi_2, \dots, \xi_n) dt d\eta d\xi_1 d\xi_2 \dots d\xi_n.$$

Observing that $f_{m,l,k}^q(t, \eta, \xi_1, \xi_2, \dots, \xi_n)$ is constant on cosets of $\mathbb{P}_{k+m} \times \mathbb{P}_{k+m+l} \times \mathbb{P}_k^n$, the above integral is equal to

$$2^{q(k+2m+l+n+4)-(2m+l+k(n+2))} \sum_{i=0}^{inf(k,n+2m+l+2)} \Gamma_i \left[\begin{matrix} (n) \\ 1+m \\ 1+m+l \end{matrix} \right] \times k 2^{-iq}.$$

Proof. The proof of Lemma 1.24 is just a generalization of the proof of Corollary 11.5. (see [1]) $\square$

Example. $n=2, m=1, l=3, k \geq 5, q=3$

$$\Gamma_i^{\left[\begin{smallmatrix} (2) \\ 2 \\ 5 \end{smallmatrix} \right] \times k} = 2^{2i} \Gamma_i^{\left[\begin{smallmatrix} 2 \\ 5 \end{smallmatrix} \right] \times k} + 3 \cdot 2^{i-1} (2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{\left[\begin{smallmatrix} 2 \\ 5 \end{smallmatrix} \right] \times k} + (2^k - 2^{i-1})(2^k - 2^{i-2}) \cdot \Gamma_{i-2}^{\left[\begin{smallmatrix} 2 \\ 5 \end{smallmatrix} \right] \times k} \quad \text{for } 0 \leq i \leq \inf(k, 9)$$

$$\Gamma_i^{\left[\begin{smallmatrix} (2) \\ 1+1 \\ 1+1+3 \end{smallmatrix} \right] \times k} = \begin{cases} 1 & \text{if } i = 0 \\ 3 \cdot 2^k + 33 & \text{if } i = 1 \\ 2^{2k} + 83 \cdot 2^k + 886 & \text{if } i = 2 \\ 33 \cdot 2^{2k} + 978 \cdot 2^k + 29352 & \text{if } i = 3 \\ 2^{3k+1} + 182 \cdot 2^{2k} + 16408 \cdot 2^k + 937408 & \text{if } i = 4 \\ 3 \cdot 2^{3k+1} + 189 \cdot 2^{2k+3} + 8191 \cdot 2^{k+6} + 12911 \cdot 2^{10} & \text{if } i = 5 \\ 3 \cdot 2^{3k+3} + 36672 \cdot 2^{2k} + 11271168 \cdot 2^k - 399015936 & \text{if } i = 6 \\ 11 \cdot 2^{3k+5} + 1022464 \cdot 2^{2k} - 100679680 \cdot 2^k + 2163212288 & \text{if } i = 7 \\ 117 \cdot 2^{3k+7} - 3354624 \cdot 2^{2k} + 214695936 \cdot 2^k - 3925868544 & \text{if } i = 8 \\ 2^{4k+5} - 480 \cdot 2^{3k+5} + 2240 \cdot 2^{2k+10} - 1920 \cdot 2^{k+16} + 2^{31} & \text{if } i = 9 \end{cases}$$

If $k = 5$ we obtain

$$\Gamma_i^{\left[\begin{smallmatrix} (2) \\ 1+1 \\ 1+1+3 \end{smallmatrix} \right] \times 5} = \begin{cases} 1 & \text{if } i = 0 \\ 129 & \text{if } i = 1 \\ 4566 & \text{if } i = 2 \\ 94440 & \text{if } i = 3 \\ 1714368 & \text{if } i = 4 \\ 31740928 & \text{if } i = 5 \end{cases}$$

Then the number of solutions

$$(Y_1, Z_1, U_1, V_1^{(1)}, V_2^{(1)}, Y_2, Z_2, U_2, V_1^{(2)}, V_2^{(2)}, Y_3, Z_3, U_3, V_1^{(3)}, V_2^{(3)})$$

of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + Y_3 Z_3 = 0 \\ Y_1 U_1 + Y_2 U_2 + Y_3 U_3 = 0 \\ Y_1 V_1^{(1)} + Y_2 V_1^{(2)} + Y_3 V_1^{(3)} = 0 \\ Y_1 V_2^{(1)} + Y_2 V_2^{(2)} + Y_3 V_2^{(3)} = 0 \end{cases}$$

satisfying the degree conditions

$$\deg Y_i \leq 4, \quad \deg Z_i \leq 2, \quad \deg U_i \leq 5, \quad \deg V_j^i \leq 0, \quad \text{for } 1 \leq j \leq 2, 1 \leq i \leq 3$$

is equal to

$$\int_{\mathbb{P}^4} f_{1,3,5}^3(t, \eta, \xi_1, \xi_2) dt d\eta d\xi_1 d\xi_2 = 2^{23} \sum_{i=0}^5 \Gamma_i^{\left[\begin{smallmatrix} (2) \\ 2 \\ 5 \end{smallmatrix} \right] \times k} 2^{-3i} = 13281 \cdot 2^{20}.$$

where for $(t, \eta, \xi_1, \xi_2, \xi_2) = (\sum_{i \geq 1} \alpha_i T^{-i}, \sum_{i \geq 1} \beta_i T^{-i}, \sum_{i \geq 1} \gamma_{1i} T^{-i}, \sum_{i \geq 1} \gamma_{2i} T^{-i}) \in \mathbb{P}^4$.

$$\begin{aligned} f_{1,3,5}(t, \eta, \xi_1, \xi_2) &= \sum_{\deg Y \leq 4} \sum_{\deg Z \leq 2} E(tYZ) \sum_{\deg U \leq 5} E(\eta YU) \sum_{\deg V_1 \leq 0} E(\xi_1 YV_1) \sum_{\deg V_2 \leq 0} E(\xi_2 YV_2) \\ &= 2^{16-r(D \left[\begin{smallmatrix} (2) \\ 2 \\ 5 \end{smallmatrix} \right] \times k} (\xi_1, \xi_2, t, \eta)) \end{aligned}$$

and

$$D \left[\begin{smallmatrix} (2) \\ 2 \\ 5 \end{smallmatrix} \right] \times k (\xi_1, \xi_2, t, \eta) = \begin{pmatrix} \begin{array}{ccccc} \gamma_{11} & \gamma_{12} & \gamma_{13} & \gamma_{1,4} & \gamma_{1,5} \\ \gamma_{21} & \gamma_{22} & \gamma_{23} & \gamma_{2,4} & \gamma_{2,5} \\ \hline \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 \\ \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 \\ \hline \beta_1 & \beta_2 & \beta_3 & \beta_4 & \beta_5 \\ \beta_2 & \beta_3 & \beta_4 & \beta_5 & \beta_6 \\ \beta_3 & \beta_4 & \beta_5 & \beta_6 & \beta_7 \\ \beta_4 & \beta_5 & \beta_6 & \beta_7 & \beta_8 \\ \beta_5 & \beta_6 & \beta_7 & \beta_8 & \beta_9 \end{array} \end{pmatrix}$$

1.5. Proofs of Theorems 1.4,1.5,1.6,1.7 and 1.8.

1.5.1. *Proof of Theorem 1.4.* . Follows from Lemma 1.15

1.5.2. *Proof of Theorem 1.5.* . Follows from Lemma 1.16

1.5.3. *Proof of Theorem 1.6.* . Follows from Lemma 1.18

1.5.4. *Proof of Theorem 1.7.* . Follows from Lemma 1.23

1.5.5. *Proof of Theorem 1.8.* . Follows from Lemma 1.24

2. A recurrent formula for the number $\Gamma_i \left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right] \times k$ of rank i matrices of the form $\left[\frac{A}{B} \right]$ such that A is a $s \times k$ persymmetric matrix over $\mathbb{F}_2$, B a $(s+m) \times k$ persymmetric matrix and C a $(s+m+l) \times k$ persymmetric matrix

2.1. Notation.

Definition 2.1. We introduce the following definitions in the three - dimensional $\mathbb{K}$ -vectorspace.

- Let k,s,m and l denote rational integers such that $k \geq 1$, $s \geq 2$ and $m \geq 0$, $l \geq 0$
- We denote by $\mathbb{P}/\mathbb{P}_i \times \mathbb{P}/\mathbb{P}_j \times \mathbb{P}/\mathbb{P}_r$ a complete set of coset representatives of $\mathbb{P}_i \times \mathbb{P}_j \times \mathbb{P}_r$ in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$, for instance $\mathbb{P}/\mathbb{P}_{s+k-1} \times \mathbb{P}/\mathbb{P}_{s+m+k-1} \times \mathbb{P}/\mathbb{P}_{s+m+l+k-1}$ denotes a complete set of coset representatives of $\mathbb{P}_{s+k-1} \times \mathbb{P}_{s+m+k-1} \times \mathbb{P}_{s+m+l+k-1}$ in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$.
- Set $(t, \eta, \xi) = (\sum_{i \geq 1} \alpha_i T^{-i}, \sum_{i \geq 1} \beta_i T^{-i}, \sum_{i \geq 1} \gamma_i T^{-i}) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P}$.

- We denote by $D \begin{bmatrix} s \\ s+m+l \end{bmatrix} \times k (t, \eta, \xi)$ any $(2s+m+l) \times k$ matrix, such that after a rearrangement of the rows, if necessary, we can obtain the following triple persymmetric matrix $\begin{bmatrix} D_{s \times k}(t) \\ \frac{D_{(s+m) \times k}(\eta)}{D_{(s+m+l) \times k}(\xi)} \end{bmatrix}$

$$\left(\begin{array}{cccccc} \alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \dots & \alpha_{s+k-2} & \alpha_{s+k-1} \\ \hline \beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \dots & \beta_{k+m-1} & \beta_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \\ \hline \gamma_1 & \gamma_2 & \gamma_3 & \dots & \gamma_{k-1} & \gamma_k \\ \gamma_2 & \gamma_3 & \gamma_4 & \dots & \gamma_k & \gamma_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{m+l+1} & \gamma_{m+l+2} & \gamma_{m+l+3} & \dots & \gamma_{k+m+l-1} & \beta_{k+m+l} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{s+m+l-1} & \gamma_{s+m+l} & \gamma_{s+m+l+1} & \dots & \gamma_{s+m+l+k-3} & \gamma_{s+m+l+k-2} \\ \gamma_{s+m+l} & \gamma_{s+m+l+1} & \gamma_{s+m+l+2} & \dots & \gamma_{s+m+l+k-2} & \gamma_{s+m+l+k-1} \end{array} \right)$$

- We denote by $D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-1} + \beta_{s+m-1} \end{bmatrix} \times k (t, \eta, \xi)$ any $(3s+2m+l-2) \times k$ matrix, such that after a rearrangement of the rows, if necessary, we can obtain a matrix where the first $(3s+2m+l-3)$ rows form the following triple persymmetric matrix $\begin{bmatrix} D_{s-1 \times k}(t) \\ \frac{D_{(s+m-1) \times k}(\eta)}{D_{(s+m+l-1) \times k}(\xi)} \end{bmatrix}$ and the last row is equal to $(\alpha_s + \beta_{s+m}, \alpha_{s+1} + \beta_{s+m+1}, \dots, \alpha_{s+k-1} + \beta_{s+m+k-1})$

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\hline
\beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \dots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
\gamma_1 & \gamma_2 & \gamma_3 & \dots & \gamma_{k-1} & \gamma_k \\
\gamma_2 & \gamma_3 & \gamma_4 & \dots & \gamma_k & \gamma_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\gamma_{m+l+1} & \gamma_{m+l+2} & \gamma_{m+l+3} & \dots & \gamma_{k+m+l-1} & \gamma_{k+m+l} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\gamma_{s+m+l-1} & \gamma_{s+m+l} & \gamma_{s+m+l+1} & \dots & \gamma_{s+m+l+k-3} & \gamma_{s+m+l+k-2} \\
\hline
\alpha_s + \beta_{s+m} & \alpha_{s+1} + \beta_{s+m+1} & \alpha_{s+2} + \beta_{s+m+2} & \dots & \alpha_{s+k-2} + \beta_{s+m+k-2} & \alpha_{s+k-1} + \beta_{s+m+k-1}
\end{pmatrix}.$$

- We denote by $D \begin{bmatrix} \frac{s-1}{s+m+l-1} \\ \frac{s+m+l-1}{\alpha_s + \gamma_{s+m+l-1}} \end{bmatrix} \times k$ (t, η, ξ) any $(3s + 2m + l - 2) \times k$ matrix, such that after a rearrangement of the rows, if necessary, we can obtain a matrix where the first $(3s + 2m + l - 3)$ rows form the following tripersymmetric matrix

$$\begin{bmatrix} \frac{D_{s-1 \times k}(t)}{D_{(s+m-1) \times k}(\eta)} \\ \frac{D_{(s+m+l-1) \times k}(\xi)}{D_{(s+m+l-1) \times k}(\xi)} \end{bmatrix} \text{ and the last row is equal to } (\alpha_s + \gamma_{s+m+l}, \alpha_{s+1} + \gamma_{s+m+l+1}, \dots, \alpha_{s+k-1} + \gamma_{s+m+l+k-1})$$

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\hline
\beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \dots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
\gamma_1 & \gamma_2 & \gamma_3 & \dots & \gamma_{k-1} & \gamma_k \\
\gamma_2 & \gamma_3 & \gamma_4 & \dots & \gamma_k & \gamma_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\gamma_{m+l+1} & \gamma_{m+l+2} & \gamma_{m+l+3} & \dots & \gamma_{k+m+l-1} & \gamma_{k+m+l} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\gamma_{s+m+l-1} & \gamma_{s+m+l} & \gamma_{s+m+l+1} & \dots & \gamma_{s+m+l+k-3} & \gamma_{s+m+l+k-2} \\
\hline
\alpha_s + \gamma_{s+m+l} & \alpha_{s+1} + \gamma_{s+m+l+1} & \alpha_{s+2} + \gamma_{s+m+l+2} & \dots & \alpha_{s+k-2} + \gamma_{s+m+l+k-2} & \alpha_{s+k-1} + \gamma_{s+m+l+k-1}
\end{pmatrix}.$$

We denote by $D \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_s + \beta_{s+m} + \gamma_{s+m+l} \end{smallmatrix} \right] \times k$ (t, η, ξ) any $(3s+2m+l-2) \times k$ matrix, such that after a rearrangement of the rows, if necessary, we can obtain a matrix where the first $(3s+2m+l-3)$ rows form the following triple persymmetric matrix

$$\left[\begin{array}{c} \frac{D_{s-1 \times k}(t)}{D_{(s+m-1) \times k}(\eta)} \\ \frac{D_{(s+m+l-1) \times k}(\xi)}{D_{(s+m+l-1) \times k}(\xi)} \end{array} \right] \text{ and the last row is equal to }$$

$$(\alpha_s + \beta_{s+m} + \gamma_{s+m+l}, \alpha_{s+1} + \beta_{s+m+1} + \gamma_{s+m+l+1}, \dots, \alpha_{s+k-1} + \beta_{s+m+k-1} + \gamma_{s+m+l+k-1})$$

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \dots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \dots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\hline
\beta_1 & \beta_2 & \dots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \dots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \dots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
\gamma_1 & \gamma_2 & \dots & \gamma_{k-1} & \gamma_k \\
\gamma_2 & \gamma_3 & \dots & \gamma_k & \gamma_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\gamma_{m+l+1} & \gamma_{m+l+2} & \dots & \gamma_{k+m+l-1} & \gamma_{k+m+l} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\gamma_{s+m+l-1} & \gamma_{s+m+l} & \dots & \gamma_{s+m+l+k-3} & \gamma_{s+m+l+k-2} \\
\hline
\alpha_s + \beta_{s+m} + \gamma_{s+m+l} & \alpha_{s+1} + \beta_{s+m+1} + \gamma_{s+m+l+1} & \dots & \alpha_{s+k-2} + \beta_{s+m+k-2} + \gamma_{s+m+l+k-2} & \alpha_{s+k-1} + \beta_{s+m+k-1} + \gamma_{s+m+l+k-1}
\end{pmatrix}$$

We denote by $D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_s - \\ \beta_{s+m} - \end{bmatrix} \times k$ (t, η, ξ) any $(3s+2m+l-1) \times k$ matrix, such that after a rearrangement of the rows, if necessary, we can obtain a matrix where the first $(3s+2m+l-3)$ rows form the following triple persymmetric matrix $\begin{bmatrix} D_{s-1 \times k}(t) \\ \frac{D_{(s+m-1) \times k}(\eta)}{D_{(s+m+l-1) \times k}(\xi)} \end{bmatrix}$ and the last two rows is equal to the following $2 \times k$ matrix $\begin{pmatrix} \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-1} \\ \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-1} \end{pmatrix}$, that is we get a matrix of the form

$$\left(\begin{array}{ccccc} \alpha_1 & \alpha_2 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\ \hline \beta_1 & \beta_2 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \dots & \beta_{k+m-1} & \beta_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \hline \gamma_1 & \gamma_2 & \dots & \gamma_{k-1} & \gamma_k \\ \gamma_2 & \gamma_3 & \dots & \gamma_k & \gamma_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{m+l+1} & \gamma_{m+l+2} & \dots & \gamma_{k+m+l-1} & \gamma_{k+m+l} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{s+m+l-1} & \gamma_{s+m+l} & \dots & \gamma_{s+m+l+k-3} & \gamma_{s+m+l+k-2} \\ \hline \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-2} & \alpha_{s+k-1} \\ \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \end{array} \right).$$

We introduce in a similar way the following notations

$$D \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \beta_{s+m-} \\ \alpha_{s-} \end{array} \right] \times k (t, \eta, \xi), \quad D \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} + \beta_{s+m-} \\ \gamma_{s+m+l-} \end{array} \right] \times k (t, \eta, \xi), \quad D \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \beta_{s+m-} + \gamma_{s+m+l-} \end{array} \right] \times k, \quad D \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} + \gamma_{s+m+l-} \\ \beta_{s+m-} \end{array} \right] \times k,$$

$$D \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \gamma_{s+m+l-} \end{array} \right] \times k (t, \eta, \xi), \quad D \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \gamma_{s+m+l-} \\ \alpha_{s-} \end{array} \right] \times k (t, \eta, \xi) \quad \text{and} \quad D \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} + \gamma_{s+m+l-} \end{array} \right] \times k (t, \eta, \xi)$$

- We define

$$\sigma_{i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \end{array} \right] \times k$$

to be the cardinality of the following set

$$\left\{ (t, \eta, \xi) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-2} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-2} \mid r(D \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \end{array} \right] \times k (t, \eta, \xi)) = r(D \left[\begin{array}{c} s \\ s+m-1 \\ s+m+l-1 \end{array} \right] \times k (t, \eta, \xi)) = i \right\}.$$

- We define

$$\sigma_{i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \alpha_{s-} + \beta_{s+m-} \end{array} \right] \times k$$

to be the cardinality of the following set

$$\left\{ (t, \eta, \xi) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-2} \mid r(D \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \end{array} \right] \times k (t, \eta, \xi)) = r(D \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \alpha_{s-} + \beta_{s+m-} \end{array} \right] \times k (t, \eta, \xi)) = i \right\}.$$

- We define

$$\sigma_{i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \alpha_{s-} + \gamma_{s+m+l-} \end{array} \right] \times k$$

to be the cardinality of the following set

$$\left\{ (t, \eta, \xi) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-2} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1} \mid r(D \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \end{array} \right] \times k (t, \eta, \xi)) = r(D \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \alpha_{s-} + \gamma_{s+m+l-} \end{array} \right] \times k (t, \eta, \xi)) = i \right\}.$$

- We define

$$\sigma_{i,i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \end{array} \right] \times k = \sigma_{i,i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \beta_{s+m-} \\ \alpha_{s-} \end{array} \right] \times k$$

to be the cardinality of the following set

$$\left\{ (t, \eta, \xi) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-2} \mid r(D \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \end{array} \right] \times k (t, \eta, \xi)) = r(D \left[\begin{array}{c} s \\ s+m-1 \\ s+m+l-1 \end{array} \right] \times k (t, \eta, \xi)) = r(D \left[\begin{array}{c} s \\ s+m-1 \\ s+m+l-1 \end{array} \right] \times k (t, \eta, \xi)) = i \right\}.$$

- We define

$$\sigma_{i,i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} + \gamma_{s+m+l-} \end{array} \right] \times k$$

to be the cardinality of the following set

$$\left\{ (t, \eta, \xi) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1} \mid r(D \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \end{array} \right] \times k (t, \eta, \xi)) = r(D \left[\begin{array}{c} s \\ s+m-1 \\ s+m+l-1 \end{array} \right] \times k (t, \eta, \xi)) = r(D \left[\begin{array}{c} s \\ s+m-1 \\ s+m+l-1 \\ \hline \beta_{s+m-} + \gamma_{s+m+l-} \end{array} \right] \times k (t, \eta, \xi)) = i \right\}.$$

- We introduce in a similar way the following definitions

$$\sigma_{i,i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \alpha_{s-} + \beta_{s+m-} \\ \gamma_{s+m+l-} \end{array} \right] \times k, \quad \sigma_{i,i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \alpha_{s-} + \beta_{s+m-} \\ \beta_{s+m-} + \gamma_{s+m+l-} \end{array} \right] \times k, \quad \sigma_{i,i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \alpha_{s-} \\ \gamma_{s+m+l-} \end{array} \right] \times k, \quad \text{and} \quad \sigma_{i,i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \alpha_{s-} + \gamma_{s+m+l-} \\ \beta_{s+m-} \end{array} \right] \times k.$$

- Let $(j_1, j_2, j_3, j_4) \in \mathbb{N}^4$ we define

$$\sigma_{j_1, j_2, j_3, j_4} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{array} \right] \times k = \sigma_{j_1, j_2, j_3, j_4} \left[\begin{array}{c} \alpha \\ \beta \\ \gamma \end{array} \right] \times k$$

to be the cardinality of the following set

$$\left\{ (t, \eta, \xi) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1} \mid r(D \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \end{array} \right] \times k (t, \eta, \xi)) = j_1 \right. \\ \left. r(D \left[\begin{array}{c} s \\ s+m-1 \end{array} \right] \times k (t, \eta, \xi)) = j_2, \quad r(D \left[\begin{array}{c} s \\ s+m+l-1 \end{array} \right] \times k (t, \eta, \xi)) = j_3, \quad r(D \left[\begin{array}{c} s \\ s+m+l \end{array} \right] \times k (t, \eta, \xi)) = j_4 \right\}$$

For any permutation of $\{\alpha, \beta, \gamma\}$ we define in a similar manner

$$\begin{aligned} \sigma_{j_1, j_2, j_3, j_4} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \beta_{s+m-} \\ \alpha_{s-} \\ \gamma_{s+m+l-} \end{array} \right] \times k &= \sigma_{j_1, j_2, j_3, j_4} \left[\begin{array}{c} \beta \\ \alpha \\ \gamma \end{array} \right] \times k, \\ \sigma_{j_1, j_2, j_3, j_4} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \beta_{s+m-} \\ \gamma_{s+m+l-} \\ \alpha_{s-} \end{array} \right] \times k &= \sigma_{j_1, j_2, j_3, j_4} \left[\begin{array}{c} \beta \\ \gamma \\ \alpha \end{array} \right] \times k, \\ \sigma_{j_1, j_2, j_3, j_4} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \alpha_{s-} \\ \gamma_{s+m+l-} \\ \beta_{s+m-} \end{array} \right] \times k &= \sigma_{j_1, j_2, j_3, j_4} \left[\begin{array}{c} \alpha \\ \gamma \\ \beta \end{array} \right] \times k, \\ \sigma_{j_1, j_2, j_3, j_4} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \gamma_{s+m+l-} \\ \alpha_{s-} \\ \beta_{s+m-} \end{array} \right] \times k &= \sigma_{j_1, j_2, j_3, j_4} \left[\begin{array}{c} \gamma \\ \alpha \\ \beta \end{array} \right] \times k, \\ \sigma_{j_1, j_2, j_3, j_4} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \beta_{s+m-} \\ \gamma_{s+m+l-} \\ \alpha_{s-} \end{array} \right] \times k &= \sigma_{j_1, j_2, j_3, j_4} \left[\begin{array}{c} \gamma \\ \beta \\ \alpha \end{array} \right] \times k, \end{aligned}$$

•

The sum $\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix}\right] \times k} + \sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix}\right] \times k} + \sigma_{i-1,i-1,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix}\right] \times k} + \sigma_{i-2,i-1,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix}\right] \times k}$
 is equal to the sum $\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \beta \\ \gamma \\ \alpha \end{smallmatrix}\right] \times k} + \sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \beta \\ \gamma \\ \alpha \end{smallmatrix}\right] \times k} + \sigma_{i-1,i-1,i,i}^{\left[\begin{smallmatrix} \beta \\ \gamma \\ \alpha \end{smallmatrix}\right] \times k} + \sigma_{i-2,i-1,i,i}^{\left[\begin{smallmatrix} \beta \\ \gamma \\ \alpha \end{smallmatrix}\right] \times k}$
 and is also equal to the sum $\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \gamma \\ \beta \\ \alpha \end{smallmatrix}\right] \times k} + \sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \gamma \\ \beta \\ \alpha \end{smallmatrix}\right] \times k} + \sigma_{i-1,i-1,i,i}^{\left[\begin{smallmatrix} \gamma \\ \beta \\ \alpha \end{smallmatrix}\right] \times k} + \sigma_{i-2,i-1,i,i}^{\left[\begin{smallmatrix} \gamma \\ \beta \\ \alpha \end{smallmatrix}\right] \times k}.$

Analogous sums are defined in a similar way

•

$$\Gamma_i^{\left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix}\right] \times k} = \text{Card} \left\{ (t, \eta, \xi) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1} \mid r(D^{\left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix}\right] \times k}(t, \eta, \xi)) = i \right\}.$$

We recall that the rank of a matrix does not change under elementary row operations.

- We denote by $\ker D$ the nullspace of the matrix D and $r(D)$ the rank of the matrix D .
- To simplify the notations concerning the exponential sums used in the proofs, we introduce the following definitions.
- Let $g_1(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$ defined by

$$(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P} \mapsto \sum_{\deg Y \leq k-1} \sum_{\deg Z = s-1} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V \leq s+m+l-1} E(\xi YV) \in \mathbb{Z}$$

- Let $h_1(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$ defined by

$$(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P} \mapsto \sum_{\deg Y \leq k-1} E(tYT^{s-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \in \mathbb{Z}$$

- Let $h_2(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$ defined by

$$(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P} \mapsto$$

$$\sum_{\deg Y \leq k-1} E(tYT^{s-1}) E(\eta YT^{s+m-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \in \mathbb{Z}$$

- Let $h_3(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$ defined by

$$(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P} \mapsto$$

$$\sum_{\deg Y \leq k-1} E(tYT^{s-1}) E(\xi YT^{s+m+l-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \in \mathbb{Z}$$

- Let $h_4(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$ defined by

$$(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P} \mapsto \sum_{\deg Y \leq k-1} E(tYT^{s-1})E(\eta YT^{s+m-1})E(\xi YT^{s+m+l-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \in \mathbb{Z}$$

- Let $\varphi(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$ defined by

$$(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P} \mapsto \sum_{\deg Y \leq k-1} \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \in \mathbb{Z}$$

- Let $\varphi_1(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$ defined by

$$(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P} \mapsto \sum_{\deg Y \leq k-1} \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V \leq s+m+l-1} E(\xi YV) \in \mathbb{Z}$$

- Let $\varphi_2(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$ defined by

$$(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P} \mapsto \sum_{\deg Y \leq k-1} \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-1} E(\xi YV) \in \mathbb{Z}$$

- Let $\varphi_3(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$ defined by

$$(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P} \mapsto \sum_{\deg Y \leq k-1} \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \in \mathbb{Z}$$

- Let $\kappa_1(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$ defined by

$$(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P} \mapsto$$

$$\sum_{\deg Y \leq k-1} E(tYT^{s-1})E(\eta YT^{s+m-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-1} E(\xi YV) \in \mathbb{Z}$$

- Let $\kappa_2(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$ defined by

$$(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P} \mapsto \sum_{\deg Y \leq k-1} E(tYT^{s-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-1} E(\xi YV) \in \mathbb{Z}$$

- Let $g_2(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$ defined by

$$(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P} \mapsto \sum_{\deg Y \leq k-1} E(tYT^{s-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \in \mathbb{Z}$$

2.2. Introduction. Adapting the method used in Section 5 of [2] we establish in this section a recurrent formula for the number of rank i matrices of the form $\begin{bmatrix} A \\ B \\ C \end{bmatrix}$, where A, B and C are persymmetric.

2.3. Computation of exponential sums in $\mathbb{K}^3$.

Lemma 2.2. *Let $(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P}$ and*

$$\begin{aligned}
g_1(t, \eta, \xi) &= \sum_{\deg Y \leq k-1} \sum_{\deg Z = s-1} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V \leq s+m+l-1} E(\xi YV) \\
h_1(t, \eta, \xi) &= \sum_{\deg Y \leq k-1} E(tYT^{s-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \\
h_2(t, \eta, \xi) &= \sum_{\deg Y \leq k-1} E(tYT^{s-1}) E(\eta YT^{s+m-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \\
h_3(t, \eta, \xi) &= \sum_{\deg Y \leq k-1} E(tYT^{s-1}) E(\xi YT^{s+m+l-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \\
h_4(t, \eta, \xi) &= \sum_{\deg Y \leq k-1} E(tYT^{s-1}) E(\eta YT^{s+m-1}) E(\xi YT^{s+m+l-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \\
\varphi(t, \eta, \xi) &= \sum_{\deg Y \leq k-1} \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \\
\varphi_1(t, \eta, \xi) &= \sum_{\deg Y \leq k-1} \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V \leq s+m+l-1} E(\xi YV) \\
\varphi_2(t, \eta, \xi) &= \sum_{\deg Y \leq k-1} \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-1} E(\xi YV) \\
\varphi_3(t, \eta, \xi) &= \sum_{\deg Y \leq k-1} \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \\
\kappa_1(t, \eta, \xi) &= \sum_{\deg Y \leq k-1} E(tYT^{s-1}) E(\eta YT^{s+m-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-1} E(\xi YV) \\
\kappa_2(t, \eta, \xi) &= \sum_{\deg Y \leq k-1} E(tYT^{s-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-1} E(\xi YV) \\
g_2(t, \eta, \xi) &= \sum_{\deg Y \leq k-1} E(tYT^{s-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV)
\end{aligned}$$

Then

(2.1)

$$\begin{aligned}
g_1(t, \eta, \xi) &= 2^{3s+2m+l-1} \cdot \sum_{\substack{\deg Y \leq k-1 \\ \begin{bmatrix} s-1 \\ s+m \\ s+m+l \end{bmatrix} \times k \\ Y \in \ker D}} E(tYT^{s-1}) \\
&= \begin{cases} 2^{k+3s+2m+l-1-r(D \begin{bmatrix} s-1 \\ s+m \\ s+m+l \end{bmatrix} \times k)} (t, \eta, \xi) & \text{if } r(D \begin{bmatrix} s-1 \\ s+m \\ s+m+l \end{bmatrix} \times k) (t, \eta, \xi) = r(D \begin{bmatrix} s \\ s+m \\ s+m+l \end{bmatrix} \times k) (t, \eta, \xi) \\ 0 & \text{otherwise .} \end{cases}
\end{aligned}$$

(2.2)

$$\begin{aligned}
h_1(t, \eta, \xi) &= 2^{3s+2m+l-3} \cdot \sum_{\substack{\deg Y \leq k-1 \\ \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k \\ Y \in \ker D}} E(tYT^{s-1}) \\
&= \begin{cases} 2^{k+3s+2m+l-3-r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k)} & \text{if } r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k) = r(D \begin{bmatrix} s \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k) \\ 0 & \text{otherwise} \end{cases} \quad (t, \eta, \xi)
\end{aligned}$$

(2.3)

$$\begin{aligned}
h_2(t, \eta, \xi) &= 2^{3s+2m+l-3} \cdot \sum_{\substack{\deg Y \leq k-1 \\ \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k \\ Y \in \ker D}} E(tYT^{s-1})E(\eta YT^{s+m-1}) \\
&= \begin{cases} 2^{k+3s+2m+l-3-r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k)} & \text{if } r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k) = r(D \begin{bmatrix} s-1 \\ s+m-1 \\ \frac{s+m-1}{\alpha_{s-} + \beta_{s+m-}} \end{bmatrix} \times k) \\ 0 & \text{otherwise} \end{cases} \quad (t, \eta, \xi)
\end{aligned}$$

(2.4)

$$\begin{aligned}
h_3(t, \eta, \xi) &= 2^{3s+2m+l-3} \cdot \sum_{\substack{\deg Y \leq k-1 \\ \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k \\ Y \in \ker D}} E(tYT^{s-1})E(\xi YT^{s+m+l-1}) \\
&= \begin{cases} 2^{k+3s+2m+l-3-r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k)} & \text{if } r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k) = r(D \begin{bmatrix} s-1 \\ s+m-1 \\ \frac{s+m-1}{\alpha_{s-} + \gamma_{s+m+l-}} \end{bmatrix} \times k) \\ 0 & \text{otherwise} \end{cases} \quad (t, \eta, \xi)
\end{aligned}$$

(2.5)

$$\begin{aligned}
h_4(t, \eta, \xi) &= 2^{3s+2m+l-3} \cdot \sum_{\substack{\deg Y \leq k-1 \\ \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k \\ Y \in \ker D}} E(tYT^{s-1})E(\eta YT^{s+m-1})E(\xi YT^{s+m+l-1}) \\
&= \begin{cases} 2^{k+3s+2m+l-3-r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k)} & \text{if } r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k) = r(D \begin{bmatrix} s-1 \\ s+m-1 \\ \frac{s+m-1}{\alpha_{s-} + \beta_{s+m-} + \gamma_{s+m+l-}} \end{bmatrix} \times k) \\ 0 & \text{otherwise} \end{cases} \quad (t, \eta, \xi)
\end{aligned}$$

(2.6)

$$\varphi(t, \eta, \xi) = 2^{3s+2m+l-3} \cdot \sum_{\substack{\deg Y \leq k-1 \\ \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k \\ Y \in \ker D}} 1 = 2^{k+3s+2m+l-3-r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k)} (t, \eta, \xi)$$

(2.7)

$$\varphi_1(t, \eta, \xi) = 2^{3s+2m+l-1} \cdot \sum_{\substack{\deg Y \leq k-1 \\ \begin{bmatrix} s-1 \\ s+m \\ s+m+l \end{bmatrix} \times k \\ Y \in \ker D}} 1 = 2^{k+3s+2m+l-1-r(D \begin{bmatrix} s-1 \\ s+m \\ s+m+l \end{bmatrix} \times k)} (t, \eta, \xi)$$

(2.8)

$$\varphi_2(t, \eta, \xi) = 2^{3s+2m+l-2} \cdot \sum_{\substack{\deg Y \leq k-1 \\ \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l \end{bmatrix} \times k \\ Y \in \ker D}} 1 = 2^{k+3s+2m+l-2-r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l \end{bmatrix} \times k)} (t, \eta, \xi)$$

(2.9)

$$\varphi_3(t, \eta, \xi) = 2^{3s+2m+l-2} \cdot \sum_{\substack{\deg Y \leq k-1 \\ \begin{bmatrix} s-1 \\ s+m \\ s+m+l-1 \end{bmatrix} \times k \\ Y \in \ker D}} 1 = 2^{k+3s+2m+l-2-r(D \begin{bmatrix} s-1 \\ s+m \\ s+m+l-1 \end{bmatrix} \times k)} (t, \eta, \xi)$$

(2.10)

$$\begin{aligned} \kappa_1(t, \eta, \xi) &= 2^{3s+2m+l-2} \cdot \sum_{\substack{\deg Y \leq k-1 \\ \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l \end{bmatrix} \times k \\ Y \in \ker D}} E(tYT^{s-1})E(\eta YT^{s+m-1}) \\ &= \begin{cases} 2^{k+3s+2m+l-2-r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l \end{bmatrix} \times k)} (t, \eta, \xi) & \text{if } r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l \end{bmatrix} \times k) = r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l \\ \alpha_{s-} + \beta_{s+m-} \end{bmatrix} \times k) \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

(2.11)

$$\begin{aligned} \kappa_2(t, \eta, \xi) &= 2^{3s+2m+l-2} \cdot \sum_{\substack{\deg Y \leq k-1 \\ \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l \end{bmatrix} \times k \\ Y \in \ker D}} E(tYT^{s-1}) \\ &= \begin{cases} 2^{k+3s+2m+l-2-r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l \end{bmatrix} \times k)} (t, \eta, \xi) & \text{if } r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l \end{bmatrix} \times k) = r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l \\ \alpha_{s-} \end{bmatrix} \times k) \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

(2.12)

$$\begin{aligned}
g_2(t, \eta, \xi) &= 2^{3s+2m+l-2} \cdot \sum_{\substack{\deg Y \leq k-1 \\ \begin{bmatrix} s-1 \\ s+m \\ s+m+l-1 \end{bmatrix} \times k \\ Y \in \ker D}} E(tYT^{s-1}) \\
&= \begin{cases} 2^{k+3s+2m+l-2-r(D \begin{bmatrix} s-1 \\ s+m \\ s+m+l-1 \end{bmatrix} \times k)} & \text{if } r(D \begin{bmatrix} s-1 \\ s+m \\ s+m+l-1 \end{bmatrix} \times k) = r(D \begin{bmatrix} s-1 \\ s+m \\ s+m+l-1 \\ \alpha_{s-} \end{bmatrix} \times k) \\ 0 & \text{otherwise} \end{cases} (t, \eta, \xi)
\end{aligned}$$

Proof. The proof of Lemma 2.2 is somewhat similar to the proofs of the results obtained in [2, see section 4]. $\square$

Lemma 2.3. *Let $(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P}$ and q a rational integer ≥ 2 , then we have*

$$(2.13) \quad g_1^q(t, \eta, \xi) = g_1(t, \eta, \xi) \cdot \varphi_1^{q-1}(t, \eta, \xi)$$

$$(2.14) \quad g_1(t, \eta, \xi) = \varphi_1(t, \eta, \xi) \quad \text{if} \quad g_1(t, \eta, \xi) \neq 0$$

$$(2.15) \quad h_1^q(t, \eta, \xi) = h_1(t, \eta, \xi) \cdot \varphi^{q-1}(t, \eta, \xi)$$

$$(2.16) \quad h_1(t, \eta, \xi) = \varphi(t, \eta, \xi) \quad \text{if} \quad h_1(t, \eta, \xi) \neq 0$$

$$(2.17) \quad h_2^q(t, \eta, \xi) = h_2(t, \eta, \xi) \cdot \varphi^{q-1}(t, \eta, \xi)$$

$$(2.18) \quad h_2(t, \eta, \xi) = \varphi(t, \eta, \xi) \quad \text{if} \quad h_2(t, \eta, \xi) \neq 0$$

$$(2.19) \quad h_3^q(t, \eta, \xi) = h_3(t, \eta, \xi) \cdot \varphi^{q-1}(t, \eta, \xi)$$

$$(2.20) \quad h_3(t, \eta, \xi) = \varphi(t, \eta, \xi) \quad \text{if} \quad h_3(t, \eta, \xi) \neq 0$$

$$(2.21) \quad h_4^q(t, \eta, \xi) = h_4(t, \eta, \xi) \cdot \varphi^{q-1}(t, \eta, \xi)$$

$$(2.22) \quad h_4(t, \eta, \xi) = \varphi(t, \eta, \xi) \quad \text{if} \quad h_4(t, \eta, \xi) \neq 0$$

$$(2.23) \quad \kappa_1^q(t, \eta, \xi) = \kappa_1(t, \eta, \xi) \cdot \varphi_2^{q-1}(t, \eta, \xi)$$

$$(2.24) \quad \kappa_1(t, \eta, \xi) = \varphi_2(t, \eta, \xi) \quad \text{if} \quad \kappa_1(t, \eta, \xi) \neq 0$$

$$(2.25) \quad \kappa_2^q(t, \eta, \xi) = \kappa_2(t, \eta, \xi) \cdot \varphi_2^{q-1}(t, \eta, \xi)$$

$$(2.26) \quad \kappa_2(t, \eta, \xi) = \varphi_2(t, \eta, \xi) \quad \text{if} \quad \kappa_2(t, \eta, \xi) \neq 0$$

$$(2.27) \quad g_2^q(t, \eta, \xi) = g_2(t, \eta, \xi) \cdot \varphi_3^{q-1}(t, \eta, \xi)$$

$$(2.28) \quad g_2(t, \eta, \xi) = \varphi_3(t, \eta, \xi) \quad \text{if} \quad g_2(t, \eta, \xi) \neq 0$$

Proof. To prove (2.13) and (2.14) we proceed as in [1 section 4], that is :
From (2.1) we get

$$\begin{aligned}
g_1^2(t, \eta, \xi) &= [2^{3s+2m+l-1} \cdot \sum_{\substack{\deg Y_1 \leq k-1 \\ \begin{bmatrix} s-1 \\ s+m \\ s+m+l \end{bmatrix} \times k \\ Y_1 \in \ker D}} E(tY_1T^{s-1})] [2^{3s+2m+l-1} \cdot \sum_{\substack{\deg Y_2 \leq k-1 \\ \begin{bmatrix} s-1 \\ s+m \\ s+m+l \end{bmatrix} \times k \\ Y_2 \in \ker D}} E(tY_2T^{s-1})] \\
&\quad (t, \eta, \xi) \quad (t, \eta, \xi)
\end{aligned}$$

We set $\begin{cases} Y_1 + Y_2 = Y_3 & \deg Y_3 \leq k-1, \\ Y_2 = Y_4 & \deg Y_4 \leq k-1. \end{cases}$

Then we obtain

$$\begin{aligned}
g_1^2(t, \eta, \xi) &= 2^{(3s+2m+l-1)2} \sum_{\substack{\deg Y_1 \leq k-1 \\ Y_1 \in \ker D}} \left[\begin{smallmatrix} s-1 \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k \sum_{\substack{\deg Y_2 \leq k-1 \\ Y_2 \in \ker D}} \left[\begin{smallmatrix} s-1 \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k E(t(Y_1 + Y_2)T^{s-1}) \\
&= [2^{3s+2m+l-1} \cdot \sum_{\substack{\deg Y_3 \leq k-1 \\ Y_3 \in \ker D}} \left[\begin{smallmatrix} s-1 \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k E(tY_3T^{s-1})] [2^{3s+2m+l-1} \cdot \sum_{\substack{\deg Y_4 \leq k-1 \\ Y_4 \in \ker D}} \left[\begin{smallmatrix} s-1 \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k 1] \\
&= g_1(t, \eta, \xi) \cdot \varphi_1(t, \eta, \xi)
\end{aligned}$$

The other assertions are proved in a similar way. $\square$

Lemma 2.4. *Let $(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P}$, and q a rational integer ≥ 2 , then we have for $1 \leq i \leq q-1$*

(2.29)

$$\begin{aligned}
h_1^i(t, \eta, \xi) \cdot h_2^{q-i}(t, \eta, \xi) &= \\
&= \begin{cases} \varphi^q(t, \eta, \xi) & \text{if } r(D \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times k (t, \eta, \xi)) = r(D \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ \frac{s+m+l-1}{\alpha_{s-}} \end{smallmatrix} \right] \times k (t, \eta, \xi)) = r(D \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ \frac{s+m+l-1}{\beta_{s+m-}} \end{smallmatrix} \right] \times k (t, \eta, \xi)) \\ 0 & \text{otherwise .} \end{cases}
\end{aligned}$$

(2.30)

$$\begin{aligned}
h_1^i(t, \eta, \xi) \cdot h_3^{q-i}(t, \eta, \xi) &= \\
&= \begin{cases} \varphi^q(t, \eta, \xi) & \text{if } r(D \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times k (t, \eta, \xi)) = r(D \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ \frac{s+m+l-1}{\alpha_{s-}} \end{smallmatrix} \right] \times k (t, \eta, \xi)) = r(D \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ \frac{s+m+l-1}{\gamma_{s+m+l-}} \end{smallmatrix} \right] \times k (t, \eta, \xi)) \\ 0 & \text{otherwise .} \end{cases}
\end{aligned}$$

(2.31)

$$\begin{aligned}
h_1^i(t, \eta, \xi) \cdot h_4^{q-i}(t, \eta, \xi) &= \\
&= \begin{cases} \varphi^q(t, \eta, \xi) & \text{if } r(D \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times k (t, \eta, \xi)) = r(D \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ \frac{s+m+l-1}{\alpha_{s-}} \end{smallmatrix} \right] \times k (t, \eta, \xi)) = r(D \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ \frac{s+m+l-1}{\beta_{s+m-} + \gamma_{s+m+l-}} \end{smallmatrix} \right] \times k (t, \eta, \xi)) \\ 0 & \text{otherwise .} \end{cases}
\end{aligned}$$

(2.32)

$$\begin{aligned}
h_2^i(t, \eta, \xi) \cdot h_3^{q-i}(t, \eta, \xi) &= \\
&= \begin{cases} \varphi^q(t, \eta, \xi) & \text{if } r(D \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times k (t, \eta, \xi)) = r(D \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ \frac{s+m+l-1}{\alpha_{s-} + \beta_{s+m-}} \end{smallmatrix} \right] \times k (t, \eta, \xi)) = r(D \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ \frac{s+m+l-1}{\beta_{s+m-} + \gamma_{s+m+l-}} \end{smallmatrix} \right] \times k (t, \eta, \xi)) \\ 0 & \text{otherwise .} \end{cases}
\end{aligned}$$

(2.33)

$$h_2^i(t, \eta, \xi) \cdot h_4^{q-i}(t, \eta, \xi) = \begin{cases} \varphi^q(t, \eta, \xi) & \text{if } r\left(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k\right)(t, \eta, \xi) = r\left(D \begin{bmatrix} s-1 \\ s+m-1 \\ \frac{s+m+l-1}{\alpha_{s-}+\beta_{s+m-}} \end{bmatrix} \times k\right)(t, \eta, \xi) = r\left(D \begin{bmatrix} s-1 \\ s+m-1 \\ \frac{s+m+l-1}{\gamma_{s+m+l-}} \end{bmatrix} \times k\right)(t, \eta, \xi) \\ 0 & \text{otherwise .} \end{cases}$$

(2.34)

$$h_3^i(t, \eta, \xi) \cdot h_4^{q-i}(t, \eta, \xi) = \begin{cases} \varphi^q(t, \eta, \xi) & \text{if } r\left(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k\right)(t, \eta, \xi) = r\left(D \begin{bmatrix} s-1 \\ s+m-1 \\ \frac{s+m+l-1}{\alpha_{s-}+\gamma_{s+m+l-}} \end{bmatrix} \times k\right)(t, \eta, \xi) = r\left(D \begin{bmatrix} s-1 \\ s+m-1 \\ \frac{s+m+l-1}{\beta_{s+m-}} \end{bmatrix} \times k\right)(t, \eta, \xi) \\ 0 & \text{otherwise .} \end{cases}$$

(2.35)

$$\kappa_1^i(t, \eta, \xi) \cdot \kappa_2^{q-i}(t, \eta, \xi) = \begin{cases} \varphi_2^q(t, \eta, \xi) & \text{if } r\left(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l \end{bmatrix} \times k\right)(t, \eta, \xi) = r\left(D \begin{bmatrix} s-1 \\ s+m-1 \\ \frac{s+m+l}{\alpha_{s-}+\beta_{s+m-}} \end{bmatrix} \times k\right)(t, \eta, \xi) = r\left(D \begin{bmatrix} s-1 \\ s+m-1 \\ \frac{s+m+l}{\alpha_{s-}} \end{bmatrix} \times k\right)(t, \eta, \xi) \\ 0 & \text{otherwise .} \end{cases}$$

Proof. Follows from Lemma 2.2 and Lemma 2.3 using elementary rank considerations. $\square$

Lemma 2.5. *Let $(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P}$, and q a rational integer ≥ 3 , then we have for any integers i, j, r such that*

$$\begin{cases} i + j + r = q \\ 1 \leq i, j, r \leq q - 2 \end{cases}$$

(2.36)

$$\begin{aligned} h_1^i(t, \eta, \xi) \cdot h_2^j(t, \eta, \xi) \cdot h_3^r(t, \eta, \xi) &= h_1^i(t, \eta, \xi) \cdot h_3^j(t, \eta, \xi) \cdot h_4^r(t, \eta, \xi) \\ &= h_2^i(t, \eta, \xi) \cdot h_3^j(t, \eta, \xi) \cdot h_4^r(t, \eta, \xi) = h_1^i(t, \eta, \xi) \cdot h_2^j(t, \eta, \xi) \cdot h_4^r(t, \eta, \xi) \\ &= \begin{cases} \varphi^q(t, \eta, \xi) & \text{if } r\left(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k\right)(t, \eta, \xi) = r\left(D \begin{bmatrix} s-1 \\ s+m-1 \\ \frac{s+m+l-1}{\alpha_{s-}} \end{bmatrix} \times k\right)(t, \eta, \xi) = r\left(D \begin{bmatrix} s-1 \\ s+m-1 \\ \frac{s+m+l-1}{\beta_{s+m-}} \end{bmatrix} \times k\right)(t, \eta, \xi) = r\left(D \begin{bmatrix} s-1 \\ s+m-1 \\ \frac{s+m+l-1}{\gamma_{s+m+l-}} \end{bmatrix} \times k\right)(t, \eta, \xi) \\ 0 & \text{otherwise .} \end{cases} \end{aligned}$$

Proof. Follows from Lemma 2.2 and Lemma 2.3 using elementary row operations. $\square$

Lemma 2.6. *Let $(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P}$, and q a rational integer ≥ 4 , then we have for any integers i, j, r, p such that*

$$\begin{cases} i + j + r + p = q \\ 1 \leq i, j, r, p \leq q - 3 \end{cases}$$

(2.37)

$$h_1^i(t, \eta, \xi) \cdot h_2^j(t, \eta, \xi) \cdot h_3^r(t, \eta, \xi) \cdot h_4^p(t, \eta, \xi)$$

$$= \begin{cases} \varphi^q(t, \eta, \xi) & \text{if } r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k) (t, \eta, \xi) = r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \end{bmatrix} \times k) (t, \eta, \xi) = r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \beta_{s+m-} \end{bmatrix} \times k) (t, \eta, \xi) = r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \gamma_{s+m+l-} \end{bmatrix} \times k) (t, \eta, \xi) \\ 0 & \text{otherwise .} \end{cases}$$

Proof. Similarly to the proof of Lemma 2.5. $\square$

2.4. Rank properties of partitions on triple persymmetric matrices.

Lemma 2.7. *We have :*

$$(2.38) \quad \sigma_{i,i} \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} + \beta_{s+m-} + \gamma_{s+m+l-} \end{bmatrix} \times k = 2 \cdot \sigma_{i,i} \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} + \beta_{s+m-} \end{bmatrix} \times k = 2 \cdot \sigma_{i,i} \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} + \gamma_{s+m+l-} \end{bmatrix} \times k = 4 \cdot \sigma_{i,i} \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \end{bmatrix} \times k$$

$$(2.39) \quad \sigma_{i,i} \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l \\ \alpha_{s-} + \beta_{s+m-} \end{bmatrix} \times k = 2 \cdot \sigma_{i,i} \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l \\ \alpha_{s-} \end{bmatrix} \times k$$

$$(2.40) \quad \sigma_{i,i,i} \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} + \gamma_{s+m+l-} \end{bmatrix} \times k = 2 \cdot \sigma_{i,i,i} \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \end{bmatrix} \times k = 2 \cdot \sigma_{i,i,i} \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \beta_{s+m-} \\ \alpha_{s-} \end{bmatrix} \times k$$

$$(2.41) \quad \sigma_{i,i,i} \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} + \beta_{s+m-} \\ \beta_{s+m-} + \gamma_{s+m+l-} \end{bmatrix} \times k = 2 \cdot \sigma_{i,i,i} \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \end{bmatrix} \times k = 2 \cdot \sigma_{i,i,i} \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \beta_{s+m-} \\ \alpha_{s-} \end{bmatrix} \times k$$

$$(2.42) \quad \sigma_{i,i,i} \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} + \beta_{s+m-} \\ \gamma_{s+m+l-} \end{bmatrix} \times k = 2 \cdot \sigma_{i,i,i} \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \gamma_{s+m+l-} \end{bmatrix} \times k = 2 \cdot \sigma_{i,i,i} \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \gamma_{s+m+l-} \\ \alpha_{s-} \end{bmatrix} \times k$$

$$(2.43) \quad \sigma_{i,i,i} \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} + \gamma_{s+m+l-} \\ \beta_{s+m-} \end{bmatrix} \times k = 2 \cdot \sigma_{i,i,i} \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \end{bmatrix} \times k = 2 \cdot \sigma_{i,i,i} \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \beta_{s+m-} \\ \alpha_{s-} \end{bmatrix} \times k$$

Proof. By way of example we establish the following equality in (2.38)

$$\sigma_{i,i} \left[\begin{array}{c} \frac{s-1}{s+m+l-1} \\ \alpha_{s-} + \beta_{s+m-} + \gamma_{s+m+l-} \end{array} \right] \times k = 4 \cdot \sigma_{i,i} \left[\begin{array}{c} \frac{s-1}{s+m+l-1} \\ \alpha_{s-} \end{array} \right] \times k$$

Consider the following $(3s+2m+l-2) \times k$ matrix denoted by D $\left[\begin{array}{c} \frac{s-1}{s+m+l-1} \\ \alpha_{s-} + \beta_{s+m-} + \gamma_{s+m+l-} \end{array} \right] \times k$ (t, η, ξ)

$$\begin{array}{l} r_1 \\ r_2 \\ \vdots \\ r_{s-1} \\ r_s \\ r_{s+1} \\ \vdots \\ r_{s+m} \\ \vdots \\ r_{2s+m-2} \\ r_{2s+m-1} \\ r_{2s+m} \\ \vdots \\ r_{2s+2m+l-1} \\ \vdots \\ r_{3s+2m-3} \\ r_{3s+2m-2} \end{array} \left(\begin{array}{cccccc} \alpha_1 & \alpha_2 & \dots & \alpha_{k-1} & \alpha_k & \\ \alpha_2 & \alpha_3 & \dots & \alpha_k & \alpha_{k+1} & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \\ \alpha_{s-1} & \alpha_s & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} & \\ \beta_1 & \beta_2 & \dots & \beta_{k-1} & \beta_k & \\ \beta_2 & \beta_3 & \dots & \beta_k & \beta_{k+1} & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \\ \beta_{m+1} & \beta_{m+2} & \dots & \beta_{k+m-1} & \beta_{k+m} & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \\ \beta_{s+m-1} & \beta_{s+m} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} & \\ \gamma_1 & \gamma_2 & \dots & \gamma_{k-1} & \gamma_k & \\ \gamma_2 & \gamma_3 & \dots & \gamma_k & \gamma_{k+1} & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \\ \gamma_{m+l+1} & \gamma_{m+l+2} & \dots & \gamma_{k+m+l-1} & \gamma_{k+m+l} & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \\ \gamma_{s+m+l-1} & \gamma_{s+m+l} & \dots & \gamma_{s+m+l+k-3} & \gamma_{s+m+l+k-2} & \\ \alpha_s + \beta_{s+m} + \gamma_{s+m+l} & \alpha_{s+1} + \beta_{s+m+1} + \gamma_{s+m+l+1} & \dots & \alpha_{s+k-2} + \beta_{s+k+m-2} + \gamma_{s+m+l+k-2} & \alpha_{s+k-1} + \beta_{s+k+m-1} + \gamma_{s+m+l+k-1} & \end{array} \right)$$

We recall that the rank of a matrix does not change under elementary row operations.

On the above matrix, we add to the j -th row the $s+m+j$ -th row and the $2s+2m+l-1+j-th$ row for $0 \leq j \leq s-2$ obtaining

$$\left(\begin{array}{cccccc} \alpha_1 + \beta_{m+1} + \gamma_{m+l+1} & \alpha_2 + \beta_{m+2} + \gamma_{m+l+2} & \dots & \alpha_{k-1} + \beta_{m+k-1} + \gamma_{m+l+k-1} & \alpha_k + \beta_{m+k} + \gamma_{m+l+k} & \\ \alpha_2 + \beta_{m+2} + \gamma_{m+l+2} & \alpha_3 + \beta_{m+3} + \gamma_{m+l+3} & \dots & \alpha_k + \beta_{m+k} + \gamma_{m+l+k} & \alpha_{k+1} + \beta_{m+k+1} + \gamma_{m+l+k+1} & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \\ \alpha_{s-1} + \beta_{s-1+m} + \gamma_{s-1+m+l} & \alpha_s + \beta_{s+m} + \gamma_{s+m+l} & \dots & \alpha_{s+k-3} + \beta_{s+k+m-3} + \gamma_{s+k+m+l-3} & \alpha_{s+k-2} + \beta_{s+k+m-2} + \gamma_{s+k+m+l-2} & \\ \beta_1 & \beta_2 & \dots & \beta_{k-1} & \beta_k & \\ \beta_2 & \beta_3 & \dots & \beta_k & \beta_{k+1} & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \\ \beta_{m+1} & \beta_{m+2} & \dots & \beta_{k+m-1} & \beta_{k+m} & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \\ \beta_{s+m-1} & \beta_{s+m} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} & \\ \gamma_1 & \gamma_2 & \dots & \gamma_{k-1} & \gamma_k & \\ \gamma_2 & \gamma_3 & \dots & \gamma_k & \gamma_{k+1} & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \\ \gamma_{m+l+1} & \gamma_{m+l+2} & \dots & \gamma_{k+m+l-1} & \gamma_{k+m+l} & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \\ \gamma_{s+m+l-1} & \gamma_{s+m+l} & \dots & \gamma_{s+m+l+k-3} & \gamma_{s+m+l+k-2} & \\ \alpha_s + \beta_{s+m} + \gamma_{s+m+l} & \alpha_{s+1} + \beta_{s+m+1} + \gamma_{s+m+l+1} & \dots & \alpha_{s+k-2} + \beta_{s+m+k-2} + \gamma_{s+m+l+k-2} & \alpha_{s+k-1} + \beta_{s+m+k-1} + \gamma_{s+m+l+k-1} & \end{array} \right)$$

↑

In the above matrix we set

$$\begin{cases} \alpha_i + \beta_{i+m} + \gamma_{i+m+l} &= \alpha'_i & \text{for } 1 \leq i \leq k+s-2, \\ \beta_j &= \beta_j & \text{for } 1 \leq j \leq k+s+m-2. \\ \gamma_r &= \gamma_r & \text{for } 1 \leq r \leq k+s+m+l-2. \end{cases}$$

We remark that the

map $\kappa : \mathbb{F}_2^{3k+3s+2m+l-6} \mapsto \mathbb{F}_2^{3k+3s+2m+l-6}$ defined by

$$(\alpha_1, \alpha_2, \dots, \alpha_{k+s-2}, \beta_1, \beta_2, \dots, \beta_{k+s+m-2}, \gamma_1, \gamma_2, \dots, \gamma_{k+s+m+l-2})$$

$$\mapsto (\alpha'_1, \alpha'_2, \dots, \alpha'_{k+s-2}, \beta_1, \beta_2, \dots, \beta_{k+s+m-2}, \gamma_1, \gamma_2, \dots, \gamma_{k+s+m+l-2})$$

is an isomorphisme. We then obtain the below matrix

↓

$$\begin{pmatrix} \alpha'_1 & \alpha'_2 & \dots & \alpha'_{k-1} & \alpha'_k \\ \alpha'_2 & \alpha'_3 & \dots & \alpha'_k & \alpha'_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha'_{s-1} & \alpha'_s & \dots & \alpha'_{s+k-3} & \alpha'_{s+k-2} \\ \beta_1 & \beta_2 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \dots & \beta_{k+m-1} & \beta_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \gamma_1 & \gamma_2 & \dots & \gamma_{k-1} & \gamma_k \\ \gamma_2 & \gamma_3 & \dots & \gamma_k & \gamma_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{m+l+1} & \gamma_{m+l+2} & \dots & \gamma_{k+m+l-1} & \gamma_{k+m+l} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{s+m+l-1} & \gamma_{s+m+l} & \dots & \gamma_{s+m+l+k-3} & \gamma_{s+m+l+k-2} \\ \hline \alpha'_s & \alpha'_{s+1} & \dots & \alpha'_{s+k-2} & \alpha_{s+k-1} + \beta_{s+m+k-1} + \gamma_{s+m+l+k-1} \end{pmatrix}$$

We then get the following equivalences :

$$\text{rank} \begin{pmatrix} \alpha_1 & \alpha_2 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\ \beta_1 & \beta_2 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \dots & \beta_{k+m-1} & \beta_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \gamma_1 & \gamma_2 & \dots & \gamma_{k-1} & \gamma_k \\ \gamma_2 & \gamma_3 & \dots & \gamma_k & \gamma_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{m+l+1} & \gamma_{m+l+2} & \dots & \gamma_{k+m+l-1} & \gamma_{k+m+l} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{s+m+l-1} & \gamma_{s+m+l} & \dots & \gamma_{s+m+l+k-3} & \gamma_{s+m+l+k-2} \\ \hline \alpha_s + \beta_{s+m} + \gamma_{s+m+l} & \alpha_{s+1} + \beta_{s+m+1} + \gamma_{s+m+l+1} & \dots & \alpha_{s+k-2} + \beta_{s+m+k-2} + \gamma_{s+m+l+k-2} & \alpha_{s+k-1} + \beta_{s+m+k-1} + \gamma_{s+m+l+k-1} \end{pmatrix}$$

$$\begin{aligned}
&= \text{rank} \begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\ \beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \dots & \beta_{k+m-1} & \beta_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \gamma_1 & \gamma_2 & \gamma_3 & \dots & \gamma_{k-1} & \gamma_k \\ \gamma_2 & \gamma_3 & \gamma_4 & \dots & \gamma_k & \gamma_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{m+l+1} & \gamma_{m+l+2} & \gamma_{m+l+3} & \dots & \gamma_{k+m+l-1} & \gamma_{k+m+l} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{s+m+l-1} & \gamma_{s+m+l} & \gamma_{s+m+l+1} & \dots & \gamma_{s+m+l+k-3} & \gamma_{s+m+l+k-2} \end{pmatrix} \\
&\iff \\
&\text{rank} \begin{pmatrix} \alpha_1+\beta_{m+1}+\gamma_{m+l+1} & \alpha_2+\beta_{m+2}+\gamma_{m+l+2} & \dots & \alpha_{k-1}+\beta_{m+k-1}+\gamma_{m+l+k-1} & \alpha_k+\beta_{m+k}+\gamma_{m+l+k} \\ \alpha_2+\beta_{m+2}+\gamma_{m+l+2} & \alpha_3+\beta_{m+3}+\gamma_{m+l+3} & \dots & \alpha_k+\beta_{m+k}+\gamma_{m+l+k} & \alpha_{k+1}+\beta_{m+k+1}+\gamma_{m+l+k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1}+\beta_{s-1+m}+\gamma_{s-1+m+l} & \alpha_s+\beta_{s+m}+\gamma_{s+m+l} & \dots & \alpha_{s+k-3}+\beta_{s+k+m-3}+\gamma_{s+k+m+l-3} & \alpha_{s+k-2}+\beta_{s+k+m-2}+\gamma_{s+k+m+l-2} \\ \beta_1 & \beta_2 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \dots & \beta_{k+m-1} & \beta_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \gamma_1 & \gamma_2 & \dots & \gamma_{k-1} & \gamma_k \\ \gamma_2 & \gamma_3 & \dots & \gamma_k & \gamma_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{m+l+1} & \gamma_{m+l+2} & \dots & \gamma_{k+m+l-1} & \gamma_{k+m+l} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{s+m+l-1} & \gamma_{s+m+l} & \dots & \gamma_{s+m+l+k-3} & \gamma_{s+m+l+k-2} \\ \hline \alpha_s+\beta_{s+m}+\gamma_{s+m+l} & \alpha_{s+1}+\beta_{s+m+1}+\gamma_{s+m+l+1} & \dots & \alpha_{s+k-2}+\beta_{s+m+k-2}+\gamma_{s+m+l+k-2} & \alpha_{s+k-1}+\beta_{s+m+k-1}+\gamma_{s+m+l+k-1} \end{pmatrix} \\
&= \text{rank} \begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\ \beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \dots & \beta_{k+m-1} & \beta_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \gamma_1 & \gamma_2 & \gamma_3 & \dots & \gamma_{k-1} & \gamma_k \\ \gamma_2 & \gamma_3 & \gamma_4 & \dots & \gamma_k & \gamma_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{m+l+1} & \gamma_{m+l+2} & \gamma_{m+l+3} & \dots & \gamma_{k+m+l-1} & \gamma_{k+m+l} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{s+m+l-1} & \gamma_{s+m+l} & \gamma_{s+m+l+1} & \dots & \gamma_{s+m+l+k-3} & \gamma_{s+m+l+k-2} \end{pmatrix} \\
&\iff
\end{aligned}$$

$$\begin{aligned}
& \text{rank} \left(\begin{array}{cccccc}
\alpha'_1 & \alpha'_2 & \dots & \alpha'_{k-1} & & \alpha'_k \\
\alpha'_2 & \alpha'_3 & \dots & \alpha'_k & & \alpha'_{k+1} \\
\vdots & \vdots & \vdots & \vdots & & \vdots \\
\alpha'_{s-1} & \alpha'_s & \dots & \alpha'_{s+k-3} & & \alpha'_{s+k-2} \\
\beta_1 & \beta_2 & \dots & \beta_{k-1} & & \beta_k \\
\beta_2 & \beta_3 & \dots & \beta_k & & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & & \vdots \\
\beta_{m+1} & \beta_{m+2} & \dots & \beta_{k+m-1} & & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \dots & \beta_{s+m+k-3} & & \beta_{s+m+k-2} \\
\gamma_1 & \gamma_2 & \dots & \gamma_{k-1} & & \gamma_k \\
\gamma_2 & \gamma_3 & \dots & \gamma_k & & \gamma_{k+1} \\
\vdots & \vdots & \vdots & \vdots & & \vdots \\
\gamma_{m+l+1} & \gamma_{m+l+2} & \dots & \gamma_{k+m+l-1} & & \gamma_{k+m+l} \\
\vdots & \vdots & \vdots & \vdots & & \vdots \\
\gamma_{s+m+l-1} & \gamma_{s+m+l} & \dots & \gamma_{s+m+l+k-3} & & \gamma_{s+m+l+k-2} \\
\hline
\alpha'_s & \alpha'_{s+1} & \dots & \alpha'_{s+k-2} & \alpha_{s+k-1} + \beta_{s+m+k-1} + \gamma_{s+m+l+k-1} &
\end{array} \right) \\
& = \text{rank} \left(\begin{array}{cccccc}
\alpha'_1 & \alpha'_2 & \alpha'_3 & \dots & \alpha'_{k-1} & \alpha'_k \\
\alpha'_2 & \alpha'_3 & \alpha'_4 & \dots & \alpha'_k & \alpha'_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha'_{s-1} & \alpha'_s & \alpha'_{s+1} & \dots & \alpha'_{s+k-3} & \alpha'_{s+k-2} \\
\beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \dots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\gamma_1 & \gamma_2 & \gamma_3 & \dots & \gamma_{k-1} & \gamma_k \\
\gamma_2 & \gamma_3 & \gamma_4 & \dots & \gamma_k & \gamma_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\gamma_{m+l+1} & \gamma_{m+l+2} & \gamma_{m+l+3} & \dots & \gamma_{k+m+l-1} & \gamma_{k+m+l} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\gamma_{s+m+l-1} & \gamma_{s+m+l} & \gamma_{s+m+l+1} & \dots & \gamma_{s+m+l+k-3} & \gamma_{s+m+l+k-2}
\end{array} \right)
\end{aligned}$$

Observing that for all $\alpha'_{s+k-1} \in \mathbb{F}_2$

$$\text{Card} \{ (\alpha_{s+k-1}, \beta_{s+k+m-1}, \gamma_{s+k+m+l-1}) \in \mathbb{F}_2^3 \mid \alpha_{s+k-1} + \beta_{s+k+m-1} + \gamma_{s+k+m+l-1} = \alpha'_{s+k-1} \} = 4,$$

we obtain by combining the above results the desired equality

The other equalities are obtained in a somewhat similarly way. $\square$

2.5. Integration formulas for exponential sums over the unit interval of $\mathbb{K}^3$.

Lemma 2.8. *Let $(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P}$ and q a rational integer ≥ 2 , then we have*

(2.44)

$$\begin{aligned}
& \int_{\mathbb{P}^3} g_1^q(t, \eta, \xi) dt d\eta d\xi = \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid g_1(t, \eta, \xi) \neq 0\}} \varphi_1^q(t, \eta, \xi) dt d\eta d\xi \\
& = 2^{(k+3s+2m+l-1)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-1)} \sigma_{i,i}^{\left[\begin{array}{c} s-1 \\ s+m \\ s+m+l \\ \alpha_{s-} \end{array} \right] \times k} \cdot 2^{-iq}
\end{aligned}$$

$$= 2^{(k+3s+2m+l-1)q-(3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-1)} \left[\sigma_{i,i,i,i} \begin{bmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{bmatrix} \times k + \sigma_{i-1,i,i,i} \begin{bmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{bmatrix} \times k + \sigma_{i-1,i-1,i,i} \begin{bmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{bmatrix} \times k + \sigma_{i-2,i-1,i,i} \begin{bmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{bmatrix} \times k \right] \cdot 2^{-iq}$$

(2.45)

$$\int_{\mathbb{P}^3} h_1^q(t, \eta, \xi) dt d\eta d\xi = \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 | h_1(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi$$

$$= 2^{(k+3s+2m+l-3)q-(3k+3s+2m+l-5)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \sigma_{i,i} \begin{bmatrix} \frac{s-1}{s+m-1} \\ \frac{s+m-1}{s+m+l-1} \end{bmatrix} \times k \cdot 2^{-iq}$$

$$= 2^{(k+3s+2m+l-3)q-(3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \left[\sigma_{i,i,i,i} \begin{bmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k + \sigma_{i,i,i,i+1} \begin{bmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k + \sigma_{i,i,i+1,i+1} \begin{bmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k + \sigma_{i,i,i+1,i+2} \begin{bmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k \right] \cdot 2^{-iq}$$

(2.46)

$$\int_{\mathbb{P}^3} h_2^q(t, \eta, \xi) dt d\eta d\xi = \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 | h_2(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi$$

$$= 2^{(k+3s+2m+l-3)q-(3k+3s+2m+l-4)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \sigma_{i,i} \begin{bmatrix} \frac{s-1}{s+m-1} \\ \frac{s+m-1}{s+m+l-1} \end{bmatrix} \times k \cdot 2^{-iq}$$

$$= 2^{(k+3s+2m+l-3)q-(3k+3s+2m+l-4)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} 2 \cdot \sigma_{i,i} \begin{bmatrix} \frac{s-1}{s+m-1} \\ \frac{s+m-1}{s+m+l-1} \end{bmatrix} \times k \cdot 2^{-iq}$$

$$= 2^{(k+3s+2m+l-3)q-(3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \left[\sigma_{i,i,i,i} \begin{bmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k + \sigma_{i,i,i,i+1} \begin{bmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k + \sigma_{i,i,i+1,i+1} \begin{bmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k + \sigma_{i,i,i+1,i+2} \begin{bmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k \right] \cdot 2^{-iq}$$

(2.47)

$$\int_{\mathbb{P}^3} h_3^q(t, \eta, \xi) dt d\eta d\xi = \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 | h_3(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi$$

$$= 2^{(k+3s+2m+l-3)q-(3k+3s+2m+l-4)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \sigma_{i,i} \begin{bmatrix} \frac{s-1}{s+m-1} \\ \frac{s+m-1}{s+m+l-1} \end{bmatrix} \times k \cdot 2^{-iq}$$

$$= 2^{(k+3s+2m+l-3)q-(3k+3s+2m+l-4)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} 2 \cdot \sigma_{i,i} \begin{bmatrix} \frac{s-1}{s+m-1} \\ \frac{s+m-1}{s+m+l-1} \end{bmatrix} \times k \cdot 2^{-iq}$$

$$= 2^{(k+3s+2m+l-3)q-(3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \left[\sigma_{i,i,i,i} \begin{bmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k + \sigma_{i,i,i,i+1} \begin{bmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k + \sigma_{i,i,i+1,i+1} \begin{bmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k + \sigma_{i,i,i+1,i+2} \begin{bmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k \right] \cdot 2^{-iq}$$

(2.48)

$$\int_{\mathbb{P}^3} h_4^q(t, \eta, \xi) dt d\eta d\xi = \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 | h_4(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi$$

$$= 2^{(k+3s+2m+l-3)q-(3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \sigma_{i,i} \begin{bmatrix} \frac{s-1}{s+m-1} \\ \frac{s+m-1}{s+m+l-1} \end{bmatrix} \times k \cdot 2^{-iq}$$

$$\begin{aligned}
&= 2^{(k+3s+2m+l-3)q-(3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} 4 \cdot \sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \\ \alpha s- \end{smallmatrix} \right] \times k} \cdot 2^{-iq} \\
&= 2^{(k+3s+2m+l-3)q-(3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \left[\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \alpha | \alpha \\ \beta | \gamma \end{smallmatrix} \right] \times k} + \sigma_{i,i,i,i+1}^{\left[\begin{smallmatrix} \alpha | \alpha \\ \beta | \gamma \end{smallmatrix} \right] \times k} + \sigma_{i,i,i+1,i+1}^{\left[\begin{smallmatrix} \alpha | \alpha \\ \beta | \gamma \end{smallmatrix} \right] \times k} + \sigma_{i,i,i+1,i+2}^{\left[\begin{smallmatrix} \alpha | \alpha \\ \beta | \gamma \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq}
\end{aligned}$$

(2.49)

$$\begin{aligned}
\int_{\mathbb{P}^3} \kappa_1^q(t, \eta, \xi) dt d\eta d\xi &= \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 | \kappa_1(t, \eta, \xi) \neq 0\}} \varphi_2^q(t, \eta, \xi) dt d\eta d\xi \\
&= 2^{(k+3s+2m+l-2)q-(3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-2)} \sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \\ \alpha s- \end{smallmatrix} \right] \times k} \cdot 2^{-iq} \\
&= 2^{(k+3s+2m+l-2)q-(3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-2)} 2 \cdot \sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \\ \alpha s- \end{smallmatrix} \right] \times k} \cdot 2^{-iq} \\
&= 2^{(k+3s+2m+l-2)q-(3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-2)} \left[\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} + \sigma_{i,i,i,i+1}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} + \sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} + \sigma_{i-1,i,i,i+1}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq}
\end{aligned}$$

(2.50)

$$\begin{aligned}
\int_{\mathbb{P}^3} \kappa_2^q(t, \eta, \xi) dt d\eta d\xi &= \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 | \kappa_2(t, \eta, \xi) \neq 0\}} \varphi_2^q(t, \eta, \xi) dt d\eta d\xi \\
&= 2^{(k+3s+2m+l-2)q-(3k+3s+2m+l-4)} \sum_{i=0}^{\inf(k, 3s+2m+l-2)} \sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l \\ \alpha s- \end{smallmatrix} \right] \times k} \cdot 2^{-iq} \\
&= 2^{(k+3s+2m+l-2)q-(3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-2)} 2 \cdot \sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l \\ \alpha s- \end{smallmatrix} \right] \times k} \cdot 2^{-iq} \\
&= 2^{(k+3s+2m+l-2)q-(3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-2)} \left[\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} + \sigma_{i,i,i,i+1}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} + \sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} + \sigma_{i-1,i,i,i+1}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq}
\end{aligned}$$

(2.51)

$$\begin{aligned}
\int_{\mathbb{P}^3} g_2^q(t, \eta, \xi) dt d\eta d\xi &= \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 | g_2(t, \eta, \xi) \neq 0\}} \varphi_3^q(t, \eta, \xi) dt d\eta d\xi \\
&= 2^{(k+3s+2m+l-2)q-(3k+3s+2m+l-4)} \sum_{i=0}^{\inf(k, 3s+2m+l-2)} \sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ s+m \\ s+m+l-1 \\ \alpha s- \end{smallmatrix} \right] \times k} \cdot 2^{-iq} \\
&= 2^{(k+3s+2m+l-2)q-(3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-2)} 2 \cdot \sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ s+m \\ s+m+l-1 \\ \alpha s- \end{smallmatrix} \right] \times k} \cdot 2^{-iq} \\
&= 2^{(k+3s+2m+l-2)q-(3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-2)} \left[\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \beta \\ \alpha \\ \gamma \end{smallmatrix} \right] \times k} + \sigma_{i,i,i,i+1}^{\left[\begin{smallmatrix} \beta \\ \alpha \\ \gamma \end{smallmatrix} \right] \times k} + \sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \beta \\ \alpha \\ \gamma \end{smallmatrix} \right] \times k} + \sigma_{i-1,i,i,i+1}^{\left[\begin{smallmatrix} \beta \\ \alpha \\ \gamma \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq}
\end{aligned}$$

Proof. To prove (2.44) we have by (2.1) observing that $g_1^q(t, \eta, \xi)$ is constant on cosets of $\mathbb{P}_{k+s-1} \times \mathbb{P}_{k+s+m-1} \times \mathbb{P}_{k+s+m+l-1}$

$$\begin{aligned}
\int_{\mathbb{P}^3} g_1^q(t, \eta, \xi) dt d\eta d\xi &= \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid g_1(t, \eta, \xi) \neq 0\}} \varphi_1^q(t, \eta, \xi) dt d\eta d\xi \\
&= \sum_{\substack{(t, \eta, \xi) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1} \\ r(D \begin{bmatrix} s-1 \\ s+m \\ s+m+l \end{bmatrix} \times k) \\ (t, \eta, \xi) = r(D \begin{bmatrix} s \\ s+m+l \end{bmatrix} \times k) \\ (t, \eta, \xi) \\ \inf(k, 3s+2m+l-1)}} 2^{(k+3s+2m+l-1-r(D \begin{bmatrix} s-1 \\ s+m \\ s+m+l \end{bmatrix} \times k) (t, \eta, \xi))q} \int_{\mathbb{P}_{k+s-1}} dt \int_{\mathbb{P}_{k+s+m-1}} d\eta \int_{\mathbb{P}_{k+s+m+l-1}} d\xi \\
&= \sum_{i=0}^{\inf(k, 3s+2m+l-1)} \sum_{\substack{(t, \eta, \xi) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1} \\ r(D \begin{bmatrix} s-1 \\ s+m \\ s+m+l \end{bmatrix} \times k) \\ (t, \eta, \xi) = r(D \begin{bmatrix} s \\ s+m+l \end{bmatrix} \times k) \\ (t, \eta, \xi) = i}} 2^{(k+3s+2m+l-1-i)q} \int_{\mathbb{P}_{k+s-1}} dt \int_{\mathbb{P}_{k+s+m-1}} d\eta \int_{\mathbb{P}_{k+s+m+l-1}} d\xi \\
&= 2^{(k+3s+2m+l-1)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-1)} \sigma_{i,i} \begin{bmatrix} s-1 \\ s+m \\ s+m+l \\ \alpha s- \end{bmatrix} \times k \cdot 2^{-iq} \\
&= 2^{(k+3s+2m+l-1)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-1)} \left[\sigma_{i,i,i,i} \begin{bmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{bmatrix} \times k + \sigma_{i-1,i,i,i} \begin{bmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{bmatrix} \times k + \sigma_{i-1,i-1,i,i} \begin{bmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{bmatrix} \times k + \sigma_{i-2,i-1,i,i} \begin{bmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{bmatrix} \times k \right] \cdot 2^{-iq}
\end{aligned}$$

The other integral equalities are obtained in a similar way. $\square$

Lemma 2.9. Let $(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P}$, and q a rational integer ≥ 2 , then we have for $1 \leq i \leq q-1$

$$\begin{aligned}
(2.52) \quad \int_{\mathbb{P}^3} h_1^i(t, \eta, \xi) \cdot h_2^{q-i}(t, \eta, \xi) dt d\eta d\xi &= \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid h_1(t, \eta, \xi) \cdot h_2(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi \\
&= 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-4)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \sigma_{i,i,i} \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha s- \end{bmatrix} \times k \cdot 2^{-iq} \\
&= 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \left[\sigma_{i,i,i,i} \begin{bmatrix} \alpha|\beta \\ \beta|\alpha \\ \gamma|\gamma \end{bmatrix} \times k + \sigma_{i,i,i,i+1} \begin{bmatrix} \alpha|\beta \\ \beta|\alpha \\ \gamma|\gamma \end{bmatrix} \times k \right] \cdot 2^{-iq}
\end{aligned}$$

$$\begin{aligned}
(2.53) \quad \int_{\mathbb{P}^3} h_1^i(t, \eta, \xi) \cdot h_3^{q-i}(t, \eta, \xi) dt d\eta d\xi &= \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid h_1(t, \eta, \xi) \cdot h_3(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi \\
&= 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-4)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \sigma_{i,i,i} \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha s- \end{bmatrix} \times k \cdot 2^{-iq} \\
&= 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \left[\sigma_{i,i,i,i} \begin{bmatrix} \alpha|\gamma \\ \gamma|\alpha \\ \beta|\beta \end{bmatrix} \times k + \sigma_{i,i,i,i+1} \begin{bmatrix} \alpha|\gamma \\ \gamma|\alpha \\ \beta|\beta \end{bmatrix} \times k \right] \cdot 2^{-iq}
\end{aligned}$$

$$\begin{aligned}
(2.54) \quad & \int_{\mathbb{P}^3} h_1^i(t, \eta, \xi) \cdot h_4^{q-i}(t, \eta, \xi) dt d\eta d\xi = \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid h_1(t, \eta, \xi) \cdot h_4(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi \\
&= 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \sigma_{i,i,i} \left[\frac{\frac{s-1}{s+m-1}}{\frac{s+m+l-1}{\alpha_{s-}}} \right] \times k \cdot 2^{-iq} \\
&= 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} 2 \cdot \sigma_{i,i,i} \left[\frac{\frac{s-1}{s+m-1}}{\frac{s+m+l-1}{\alpha_{s-}}} \right] \times k \cdot 2^{-iq} \\
&= 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \left[\sigma_{i,i,i,i} \left[\frac{\alpha|\beta}{\beta|\alpha} \right] \times k + \sigma_{i,i,i,i+1} \left[\frac{\alpha|\beta}{\beta|\alpha} \right] \times k \right] \cdot 2^{-iq}
\end{aligned}$$

$$\begin{aligned}
(2.55) \quad & \int_{\mathbb{P}^3} h_2^i(t, \eta, \xi) \cdot h_3^{q-i}(t, \eta, \xi) dt d\eta d\xi = \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid h_2(t, \eta, \xi) \cdot h_3(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi \\
&= 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \sigma_{i,i,i} \left[\frac{\frac{s-1}{s+m-1}}{\frac{\alpha_{s-} + \beta_{s+m-}}{\beta_{s+m-} + \gamma_{s+m+l-}}} \right] \times k \cdot 2^{-iq} \\
&= 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} 2 \cdot \sigma_{i,i,i} \left[\frac{\frac{s-1}{s+m-1}}{\frac{s+m+l-1}{\alpha_{s-}}} \right] \times k \cdot 2^{-iq} \\
&= 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \left[\sigma_{i,i,i,i} \left[\frac{\alpha|\beta}{\beta|\alpha} \right] \times k + \sigma_{i,i,i,i+1} \left[\frac{\alpha|\beta}{\beta|\alpha} \right] \times k \right] \cdot 2^{-iq}
\end{aligned}$$

$$\begin{aligned}
(2.56) \quad & \int_{\mathbb{P}^3} h_2^i(t, \eta, \xi) \cdot h_4^{q-i}(t, \eta, \xi) dt d\eta d\xi = \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid h_2(t, \eta, \xi) \cdot h_4(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi \\
&= 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \sigma_{i,i,i} \left[\frac{\frac{s-1}{s+m-1}}{\frac{\alpha_{s-} + \beta_{s+m-}}{\gamma_{s+m+l-}}} \right] \times k \cdot 2^{-iq} \\
&= 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} 2 \cdot \sigma_{i,i,i} \left[\frac{\frac{s-1}{s+m-1}}{\frac{s+m+l-1}{\alpha_{s-}}} \right] \times k \cdot 2^{-iq} \\
&= 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \left[\sigma_{i,i,i,i} \left[\frac{\alpha|\gamma}{\gamma|\alpha} \right] \times k + \sigma_{i,i,i,i+1} \left[\frac{\alpha|\gamma}{\gamma|\alpha} \right] \times k \right] \cdot 2^{-iq}
\end{aligned}$$

$$(2.57) \quad \int_{\mathbb{P}^3} h_3^i(t, \eta, \xi) \cdot h_4^{q-i}(t, \eta, \xi) dt d\eta d\xi = \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid h_3(t, \eta, \xi) \cdot h_4(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi$$

$$\begin{aligned}
&= 2^{(k+3s+2m+l-3)q-(3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \sigma_{i,i,i} \left[\frac{\frac{s-1}{s+m-1}}{\frac{\alpha_{s-} + \gamma_{s+m+l-}}{\beta_{s+m-}}} \right] \times k \cdot 2^{-iq} \\
&= 2^{(k+3s+2m+l-3)q-(3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} 2 \cdot \sigma_{i,i,i} \left[\frac{\frac{s-1}{s+m-1}}{\frac{\alpha_{s-}}{\beta_{s+m-}}} \right] \times k \cdot 2^{-iq} \\
&= 2^{(k+3s+2m+l-3)q-(3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \left[\sigma_{i,i,i} \left[\frac{\alpha|\beta}{\beta|\alpha} \right] \times k + \sigma_{i,i,i+1} \left[\frac{\alpha|\beta}{\beta|\alpha} \right] \times k \right] \cdot 2^{-iq} \\
(2.58) \quad &\int_{\mathbb{P}^3} \kappa_1^i(t, \eta, \xi) \cdot \kappa_2^{q-i}(t, \eta, \xi) dt d\eta d\xi = \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid \kappa_1(t, \eta, \xi) \cdot \kappa_2(t, \eta, \xi) \neq 0\}} \varphi_2^q(t, \eta, \xi) dt d\eta d\xi \\
&= 2^{(k+3s+2m+l-2)q-(3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-2)} \sigma_{i,i,i} \left[\frac{\frac{s-1}{s+m-1}}{\frac{\alpha_{s-} + \beta_{s+m-}}{\alpha_{s-}}} \right] \times k \cdot 2^{-iq} \\
&= 2^{(k+3s+2m+l-2)q-(3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-2)} \left[\sigma_{i,i,i,i} \left[\frac{\gamma|\gamma}{\beta|\beta} \right] \times k + \sigma_{i-1,i,i,i} \left[\frac{\gamma|\gamma}{\beta|\beta} \right] \times k \right] \cdot 2^{-iq}
\end{aligned}$$

Proof. Proceed as in the proof of (2.44) using the results in Lemma 2.4 and Lemma 2.7. $\square$

Lemma 2.10. *Let $(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P}$ and q a rational integer ≥ 3 , then we have for any integers i, j, r such that*

$$\begin{cases} i + j + r = q \\ 1 \leq i, j, r \leq q - 2 \end{cases}$$

the following integral identities :

(2.59)

$$\begin{aligned}
&\int_{\mathbb{P}^3} h_1^i(t, \eta, \xi) \cdot h_2^j(t, \eta, \xi) \cdot h_3^r(t, \eta, \xi) dt d\eta d\xi = \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid h_1(t, \eta, \xi) \cdot h_2(t, \eta, \xi) \cdot h_3(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi \\
&= \int_{\mathbb{P}^3} h_1^i(t, \eta, \xi) \cdot h_3^j(t, \eta, \xi) \cdot h_4^r(t, \eta, \xi) dt d\eta d\xi = \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid h_1(t, \eta, \xi) \cdot h_3(t, \eta, \xi) \cdot h_4(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi \\
&= \int_{\mathbb{P}^3} h_2^i(t, \eta, \xi) \cdot h_3^j(t, \eta, \xi) \cdot h_4^r(t, \eta, \xi) dt d\eta d\xi = \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid h_2(t, \eta, \xi) \cdot h_3(t, \eta, \xi) \cdot h_4(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi \\
&= \int_{\mathbb{P}^3} h_1^i(t, \eta, \xi) \cdot h_2^j(t, \eta, \xi) \cdot h_4^r(t, \eta, \xi) dt d\eta d\xi = \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid h_1(t, \eta, \xi) \cdot h_2(t, \eta, \xi) \cdot h_4(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi \\
&= 2^{(k+3s+2m+l-3)q-(3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \sigma_{i,i,i,i} \left[\frac{\frac{s-1}{s+m-1}}{\frac{\alpha_{s-}}{\beta_{s+m-}}} \right] \times k \cdot 2^{-iq}
\end{aligned}$$

$$= 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \alpha \\ \beta \\ \gamma \end{smallmatrix} \right] \times k} \cdot 2^{-iq}$$

Proof. Proceed as in the proof of (2.44) using the results in Lemma 2.5. $\square$

Lemma 2.11. *Let $(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P}$, and q a rational integer ≥ 4 , then we have for any integers i, j, r, p such that*

$$\begin{cases} i + j + r + p = q \\ 1 \leq i, j, r, p \leq q - 3 \end{cases}$$

the following integral identities :

(2.60)

$$\begin{aligned} \int_{\mathbb{P}^3} h_1^i(t, \eta, \xi) \cdot h_2^j(t, \eta, \xi) \cdot h_3^r(t, \eta, \xi) \cdot h_4^p(t, \eta, \xi) dt d\eta d\xi &= \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid h_1(t, \eta, \xi) \cdot h_2(t, \eta, \xi) \cdot h_3(t, \eta, \xi) \cdot h_4(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi \\ &= 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \sigma_{i,i,i,i}^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-m} \\ \beta_{s+m} \\ \gamma_{s+m+l-} \end{smallmatrix} \right] \times k} \cdot 2^{-iq} \\ &= 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \alpha \\ \beta \\ \gamma \end{smallmatrix} \right] \times k} \cdot 2^{-iq} \end{aligned}$$

Proof. Proceed as in the proof of (2.44) using the results in Lemma 2.6. $\square$

2.6. Some rank formulas for partitions of triple persymmetric matrices.

Lemma 2.12. *We can write :*

$$(2.61) \quad g_1(t, \eta, \xi) = \kappa_1(t, \eta, \xi) + \kappa_2(t, \eta, \xi)$$

Let q be a rational integer ≥ 2 , then we obtain from the binomial formula

$$(2.62) \quad g_1^q = (\kappa_1 + \kappa_2)^q = \kappa_1^q + \kappa_2^q + \sum_{i=1}^{q-1} \binom{q}{i} \kappa_1^i \kappa_2^{q-i}$$

By integrating (2.62) on the unit interval of $\mathbb{K}^3$ we get

$$(2.63) \quad \int_{\mathbb{P}^3} g_1^q(t, \eta, \xi) dt d\eta d\xi = 2 \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid \kappa_1(t, \eta, \xi) \neq 0\}} \varphi_2^q(t, \eta, \xi) dt d\eta d\xi + (2^q - 2) \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid \kappa_1(t, \eta, \xi) \cdot \kappa_2(t, \eta, \xi) \neq 0\}} \varphi_2^q(t, \eta, \xi) dt d\eta d\xi$$

Proof. From Lemma 2.2 we get easily (2.61), indeed we have

$$\begin{aligned} g_1(t, \eta, \xi) &= \sum_{\deg Y \leq k-1} \sum_{\deg Z = s-1} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V \leq s+m+l-1} E(\xi YV) \\ &= \sum_{\deg Y \leq k-1} E(tYT^{s-1}) E(\eta YT^{s+m-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-1} E(\xi YV) \\ &+ \sum_{\deg Y \leq k-1} E(tYT^{s-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-1} E(\xi YV) \end{aligned}$$

$$= \kappa_1(t, \eta, \xi) + \kappa_2(t, \eta, \xi).$$

To prove (2.63) we proceed as follows

From (2.49),(2.50) we get

$$(2.64) \quad \int_{\mathbb{P}^3} (\kappa_1^q(t, \eta, \xi) + \kappa_2^q(t, \eta, \xi)) dt d\eta d\xi = 2 \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid \kappa_1(t, \eta, \xi) \neq 0\}} \varphi_2^q(t, \eta, \xi) dt d\eta d\xi$$

By (2.58) we get

$$(2.65) \quad \int_{\mathbb{P}^3} \sum_{i=1}^{q-1} \binom{q}{i} \kappa_1^i \kappa_2^{q-i} dt d\eta d\xi = \sum_{i=1}^{q-1} \binom{q}{i} \int_{\mathbb{P}^3} \kappa_1^i \kappa_2^{q-i} dt d\eta d\xi = (2^q - 2) \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid \kappa_1(t, \eta, \xi) \cdot \kappa_2(t, \eta, \xi) \neq 0\}} \varphi_2^q(t, \eta, \xi) dt d\eta d\xi,$$

(2.63) follows now from (2.64) and (2.65). $\square$

Lemma 2.13. *We have the following rank formula :*

$$(2.66) \quad \boxed{\sigma_{i-1, i-1, i, i}^{\left[\begin{smallmatrix} \gamma \\ \beta \\ \alpha \end{smallmatrix} \right] \times k} + \sigma_{i-2, i-1, i, i}^{\left[\begin{smallmatrix} \gamma \\ \beta \\ \alpha \end{smallmatrix} \right] \times k} = 2 \cdot \left[\sigma_{i-1, i-1, i-1, i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} + \sigma_{i-2, i-1, i-1, i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} \right] \text{ for } 0 \leq i \leq \inf(k, 3s + 2m + l - 1)}$$

Proof. We obtain by combining (2.44), (2.49),(2.58) and (2.62).

$$\begin{aligned} \int_{\mathbb{P}^3} g_1^q(t, \eta, \xi) dt d\eta d\xi &= 2 \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid \kappa_1(t, \eta, \xi) \neq 0\}} \varphi_2^q(t, \eta, \xi) dt d\eta d\xi + (2^q - 2) \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid \kappa_1(t, \eta, \xi) \cdot \kappa_2(t, \eta, \xi) \neq 0\}} \varphi_2^q(t, \eta, \xi) dt d\eta d\xi \\ &\iff \\ 2^{(k+3s+2m+l-1)q - (3k+3s+2m+l-3)} &\sum_{i=0}^{\inf(k, 3s+2m+l-1)} \left[\sigma_{i, i, i, i}^{\left[\begin{smallmatrix} \beta | \gamma \\ \gamma | \beta \\ \alpha | \alpha \end{smallmatrix} \right] \times k} + \sigma_{i-1, i, i, i}^{\left[\begin{smallmatrix} \beta | \gamma \\ \gamma | \beta \\ \alpha | \alpha \end{smallmatrix} \right] \times k} + \sigma_{i-1, i-1, i, i}^{\left[\begin{smallmatrix} \beta | \gamma \\ \gamma | \beta \\ \alpha | \alpha \end{smallmatrix} \right] \times k} + \sigma_{i-2, i-1, i, i}^{\left[\begin{smallmatrix} \beta | \gamma \\ \gamma | \beta \\ \alpha | \alpha \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq} \\ &= 2 \cdot 2^{(k+3s+2m+l-2)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-2)} \left[\sigma_{i, i, i, i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} + \sigma_{i, i, i, i+1}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} + \sigma_{i-1, i, i, i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} + \sigma_{i-1, i, i, i+1}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq} \\ &+ (2^q - 2) \cdot 2^{(k+3s+2m+l-2)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-2)} \left[\sigma_{i, i, i, i}^{\left[\begin{smallmatrix} \gamma | \gamma \\ \alpha | \beta \\ \beta | \alpha \end{smallmatrix} \right] \times k} + \sigma_{i-1, i, i, i}^{\left[\begin{smallmatrix} \gamma | \gamma \\ \alpha | \beta \\ \beta | \alpha \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq} \end{aligned}$$

The case $k \geq 3s + 2m + l - 1$

(2.62) $\iff$

$$\begin{aligned}
& \sum_{i=0}^{3s+2m+l-1} \left[\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix} \right] \times k} + \sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix} \right] \times k} + \sigma_{i-1,i-1,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix} \right] \times k} + \sigma_{i-2,i-1,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq} \\
&= 2 \cdot 2^{-iq} \sum_{i=0}^{3s+2m+l-2} \left[\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} + \sigma_{i,i,i,i+1}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} + \sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} + \sigma_{i-1,i,i,i+1}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq} + (2^q - 2) \cdot 2^{-q} \sum_{i=0}^{3s+2m+l-2} \left[\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \gamma|\gamma \\ \alpha|\beta \\ \beta|\alpha \end{smallmatrix} \right] \times k} + \sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \gamma|\gamma \\ \alpha|\beta \\ \beta|\alpha \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq} \\
&\iff \\
& \sum_{i=0}^{3s+2m+l-1} \left[\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix} \right] \times k} + \sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix} \right] \times k} + \sigma_{i-1,i-1,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix} \right] \times k} + \sigma_{i-2,i-1,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq} \\
&= \sum_{i=1}^{3s+2m+l-1} 2 \cdot \left[\sigma_{i-1,i-1,i-1,i-1}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} + \sigma_{i-1,i-1,i-1,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} + \sigma_{i-2,i-1,i-1,i-1}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} + \sigma_{i-2,i-1,i-1,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq} \\
&+ \sum_{i=0}^{3s+2m+l-2} \left[\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \gamma|\gamma \\ \alpha|\beta \\ \beta|\alpha \end{smallmatrix} \right] \times k} + \sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \gamma|\gamma \\ \alpha|\beta \\ \beta|\alpha \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq} - \sum_{i=1}^{3s+2m+l-1} 2 \cdot \left[\sigma_{i-1,i-1,i-1,i-1}^{\left[\begin{smallmatrix} \gamma|\gamma \\ \alpha|\beta \\ \beta|\alpha \end{smallmatrix} \right] \times k} + \sigma_{i-2,i-1,i-1,i-1}^{\left[\begin{smallmatrix} \gamma|\gamma \\ \alpha|\beta \\ \beta|\alpha \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq} \\
&\implies \\
& \sum_{i=2}^{3s+2m+l-2} \left(\sigma_{i-1,i-1,i,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} + \sigma_{i-2,i-1,i,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} - 2 \cdot \left[\sigma_{i-1,i-1,i-1,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} + \sigma_{i-2,i-1,i-1,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} \right] \right) \cdot 2^{-iq} = 0 \quad \text{for all } q \geq 2 \\
& \left(\sigma_{3s+2m+l-3,3s+2m+l-2,3s+2m+l-1,3s+2m+l-1}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} - 2 \cdot \sigma_{3s+2m+l-3,3s+2m+l-2,3s+2m+l-2,3s+2m+l-1}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} \right) \cdot 2^{-(3s+2m+l-1)q} = 0 \quad \text{for all } q \geq 2 \\
&\implies \\
& \sigma_{i-1,i-1,i,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} + \sigma_{i-2,i-1,i,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} - 2 \cdot \left[\sigma_{i-1,i-1,i-1,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} + \sigma_{i-2,i-1,i-1,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} \right] = 0 \quad \text{for } 2 \leq i \leq 3s + 2m + l - 2 \\
& \sigma_{3s+2m+l-3,3s+2m+l-2,3s+2m+l-1,3s+2m+l-1}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} - 2 \cdot \sigma_{3s+2m+l-3,3s+2m+l-2,3s+2m+l-2,3s+2m+l-1}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix} \right] \times k} = 0 \\
&\quad \square
\end{aligned}$$

Lemma 2.14. *We can write :*

$$(2.67) \quad \kappa_2(t, \eta, \xi) = h_1(t, \eta, \xi) + h_3(t, \eta, \xi)$$

Let q be a rational integer ≥ 2 , then we obtain from the binomial formula

$$(2.68) \quad \kappa_2^q = (h_1 + h_3)^q = h_1^q + h_3^q + \sum_{i=1}^{q-1} \binom{q}{i} h_1^i h_3^{q-i}$$

By integrating (2.68) on the unit interval of $\mathbb{K}^3$ we get

$$(2.69) \quad \int_{\mathbb{P}^3} \kappa_2^q(t, \eta, \xi) dt d\eta d\xi = 2 \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid h_1(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi + (2^q - 2) \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid h_1(t, \eta, \xi) \cdot h_3(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi$$

Proof. Similar to the proof of Lemma 2.12, using Lemma 2.2, (2.45), (2.47) and (2.53). $\square$

Lemma 2.15. *We have the following rank formula :*

(2.70)

$$\sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix}\right] \times k} + \sigma_{i-1,i,i,i+1}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix}\right] \times k} = 2 \cdot \left[\sigma_{i-1,i-1,i,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix}\right] \times k} + \sigma_{i-1,i-1,i,i+1}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix}\right] \times k} \right] \quad \text{for } 0 \leq i \leq \inf(k, 3s+2m+l-2)$$

Proof. We proceed as in Lemma 2.13 by combining (2.50), (2.45), (2.47) and (2.53).

$$\begin{aligned} \int_{\mathbb{P}^3} \kappa_2^q(t, \eta, \xi) dt d\eta d\xi &= 2 \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid h_1(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi + (2^q - 2) \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid h_1(t, \eta, \xi) \cdot h_3(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi \\ &\iff \\ &2^{(k+3s+2m+l-2)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-2)} \left[\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix}\right] \times k} + \sigma_{i,i,i,i+1}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix}\right] \times k} + \sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix}\right] \times k} + \sigma_{i-1,i,i,i+1}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix}\right] \times k} \right] \cdot 2^{-iq} \\ &= 2 \cdot 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \left[\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \alpha | \alpha \\ \beta | \gamma \\ \gamma | \beta \end{smallmatrix}\right] \times k} + \sigma_{i,i,i,i+1}^{\left[\begin{smallmatrix} \alpha | \alpha \\ \beta | \gamma \\ \gamma | \beta \end{smallmatrix}\right] \times k} + \sigma_{i,i,i+1,i+1}^{\left[\begin{smallmatrix} \alpha | \alpha \\ \beta | \gamma \\ \gamma | \beta \end{smallmatrix}\right] \times k} + \sigma_{i,i,i+1,i+2}^{\left[\begin{smallmatrix} \alpha | \alpha \\ \beta | \gamma \\ \gamma | \beta \end{smallmatrix}\right] \times k} \right] \cdot 2^{-iq} \\ &+ (2^q - 2) \cdot 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \left[\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \alpha | \gamma \\ \gamma | \alpha \\ \beta | \beta \end{smallmatrix}\right] \times k} + \sigma_{i,i,i,i+1}^{\left[\begin{smallmatrix} \alpha | \gamma \\ \gamma | \alpha \\ \beta | \beta \end{smallmatrix}\right] \times k} \right] \cdot 2^{-iq} \\ &\iff \\ &\sum_{i=0}^{\inf(k, 3s+2m+l-2)} \left[\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix}\right] \times k} + \sigma_{i,i,i,i+1}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix}\right] \times k} + \sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix}\right] \times k} + \sigma_{i-1,i,i,i+1}^{\left[\begin{smallmatrix} \gamma \\ \alpha \\ \beta \end{smallmatrix}\right] \times k} \right] \cdot 2^{-iq} \\ &= \sum_{i=1}^{\inf(k, 3s+2m+l-3)+1} 2 \cdot \left[\sigma_{i-1,i-1,i-1,i-1}^{\left[\begin{smallmatrix} \alpha | \alpha \\ \beta | \gamma \\ \gamma | \beta \end{smallmatrix}\right] \times k} + \sigma_{i-1,i-1,i-1,i}^{\left[\begin{smallmatrix} \alpha | \alpha \\ \beta | \gamma \\ \gamma | \beta \end{smallmatrix}\right] \times k} + \sigma_{i-1,i-1,i,i}^{\left[\begin{smallmatrix} \alpha | \alpha \\ \beta | \gamma \\ \gamma | \beta \end{smallmatrix}\right] \times k} + \sigma_{i-1,i-1,i,i+1}^{\left[\begin{smallmatrix} \alpha | \alpha \\ \beta | \gamma \\ \gamma | \beta \end{smallmatrix}\right] \times k} \right] \cdot 2^{-iq} \\ &+ \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \left[\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \alpha | \gamma \\ \gamma | \alpha \\ \beta | \beta \end{smallmatrix}\right] \times k} + \sigma_{i,i,i,i+1}^{\left[\begin{smallmatrix} \alpha | \gamma \\ \gamma | \alpha \\ \beta | \beta \end{smallmatrix}\right] \times k} \right] \cdot 2^{-iq} - \sum_{i=1}^{\inf(k, 3s+2m+l-3)+1} 2 \cdot \left[\sigma_{i-1,i-1,i-1,i-1}^{\left[\begin{smallmatrix} \alpha | \gamma \\ \gamma | \alpha \\ \beta | \beta \end{smallmatrix}\right] \times k} + \sigma_{i-1,i-1,i-1,i}^{\left[\begin{smallmatrix} \alpha | \gamma \\ \gamma | \alpha \\ \beta | \beta \end{smallmatrix}\right] \times k} \right] \cdot 2^{-iq} \\ &\quad \square \end{aligned}$$

Lemma 2.16. *We can write :*

$$(2.71) \quad g_2(t, \eta, \xi) = h_1(t, \eta, \xi) + h_2(t, \eta, \xi)$$

Let q be a rational integer ≥ 2 , then we obtain from the binomial formula

$$(2.72) \quad g_2^q = (h_1 + h_2)^q = h_1^q + h_2^q + \sum_{i=1}^{q-1} \binom{q}{i} h_1^i h_2^{q-i}$$

By integrating (2.72) on the unit interval of $\mathbb{K}^3$ we get

(2.73)

$$\int_{\mathbb{P}^3} g_2^q(t, \eta, \xi) dt d\eta d\xi = 2 \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid h_1(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi + (2^q - 2) \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid h_1(t, \eta, \xi) \cdot h_2(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi$$

Proof. Similar to the proof of Lemma 2.12, using Lemma 2.2, (2.45), (2.46) and (2.52). $\square$

Lemma 2.17. *We have the following rank formula :*

(2.74)

$$\sigma_{i-1,i,i,i}^{\begin{bmatrix} \beta \\ \alpha \\ \gamma \end{bmatrix} \times k} + \sigma_{i-1,i,i,i+1}^{\begin{bmatrix} \beta \\ \alpha \\ \gamma \end{bmatrix} \times k} = 2 \cdot \left[\sigma_{i-1,i-1,i,i}^{\begin{bmatrix} \beta \\ \alpha \\ \gamma \end{bmatrix} \times k} + \sigma_{i-1,i-1,i,i+1}^{\begin{bmatrix} \beta \\ \alpha \\ \gamma \end{bmatrix} \times k} \right] \quad \text{for } 0 \leq i \leq \inf(k, 3s+2m+l-2)$$

Proof. We proceed as in Lemma 2.13 by combining (2.51), (2.45), (2.46) and (2.52).

$$\begin{aligned} \int_{\mathbb{P}^3} g_2^q(t, \eta, \xi) dt d\eta d\xi &= 2 \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid h_1(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi + (2^q - 2) \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 \mid h_1(t, \eta, \xi) \cdot h_2(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi \\ &\iff \\ &2^{(k+3s+2m+l-2)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-2)} \left[\sigma_{i,i,i,i}^{\begin{bmatrix} \beta \\ \alpha \\ \gamma \end{bmatrix} \times k} + \sigma_{i,i,i,i+1}^{\begin{bmatrix} \beta \\ \alpha \\ \gamma \end{bmatrix} \times k} + \sigma_{i-1,i,i,i}^{\begin{bmatrix} \beta \\ \alpha \\ \gamma \end{bmatrix} \times k} + \sigma_{i-1,i,i,i+1}^{\begin{bmatrix} \beta \\ \alpha \\ \gamma \end{bmatrix} \times k} \right] \cdot 2^{-iq} \\ &= 2 \cdot 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \left[\sigma_{i,i,i,i}^{\begin{bmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k} + \sigma_{i,i,i,i+1}^{\begin{bmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k} + \sigma_{i,i,i+1,i+1}^{\begin{bmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k} + \sigma_{i,i,i+1,i+2}^{\begin{bmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k} \right] \cdot 2^{-iq} \\ &+ (2^q - 2) \cdot 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \left[\sigma_{i,i,i,i}^{\begin{bmatrix} \alpha|\beta \\ \beta|\gamma \\ \gamma|\gamma \end{bmatrix} \times k} + \sigma_{i,i,i,i+1}^{\begin{bmatrix} \alpha|\beta \\ \beta|\gamma \\ \gamma|\gamma \end{bmatrix} \times k} \right] \cdot 2^{-iq} \\ &\iff \\ &\inf(k, 3s+2m+l-2) \sum_{i=0}^{\inf(k, 3s+2m+l-2)} \left[\sigma_{i,i,i,i}^{\begin{bmatrix} \beta \\ \alpha \\ \gamma \end{bmatrix} \times k} + \sigma_{i,i,i,i+1}^{\begin{bmatrix} \beta \\ \alpha \\ \gamma \end{bmatrix} \times k} + \sigma_{i-1,i,i,i}^{\begin{bmatrix} \beta \\ \alpha \\ \gamma \end{bmatrix} \times k} + \sigma_{i-1,i,i,i+1}^{\begin{bmatrix} \beta \\ \alpha \\ \gamma \end{bmatrix} \times k} \right] \cdot 2^{-iq} \\ &= \sum_{i=1}^{\inf(k, 3s+2m+l-3)+1} 2 \cdot \left[\sigma_{i-1,i-1,i-1,i-1}^{\begin{bmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k} + \sigma_{i-1,i-1,i-1,i}^{\begin{bmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k} + \sigma_{i-1,i-1,i,i}^{\begin{bmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k} + \sigma_{i-1,i-1,i,i+1}^{\begin{bmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k} \right] \cdot 2^{-iq} \\ &+ \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \left[\sigma_{i,i,i,i}^{\begin{bmatrix} \alpha|\beta \\ \beta|\gamma \\ \gamma|\alpha \end{bmatrix} \times k} + \sigma_{i,i,i,i+1}^{\begin{bmatrix} \alpha|\beta \\ \beta|\gamma \\ \gamma|\alpha \end{bmatrix} \times k} \right] \cdot 2^{-iq} - \sum_{i=1}^{\inf(k, 3s+2m+l-3)+1} 2 \cdot \left[\sigma_{i-1,i-1,i-1,i-1}^{\begin{bmatrix} \alpha|\beta \\ \beta|\gamma \\ \gamma|\alpha \end{bmatrix} \times k} + \sigma_{i-1,i-1,i-1,i}^{\begin{bmatrix} \alpha|\beta \\ \beta|\gamma \\ \gamma|\alpha \end{bmatrix} \times k} \right] \cdot 2^{-iq} \\ &\quad \square \end{aligned}$$

Lemma 2.18. *We can write :*

$$(2.75) \quad g_1(t, \eta, \xi) = h_1(t, \eta, \xi) + h_2(t, \eta, \xi) + h_3(t, \eta, \xi) + h_4(t, \eta, \xi)$$

Let q be a rational integer ≥ 2 , then we obtain from the binomial formula

(2.76)

$$\begin{aligned} g_1^q &= (h_1 + h_2 + h_3 + h_4)^q = (h_1 + h_2)^q + (h_3 + h_4)^q + \sum_{i=1}^{q-1} \binom{q}{i} (h_1 + h_2)^i (h_3 + h_4)^{q-i} \\ &= h_1^q + h_2^q + h_3^q + h_4^q + \sum_{i=1}^{q-1} \binom{q}{i} h_1^i h_2^{q-i} + \sum_{i=1}^{q-1} \binom{q}{i} h_3^i h_4^{q-i} + \sum_{i=1}^{q-1} \binom{q}{i} h_1^i h_3^{q-i} + \sum_{i=1}^{q-1} \binom{q}{i} h_1^i h_4^{q-i} + \sum_{i=1}^{q-1} \binom{q}{i} h_2^i h_3^{q-i} + \sum_{i=1}^{q-1} \binom{q}{i} h_2^i h_4^{q-i} \\ &+ \sum_{i=1}^{q-1} \sum_{j=1}^{i-1} \binom{q}{i} \binom{i}{j} h_1^j h_2^{i-j} h_3^{q-i} + \sum_{i=1}^{q-1} \sum_{j=1}^{i-1} \binom{q}{i} \binom{i}{j} h_1^j h_2^{i-j} h_4^{q-i} + \sum_{i=1}^{q-1} \sum_{r=1}^{q-i-1} \binom{q}{i} \binom{q-i}{r} h_1^i h_3^r h_4^{q-i-r} \\ &+ \sum_{i=1}^{q-1} \sum_{r=1}^{q-i-1} \binom{q}{i} \binom{q-i}{r} h_2^i h_3^r h_4^{q-i-r} + \sum_{i=1}^{q-1} \sum_{j=1}^{i-1} \sum_{r=1}^{q-i-1} \binom{q}{i} \binom{i}{j} \binom{q-i}{r} h_1^j h_2^{i-j} h_3^r h_4^{q-i-r} \end{aligned}$$

By integrating (2.76) on the unit interval of $\mathbb{K}^3$ we get

(2.77)

$$\begin{aligned}
& \int_{\mathbb{P}^3} g_1^q(t, \eta, \xi) dt d\eta d\xi = 4 \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 | h_1(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi + 4(2^q - 2) \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 | h_1(t, \eta, \xi) \cdot h_2(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi \\
& + 2(2^q - 2) \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 | h_1(t, \eta, \xi) \cdot h_3(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi \\
& + (2^{2q} - 6 \cdot 2^q + 8) \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 | h_1(t, \eta, \xi) \cdot h_2(t, \eta, \xi) \cdot h_3(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi \\
& \iff \\
& 2^{(k+3s+2m+l-1)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-1)} \left[\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix} \right] \times k} + \sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix} \right] \times k} + \sigma_{i-1,i-1,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix} \right] \times k} + \sigma_{i-2,i-1,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq} \\
& = 4 \cdot 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \left[\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{smallmatrix} \right] \times k} + \sigma_{i,i,i,i+1}^{\left[\begin{smallmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{smallmatrix} \right] \times k} + \sigma_{i,i,i+1,i+1}^{\left[\begin{smallmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{smallmatrix} \right] \times k} + \sigma_{i,i,i+1,i+2}^{\left[\begin{smallmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq} \\
& + 4 \cdot (2^q - 2) \cdot 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \left[\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \alpha|\beta \\ \gamma|\gamma \end{smallmatrix} \right] \times k} + \sigma_{i,i,i,i+1}^{\left[\begin{smallmatrix} \alpha|\beta \\ \gamma|\gamma \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq} \\
& + 2 \cdot (2^q - 2) \cdot 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \left[\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \alpha|\gamma \\ \gamma|\alpha \\ \beta|\beta \end{smallmatrix} \right] \times k} + \sigma_{i,i,i,i+1}^{\left[\begin{smallmatrix} \alpha|\gamma \\ \gamma|\alpha \\ \beta|\beta \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq} \\
& + (2^{2q} - 6 \cdot 2^q + 8) \cdot 2^{(k+3s+2m+l-3)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \alpha \\ \beta \\ \gamma \end{smallmatrix} \right] \times k} \cdot 2^{-iq} \\
& \iff \\
& \inf(k, 3s+2m+l-1) \sum_{i=0}^{\inf(k, 3s+2m+l-1)} \left[\sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix} \right] \times k} + \sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix} \right] \times k} + \sigma_{i-1,i-1,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix} \right] \times k} + \sigma_{i-2,i-1,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq} \\
& = 4 \cdot \sum_{i=2}^{\inf(k, 3s+2m+l-3)+2} \left[\sigma_{i-2,i-2,i-2,i-2}^{\left[\begin{smallmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{smallmatrix} \right] \times k} + \sigma_{i-2,i-2,i-2,i-1}^{\left[\begin{smallmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{smallmatrix} \right] \times k} + \sigma_{i-2,i-2,i-1,i-1}^{\left[\begin{smallmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{smallmatrix} \right] \times k} + \sigma_{i-2,i-2,i-1,i}^{\left[\begin{smallmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq} \\
& + \sum_{i=1}^{\inf(k, 3s+2m+l-3)+1} \left[4 \cdot \sigma_{i-1,i-1,i-1,i-1}^{\left[\begin{smallmatrix} \alpha|\beta \\ \gamma|\gamma \end{smallmatrix} \right] \times k} + 4 \cdot \sigma_{i-1,i-1,i-1,i}^{\left[\begin{smallmatrix} \alpha|\beta \\ \gamma|\gamma \end{smallmatrix} \right] \times k} - \sum_{i=2}^{\inf(k, 3s+2m+l-3)+2} \left[8 \cdot \sigma_{i-2,i-2,i-2,i-2}^{\left[\begin{smallmatrix} \alpha|\beta \\ \gamma|\gamma \end{smallmatrix} \right] \times k} + 8 \cdot \sigma_{i-2,i-2,i-2,i-1}^{\left[\begin{smallmatrix} \alpha|\beta \\ \gamma|\gamma \end{smallmatrix} \right] \times k} \right] \right] \cdot 2^{-iq} \\
& + \sum_{i=1}^{\inf(k, 3s+2m+l-3)+1} \left[2 \cdot \sigma_{i-1,i-1,i-1,i-1}^{\left[\begin{smallmatrix} \alpha|\gamma \\ \gamma|\beta \\ \beta|\beta \end{smallmatrix} \right] \times k} + 2 \cdot \sigma_{i-1,i-1,i-1,i}^{\left[\begin{smallmatrix} \alpha|\gamma \\ \gamma|\beta \\ \beta|\beta \end{smallmatrix} \right] \times k} - \sum_{i=2}^{\inf(k, 3s+2m+l-3)+2} \left[4 \cdot \sigma_{i-2,i-2,i-2,i-2}^{\left[\begin{smallmatrix} \alpha|\gamma \\ \gamma|\beta \\ \beta|\beta \end{smallmatrix} \right] \times k} + 4 \cdot \sigma_{i-2,i-2,i-2,i-1}^{\left[\begin{smallmatrix} \alpha|\gamma \\ \gamma|\beta \\ \beta|\beta \end{smallmatrix} \right] \times k} \right] \right] \cdot 2^{-iq} \\
& + \sum_{i=0}^{\inf(k, 3s+2m+l-3)} \sigma_{i,i,i,i}^{\left[\begin{smallmatrix} \alpha \\ \beta \\ \gamma \end{smallmatrix} \right] \times k} \cdot 2^{-iq} - 6 \cdot \sum_{i=1}^{\inf(k, 3s+2m+l-3)+1} \sigma_{i-1,i-1,i-1,i-1}^{\left[\begin{smallmatrix} \alpha \\ \beta \\ \gamma \end{smallmatrix} \right] \times k} \cdot 2^{-iq} + 8 \cdot \sum_{i=2}^{\inf(k, 3s+2m+l-3)+2} \sigma_{i-2,i-2,i-2,i-2}^{\left[\begin{smallmatrix} \alpha \\ \beta \\ \gamma \end{smallmatrix} \right] \times k} \cdot 2^{-iq} \\
& \implies
\end{aligned}$$

In the case $k \leq 3s + 2m + l - 3$

$$\sum_{i=2}^{k+1} \left[\sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix} \right] \times k} + \sigma_{i-1,i-1,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix} \right] \times k} + \sigma_{i-2,i-1,i,i}^{\left[\begin{smallmatrix} \beta|\gamma \\ \gamma|\beta \\ \alpha|\alpha \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq} = 4 \cdot \sum_{i=2}^{k+1} \left[\sigma_{i-2,i-2,i-2,i-1}^{\left[\begin{smallmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{smallmatrix} \right] \times k} + \sigma_{i-2,i-2,i-1,i-1}^{\left[\begin{smallmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{smallmatrix} \right] \times k} + \sigma_{i-2,i-2,i-1,i}^{\left[\begin{smallmatrix} \alpha|\alpha \\ \beta|\gamma \\ \gamma|\beta \end{smallmatrix} \right] \times k} \right] \cdot 2^{-iq}$$

$$+ \sum_{i=2}^{k+1} \left[4 \cdot \sigma_{i-1,i-1,i-1,i}^{\begin{bmatrix} \alpha|\beta \\ \gamma|\alpha \end{bmatrix} \times k} - 8 \cdot \sigma_{i-2,i-2,i-2,i-1}^{\begin{bmatrix} \alpha|\beta \\ \gamma|\alpha \end{bmatrix} \times k} \right] \cdot 2^{-iq} + \sum_{i=2}^{k+1} \left[2 \cdot \sigma_{i-1,i-1,i-1,i}^{\begin{bmatrix} \alpha|\gamma \\ \beta|\alpha \end{bmatrix} \times k} - 4 \cdot \sigma_{i-2,i-2,i-2,i-1}^{\begin{bmatrix} \alpha|\gamma \\ \beta|\alpha \end{bmatrix} \times k} \right] \cdot 2^{-iq} \quad \text{for all } q \geq 2$$

In the case $k \geq 3s+2m+l-2$

$$\begin{aligned} & \sum_{i=2}^{3s+2m+l-1} \left[\sigma_{i-1,i,i,i}^{\begin{bmatrix} \beta|\gamma \\ \alpha|\alpha \end{bmatrix} \times k} + \sigma_{i-1,i-1,i,i}^{\begin{bmatrix} \beta|\gamma \\ \gamma|\beta \end{bmatrix} \times k} + \sigma_{i-2,i-1,i,i}^{\begin{bmatrix} \beta|\gamma \\ \alpha|\alpha \end{bmatrix} \times k} \right] \cdot 2^{-iq} = 4 \cdot \sum_{i=2}^{3s+2m+l-1} \left[\sigma_{i-2,i-2,i-2,i-1}^{\begin{bmatrix} \alpha|\gamma \\ \beta|\beta \end{bmatrix} \times k} + \sigma_{i-2,i-2,i-1,i-1}^{\begin{bmatrix} \alpha|\gamma \\ \beta|\beta \end{bmatrix} \times k} + \sigma_{i-2,i-2,i-1,i}^{\begin{bmatrix} \alpha|\gamma \\ \beta|\beta \end{bmatrix} \times k} \right] \cdot 2^{-iq} \\ & + \sum_{i=2}^{3s+2m+l-1} \left[4 \cdot \sigma_{i-1,i-1,i-1,i}^{\begin{bmatrix} \alpha|\beta \\ \gamma|\alpha \end{bmatrix} \times k} - 8 \cdot \sigma_{i-2,i-2,i-2,i-1}^{\begin{bmatrix} \alpha|\beta \\ \gamma|\alpha \end{bmatrix} \times k} \right] \cdot 2^{-iq} + \sum_{i=2}^{3s+2m+l-1} \left[2 \cdot \sigma_{i-1,i-1,i-1,i}^{\begin{bmatrix} \alpha|\gamma \\ \beta|\beta \end{bmatrix} \times k} - 4 \cdot \sigma_{i-2,i-2,i-2,i-1}^{\begin{bmatrix} \alpha|\gamma \\ \beta|\beta \end{bmatrix} \times k} \right] \cdot 2^{-iq} \quad \text{for all } q \geq 2 \end{aligned}$$

Proof. From Lemma 2.2 we get easily (2.75), indeed we have

$$\begin{aligned} g_1(t, \eta, \xi) &= \sum_{\deg Y \leq k-1} \sum_{\deg Z = s-1} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V \leq s+m+l-1} E(\xi YV) \\ &= \sum_{\deg Y \leq k-1} E(tYT^{s-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V \leq s+m+l-1} E(\xi YV) \\ &= \sum_{\deg Y \leq k-1} E(tYT^{s-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \\ &+ \sum_{\deg Y \leq k-1} E(tYT^{s-1}) E(\eta YT^{s+m-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \\ &+ \sum_{\deg Y \leq k-1} E(tYT^{s-1}) E(\xi YT^{s+m+l-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \\ &+ \sum_{\deg Y \leq k-1} E(tYT^{s-1}) E(\eta YT^{s+m-1}) E(\xi YT^{s+m+l-1}) \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \\ &= h_1(t, \eta, \xi) + h_2(t, \eta, \xi) + h_3(t, \eta, \xi) + h_4(t, \eta, \xi). \end{aligned}$$

To prove (2.76) we observe that

$$\sum_{i=1}^{q-1} \binom{q}{i} (h_1 + h_2)^i (h_3 + h_4)^{q-i} = \sum_{i=1}^{q-1} \binom{q}{i} \left[h_1^i + h_2^i + \sum_{j=1}^{i-1} \binom{i}{j} h_1^j h_2^{i-j} \right] \left[h_3^{q-i} + h_4^{q-i} + \sum_{r=1}^{q-i-1} \binom{q-i}{r} h_3^r h_4^{q-i-r} \right].$$

To prove (2.77) we proceed as follows

From (2.45), (2.45), (2.45) and (2.45) we get

$$(2.78) \quad \int_{\mathbb{P}^3} (h_1^q(t, \eta, \xi) + h_2^q(t, \eta, \xi) + h_3^q(t, \eta, \xi) + h_4^q(t, \eta, \xi)) dt d\eta d\xi = 4 \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 | h_1(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi$$

By (2.52), (2.53), (2.54), (2.55), (2.56) and (2.57) we obtain

$$\begin{aligned} (2.79) \quad & \int_{\mathbb{P}^3} \left(\sum_{i=1}^{q-1} \binom{q}{i} h_1^i h_2^{q-i} + \sum_{i=1}^{q-1} \binom{q}{i} h_3^i h_4^{q-i} + \sum_{i=1}^{q-1} \binom{q}{i} h_1^i h_3^{q-i} + \sum_{i=1}^{q-1} \binom{q}{i} h_1^i h_4^{q-i} + \sum_{i=1}^{q-1} \binom{q}{i} h_2^i h_3^{q-i} + \sum_{i=1}^{q-1} \binom{q}{i} h_2^i h_4^{q-i} \right) dt d\eta d\xi \\ &= 4(2^q - 2) \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 | h_1(t, \eta, \xi) \cdot h_2(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi + 2(2^q - 2) \int_{\{(t, \eta, \xi) \in \mathbb{P}^3 | h_1(t, \eta, \xi) \cdot h_3(t, \eta, \xi) \neq 0\}} \varphi^q(t, \eta, \xi) dt d\eta d\xi \end{aligned}$$

From (2.59), (2.60) we get

$$\begin{aligned}
 (2.80) \quad & \int_{\mathbb{P}^3} \left[\sum_{i=1}^{q-1} \sum_{j=1}^{i-1} \binom{q}{i} \binom{i}{j} h_1^j h_2^{i-j} h_3^{q-i} + \sum_{i=1}^{q-1} \sum_{j=1}^{i-1} \binom{q}{i} \binom{i}{j} h_1^j h_2^{i-j} h_4^{q-i} + \sum_{i=1}^{q-1} \sum_{r=1}^{q-i-1} \binom{q}{i} \binom{q-i}{r} h_1^i h_3^r h_4^{q-i-r} \right. \\
 & + \sum_{i=1}^{q-1} \sum_{r=1}^{q-i-1} \binom{q}{i} \binom{q-i}{r} h_2^i h_3^r h_4^{q-i-r} + \sum_{i=1}^{q-1} \sum_{j=1}^{i-1} \sum_{r=1}^{q-i-1} \binom{q}{i} \binom{i}{j} \binom{q-i}{r} h_1^j h_2^{i-j} h_3^r h_4^{q-i-r} \left. \right] dtd\eta d\xi \\
 & = (2^{2q} - 6 \cdot 2^q + 8) \int_{\{(t,\eta,\xi) \in \mathbb{P}^3 \mid h_1(t,\eta,\xi) \cdot h_2(t,\eta,\xi) \cdot h_3(t,\eta,\xi) \neq 0\}} \varphi^q(t, \eta, \xi) dtd\eta d\xi
 \end{aligned}$$

(2.77) follows now from (2.78), (2.79) and (2.80) follows by (2.44), (2.45), (2.52), (2.53) and (2.59). $\square$

Lemma 2.19. *We have the following rank formula :*

$$\begin{aligned}
 (2.81) \quad & \sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \gamma \\ \beta \end{smallmatrix}\right] \times k} + \sigma_{i-1,i-1,i,i}^{\left[\begin{smallmatrix} \gamma \\ \beta \end{smallmatrix}\right] \times k} + \sigma_{i-2,i-1,i,i}^{\left[\begin{smallmatrix} \gamma \\ \beta \end{smallmatrix}\right] \times k} = 4 \cdot \left[\sigma_{i-2,i-2,i-2,i-1}^{\left[\begin{smallmatrix} \alpha \\ \beta \end{smallmatrix}\right] \times k} + \sigma_{i-2,i-2,i-1,i-1}^{\left[\begin{smallmatrix} \alpha \\ \beta \end{smallmatrix}\right] \times k} + \sigma_{i-2,i-2,i-1,i}^{\left[\begin{smallmatrix} \alpha \\ \beta \end{smallmatrix}\right] \times k} \right] \\
 & + \left[4 \cdot \sigma_{i-1,i-1,i-1,i}^{\left[\begin{smallmatrix} \alpha \\ \beta \end{smallmatrix}\right] \times k} - 8 \cdot \sigma_{i-2,i-2,i-2,i-1}^{\left[\begin{smallmatrix} \alpha \\ \beta \end{smallmatrix}\right] \times k} \right] + \left[2 \cdot \sigma_{i-1,i-1,i-1,i}^{\left[\begin{smallmatrix} \gamma \\ \beta \end{smallmatrix}\right] \times k} - 4 \cdot \sigma_{i-2,i-2,i-2,i-1}^{\left[\begin{smallmatrix} \gamma \\ \beta \end{smallmatrix}\right] \times k} \right] \quad \text{for } 1 \leq i \leq \inf(k, 3s + 2m + l - 2) + 1
 \end{aligned}$$

Proof. The rank formula follows from (2.77). $\square$

Lemma 2.20. *We have the following rank formula*

$$(2.82) \quad \sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \gamma \\ \beta \end{smallmatrix}\right] \times k} = 4 \cdot \sigma_{i-1,i-1,i-1,i}^{\left[\begin{smallmatrix} \alpha \\ \beta \end{smallmatrix}\right] \times k} \quad \text{for } 1 \leq i \leq \inf(k, 3s + 2m + l - 2)$$

Proof. We obtain by combining (2.66), (2.70) and (2.81).

$$\begin{aligned}
 & \sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \gamma \\ \beta \end{smallmatrix}\right] \times k} + \sigma_{i-1,i-1,i,i}^{\left[\begin{smallmatrix} \gamma \\ \beta \end{smallmatrix}\right] \times k} + \sigma_{i-2,i-1,i,i}^{\left[\begin{smallmatrix} \gamma \\ \beta \end{smallmatrix}\right] \times k} = 4 \cdot \left[\sigma_{i-2,i-2,i-2,i-1}^{\left[\begin{smallmatrix} \alpha \\ \beta \end{smallmatrix}\right] \times k} + \sigma_{i-2,i-2,i-1,i-1}^{\left[\begin{smallmatrix} \alpha \\ \beta \end{smallmatrix}\right] \times k} + \sigma_{i-2,i-2,i-1,i}^{\left[\begin{smallmatrix} \alpha \\ \beta \end{smallmatrix}\right] \times k} \right] \\
 & + \left[4 \cdot \sigma_{i-1,i-1,i-1,i}^{\left[\begin{smallmatrix} \alpha \\ \beta \end{smallmatrix}\right] \times k} - 8 \cdot \sigma_{i-2,i-2,i-2,i-1}^{\left[\begin{smallmatrix} \alpha \\ \beta \end{smallmatrix}\right] \times k} \right] + \left[2 \cdot \sigma_{i-1,i-1,i-1,i}^{\left[\begin{smallmatrix} \gamma \\ \beta \end{smallmatrix}\right] \times k} - 4 \cdot \sigma_{i-2,i-2,i-2,i-1}^{\left[\begin{smallmatrix} \gamma \\ \beta \end{smallmatrix}\right] \times k} \right] \\
 & \implies \\
 & \sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \gamma \\ \beta \end{smallmatrix}\right] \times k} + 2 \cdot \left[\sigma_{i-1,i-1,i-1,i}^{\left[\begin{smallmatrix} \gamma \\ \beta \end{smallmatrix}\right] \times k} + \sigma_{i-2,i-1,i-1,i}^{\left[\begin{smallmatrix} \gamma \\ \beta \end{smallmatrix}\right] \times k} \right] = 4 \cdot \sigma_{i-2,i-2,i-2,i-1}^{\left[\begin{smallmatrix} \alpha \\ \beta \end{smallmatrix}\right] \times k} + 2 \cdot \left[\sigma_{i-2,i-1,i-1,i-1}^{\left[\begin{smallmatrix} \gamma \\ \beta \end{smallmatrix}\right] \times k} + \sigma_{i-2,i-1,i-1,i}^{\left[\begin{smallmatrix} \gamma \\ \beta \end{smallmatrix}\right] \times k} \right] \\
 & + 4 \cdot \sigma_{i-1,i-1,i-1,i}^{\left[\begin{smallmatrix} \alpha \\ \beta \end{smallmatrix}\right] \times k} - 8 \cdot \sigma_{i-2,i-2,i-2,i-1}^{\left[\begin{smallmatrix} \alpha \\ \beta \end{smallmatrix}\right] \times k} + 2 \cdot \sigma_{i-1,i-1,i-1,i}^{\left[\begin{smallmatrix} \gamma \\ \beta \end{smallmatrix}\right] \times k} - 4 \cdot \sigma_{i-2,i-2,i-2,i-1}^{\left[\begin{smallmatrix} \gamma \\ \beta \end{smallmatrix}\right] \times k} \\
 & \implies \\
 & \sigma_{i-1,i,i,i}^{\left[\begin{smallmatrix} \gamma \\ \beta \end{smallmatrix}\right] \times k} - 4 \cdot \sigma_{i-1,i-1,i-1,i}^{\left[\begin{smallmatrix} \alpha \\ \beta \end{smallmatrix}\right] \times k} = 2 \cdot \left[\sigma_{i-2,i-1,i-1,i-1}^{\left[\begin{smallmatrix} \gamma \\ \beta \end{smallmatrix}\right] \times k} - 4 \cdot \sigma_{i-2,i-2,i-2,i-1}^{\left[\begin{smallmatrix} \alpha \\ \beta \end{smallmatrix}\right] \times k} \right] \quad \text{for } 2 \leq i \leq \inf(k, 3s + 2m + l - 1),
 \end{aligned}$$

which implies

$$\sigma_{i-1,i,i,i}^{\begin{bmatrix} \gamma \\ \beta \\ \alpha \end{bmatrix} \times k} - 4 \cdot \sigma_{i-1,i-1,i-1,i}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} = 2^{i-1} \cdot [\sigma_{0,1,1,1}^{\begin{bmatrix} \gamma \\ \alpha \\ \beta \end{bmatrix} \times k} - 4 \cdot \sigma_{0,0,0,1}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k}] = 0 \quad \text{for } 1 \leq i \leq \inf(k, 3s+2m+l-2)$$

□

2.7. A recurrent formula for the number of rank i matrices of the form $\begin{bmatrix} \frac{A}{B} \\ \frac{C}{C} \end{bmatrix}$, where A, B and C are persymmetric matrices over $\mathbb{F}_2$.

Lemma 2.21. *Let $s \geq 2$, $m \geq 0$, $l \geq 0$, $k \geq 1$ and $0 \leq i \leq \inf(3s+2m+l, k)$. Then we*

have the following recurrent formula for the number $\Gamma_i^{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k}$ of rank i matrices of the form $\begin{bmatrix} \frac{A}{B} \\ \frac{C}{C} \end{bmatrix}$ such that A is a $s \times k$ persymmetric matrix over $\mathbb{F}_2$, B a $(s+m) \times k$ persymmetric matrix and C a $(s+m+l) \times k$ persymmetric matrix

$$(2.83) \quad \begin{aligned} & \Gamma_i^{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k} \\ &= \left[2 \cdot \Gamma_{i-1}^{\begin{bmatrix} s-1 \\ s+m \\ s+m+l \end{bmatrix} \times k} + 4 \cdot \Gamma_{i-1}^{\begin{bmatrix} s \\ s+m-1 \\ s+m+l \end{bmatrix} \times k} + 8 \cdot \Gamma_{i-1}^{\begin{bmatrix} s \\ s+m \\ s+m+l-1 \end{bmatrix} \times k} - \left[8 \cdot \Gamma_{i-2}^{\begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l \end{bmatrix} \times k} + 16 \cdot \Gamma_{i-2}^{\begin{bmatrix} s-1 \\ s+m \\ s+m+l-1 \end{bmatrix} \times k} + 32 \cdot \Gamma_{i-2}^{\begin{bmatrix} s \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k} \right] \right. \\ & \left. + 64 \cdot \Gamma_{i-3}^{\begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k} + \Delta_i^{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k} \right] \end{aligned}$$

where

$$(2.84) \quad \Delta_i^{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k} = \sigma_{i,i,i,i}^{\begin{bmatrix} s \\ \gamma \\ \beta \end{bmatrix} \times k} - 7 \cdot \sigma_{i-1,i-1,i-1,i-1}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} + 14 \cdot \sigma_{i-2,i-2,i-2,i-2}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} - 8 \cdot \sigma_{i-3,i-3,i-3,i-3}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k}$$

Recall that

$$\sigma_{i,i,i,i}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k}$$

denotes the cardinality of the following set

$$\left\{ (t, \eta, \xi) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1} \mid r(D^{\begin{bmatrix} s-1 \\ s+m-1 \end{bmatrix} \times k}(t, \eta, \xi)) = i \right. \\ \left. r(D^{\begin{bmatrix} s+m-1 \\ s+m+l-1 \end{bmatrix} \times k}(t, \eta, \xi)) = i, \quad r(D^{\begin{bmatrix} s+m \\ s+m+l-1 \end{bmatrix} \times k}(t, \eta, \xi)) = i, \quad r(D^{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k}(t, \eta, \xi)) = i \right\}$$

Proof. We have obviously

•

$$\begin{aligned} & \Gamma_i^{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k} \\ &= \sigma_{i,i,i,i}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} + \sigma_{i-1,i,i,i}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} + \sigma_{i-2,i-1,i,i}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} + \sigma_{i-1,i-1,i,i}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} \\ &+ \sigma_{i-1,i-1,i-1,i}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} + \sigma_{i-2,i-1,i-1,i}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} + \sigma_{i-2,i-2,i-1,i}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} + \sigma_{i-3,i-2,i-1,i}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} \\ &= \sigma_{i,i,i,i}^{\begin{bmatrix} \alpha \\ \gamma \\ \beta \end{bmatrix} \times k} + \sigma_{i-1,i,i,i}^{\begin{bmatrix} \alpha \\ \gamma \\ \beta \end{bmatrix} \times k} + \sigma_{i-2,i-1,i,i}^{\begin{bmatrix} \alpha \\ \gamma \\ \beta \end{bmatrix} \times k} + \sigma_{i-1,i-1,i,i}^{\begin{bmatrix} \alpha \\ \gamma \\ \beta \end{bmatrix} \times k} \\ &+ \sigma_{i-1,i-1,i-1,i}^{\begin{bmatrix} \alpha \\ \gamma \\ \beta \end{bmatrix} \times k} + \sigma_{i-2,i-1,i-1,i}^{\begin{bmatrix} \alpha \\ \gamma \\ \beta \end{bmatrix} \times k} + \sigma_{i-2,i-2,i-1,i}^{\begin{bmatrix} \alpha \\ \gamma \\ \beta \end{bmatrix} \times k} + \sigma_{i-3,i-2,i-1,i}^{\begin{bmatrix} \alpha \\ \gamma \\ \beta \end{bmatrix} \times k} \end{aligned}$$

●

$$\begin{aligned}
& 16 \cdot \Gamma_{i-2}^{\left[\begin{smallmatrix} s-1 \\ s+m+l-1 \end{smallmatrix} \right] \times k} \\
&= 4 \cdot \left[\sigma_{i-2, i-2, i-2, i-2}^{\left[\begin{smallmatrix} \beta \\ \gamma \end{smallmatrix} \right] \times k} + \sigma_{i-2, i-2, i-2, i-1}^{\left[\begin{smallmatrix} \beta \\ \gamma \end{smallmatrix} \right] \times k} + \sigma_{i-2, i-2, i-1, i-1}^{\left[\begin{smallmatrix} \beta \\ \gamma \end{smallmatrix} \right] \times k} + \sigma_{i-2, i-2, i-1, i}^{\left[\begin{smallmatrix} \beta \\ \gamma \end{smallmatrix} \right] \times k} \right] \\
&+ 4 \cdot \left[\sigma_{i-3, i-2, i-1, i-1}^{\left[\begin{smallmatrix} \beta \\ \gamma \end{smallmatrix} \right] \times k} + \sigma_{i-3, i-2, i-1, i}^{\left[\begin{smallmatrix} \beta \\ \gamma \end{smallmatrix} \right] \times k} + \sigma_{i-3, i-2, i-2, i-2}^{\left[\begin{smallmatrix} \beta \\ \gamma \end{smallmatrix} \right] \times k} + \sigma_{i-3, i-2, i-2, i-1}^{\left[\begin{smallmatrix} \beta \\ \gamma \end{smallmatrix} \right] \times k} \right] \\
&= 4 \cdot \left[\sigma_{i-2, i-2, i-2, i-2}^{\left[\begin{smallmatrix} \beta \\ \alpha \end{smallmatrix} \right] \times k} + \sigma_{i-2, i-2, i-2, i-1}^{\left[\begin{smallmatrix} \beta \\ \alpha \end{smallmatrix} \right] \times k} + \sigma_{i-2, i-2, i-1, i-1}^{\left[\begin{smallmatrix} \beta \\ \alpha \end{smallmatrix} \right] \times k} + \sigma_{i-2, i-2, i-1, i}^{\left[\begin{smallmatrix} \beta \\ \alpha \end{smallmatrix} \right] \times k} \right] \\
&+ 4 \cdot \left[\sigma_{i-3, i-2, i-1, i-1}^{\left[\begin{smallmatrix} \beta \\ \alpha \end{smallmatrix} \right] \times k} + \sigma_{i-3, i-2, i-1, i}^{\left[\begin{smallmatrix} \beta \\ \alpha \end{smallmatrix} \right] \times k} + \sigma_{i-3, i-2, i-2, i-2}^{\left[\begin{smallmatrix} \beta \\ \alpha \end{smallmatrix} \right] \times k} + \sigma_{i-3, i-2, i-2, i-1}^{\left[\begin{smallmatrix} \beta \\ \alpha \end{smallmatrix} \right] \times k} \right]
\end{aligned}$$

●

[illegible]

$$\begin{aligned}
& -8 \cdot \left[\sigma_{i-3,i-2,i-2,i-2} \begin{bmatrix} \gamma \\ \beta \end{bmatrix} \times k + \sigma_{i-3,i-2,i-2,i-1} \begin{bmatrix} \gamma \\ \beta \end{bmatrix} \times k + \sigma_{i-3,i-2,i-1,i-1} \begin{bmatrix} \gamma \\ \beta \end{bmatrix} \times k + \sigma_{i-3,i-2,i-1,i} \begin{bmatrix} \gamma \\ \beta \end{bmatrix} \times k \right] \\
& = -8 \cdot \left[\sigma_{i-3,i-3,i-3,i-3} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-3,i-3,i-3,i-2} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-3,i-3,i-2,i-2} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-3,i-3,i-2,i-1} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k \right] \\
& -8 \cdot \left[\sigma_{i-3,i-2,i-2,i-2} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-3,i-2,i-2,i-1} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-3,i-2,i-1,i-1} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-3,i-2,i-1,i} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k \right]
\end{aligned}$$

For every term in the recurrent formula (2.83) we choose in view of (2.66),(2.74) and (2.82) an appropriate permutation of $\{\alpha, \beta, \gamma\}$, hence we get

$$\begin{aligned}
& \Gamma_i \begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k - \left[2 \cdot \Gamma_{i-1} \begin{bmatrix} s-1 \\ s+m+l \end{bmatrix} \times k + 4 \cdot \Gamma_{i-1} \begin{bmatrix} s+m-1 \\ s+m+l \end{bmatrix} \times k + 8 \cdot \Gamma_{i-1} \begin{bmatrix} s+m \\ s+m+l-1 \end{bmatrix} \times k \right] \\
& + \left[8 \cdot \Gamma_{i-2} \begin{bmatrix} s-1 \\ s+m+l \end{bmatrix} \times k + 16 \cdot \Gamma_{i-2} \begin{bmatrix} s+m-1 \\ s+m+l-1 \end{bmatrix} \times k + 32 \cdot \Gamma_{i-2} \begin{bmatrix} s+m \\ s+m+l-1 \end{bmatrix} \times k - 64 \cdot \Gamma_{i-3} \begin{bmatrix} s-1 \\ s+m+l-1 \end{bmatrix} \times k - \Delta_i \begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k \right] = 0 \\
& \iff \\
& \left[\sigma_{i,i,i,i} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-1,i,i,i} \begin{bmatrix} \gamma \\ \beta \end{bmatrix} \times k + \sigma_{i-2,i-1,i,i} \begin{bmatrix} \gamma \\ \beta \end{bmatrix} \times k + \sigma_{i-1,i-1,i,i} \begin{bmatrix} \gamma \\ \beta \end{bmatrix} \times k + \sigma_{i-1,i-1,i-1,i} \begin{bmatrix} \gamma \\ \beta \end{bmatrix} \times k + \sigma_{i-2,i-1,i-1,i} \begin{bmatrix} \gamma \\ \beta \end{bmatrix} \times k + \sigma_{i-2,i-2,i-1,i} \begin{bmatrix} \gamma \\ \beta \end{bmatrix} \times k + \sigma_{i-3,i-2,i-1,i} \begin{bmatrix} \gamma \\ \beta \end{bmatrix} \times k \right] \\
& - \left[\sigma_{i-1,i-1,i-1,i-1} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-2,i-1,i-1,i-1} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-2,i-2,i-1,i-1} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-3,i-2,i-1,i-1} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k \right] \\
& - \left[\sigma_{i-1,i-1,i-1,i} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-2,i-1,i-1,i} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-2,i-2,i-1,i} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-3,i-2,i-1,i} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k \right] \\
& - 2 \cdot \left[\sigma_{i-1,i-1,i-1,i-1} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-2,i-1,i-1,i-1} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-2,i-2,i-1,i-1} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-3,i-2,i-1,i-1} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k \right] \\
& - 2 \cdot \left[\sigma_{i-1,i-1,i-1,i} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-2,i-1,i-1,i} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-2,i-2,i-1,i} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-3,i-2,i-1,i} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k \right] \\
& - 4 \cdot \left[\sigma_{i-1,i-1,i-1,i-1} \begin{bmatrix} \beta \\ \alpha \end{bmatrix} \times k + \sigma_{i-2,i-1,i-1,i-1} \begin{bmatrix} \beta \\ \alpha \end{bmatrix} \times k + \sigma_{i-2,i-2,i-1,i-1} \begin{bmatrix} \beta \\ \alpha \end{bmatrix} \times k + \sigma_{i-3,i-2,i-1,i-1} \begin{bmatrix} \beta \\ \alpha \end{bmatrix} \times k \right] \\
& - 4 \cdot \left[\sigma_{i-1,i-1,i-1,i} \begin{bmatrix} \beta \\ \alpha \end{bmatrix} \times k + \sigma_{i-2,i-1,i-1,i} \begin{bmatrix} \beta \\ \alpha \end{bmatrix} \times k + \sigma_{i-2,i-2,i-1,i} \begin{bmatrix} \beta \\ \alpha \end{bmatrix} \times k + \sigma_{i-3,i-2,i-1,i} \begin{bmatrix} \beta \\ \alpha \end{bmatrix} \times k \right] \\
& + 2 \cdot \left[\sigma_{i-2,i-2,i-2,i-2} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-2,i-2,i-2,i-1} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-2,i-2,i-1,i-1} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-2,i-2,i-1,i} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k \right] \\
& + 2 \cdot \left[\sigma_{i-3,i-2,i-1,i-1} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-3,i-2,i-1,i} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-3,i-2,i-2,i-2} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k + \sigma_{i-3,i-2,i-2,i-1} \begin{bmatrix} \gamma \\ \alpha \end{bmatrix} \times k \right]
\end{aligned}$$

$$-8 \cdot \left[\sigma_{i-3,i-3,i-3,i-2}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} + \sigma_{i-3,i-3,i-2,i-2}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} + \sigma_{i-3,i-3,i-2,i-1}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} \right] = 0$$

Combining (2.66), (2.74) and (2.82) we get

$$\begin{aligned}
(2.83) \iff & \left[\sigma_{i-1,i,i,i}^{\begin{bmatrix} \gamma \\ \beta \\ \alpha \end{bmatrix} \times k} + 2 \cdot \left[\sigma_{i-1,i-1,i-1,i}^{\begin{bmatrix} \gamma \\ \alpha \\ \beta \end{bmatrix} \times k} + \sigma_{i-2,i-1,i-1,i}^{\begin{bmatrix} \gamma \\ \alpha \\ \beta \end{bmatrix} \times k} \right] \right. \\
& - \left[\sigma_{i-2,i-1,i-1,i-1}^{\begin{bmatrix} \gamma \\ \beta \\ \alpha \end{bmatrix} \times k} + 2 \cdot \left[\sigma_{i-2,i-2,i-2,i-1}^{\begin{bmatrix} \gamma \\ \alpha \\ \beta \end{bmatrix} \times k} + \sigma_{i-3,i-2,i-2,i-1}^{\begin{bmatrix} \gamma \\ \alpha \\ \beta \end{bmatrix} \times k} \right] \right. \\
& - 2 \cdot \left[\sigma_{i-2,i-1,i-1,i-1}^{\begin{bmatrix} \gamma \\ \alpha \\ \beta \end{bmatrix} \times k} + \sigma_{i-1,i-1,i-1,i}^{\begin{bmatrix} \gamma \\ \alpha \\ \beta \end{bmatrix} \times k} + \sigma_{i-2,i-1,i-1,i}^{\begin{bmatrix} \gamma \\ \alpha \\ \beta \end{bmatrix} \times k} \right] \\
& - 4 \cdot \left[\sigma_{i-1,i-1,i-1,i}^{\begin{bmatrix} \beta \\ \alpha \\ \gamma \end{bmatrix} \times k} + 2 \cdot \left[\sigma_{i-2,i-2,i-1,i-1}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} + \sigma_{i-2,i-2,i-1,i}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} \right] \right. \\
& + 2 \cdot \left[\sigma_{i-2,i-2,i-2,i-1}^{\begin{bmatrix} \gamma \\ \beta \\ \alpha \end{bmatrix} \times k} + \sigma_{i-3,i-2,i-2,i-2}^{\begin{bmatrix} \gamma \\ \beta \\ \alpha \end{bmatrix} \times k} + \sigma_{i-3,i-2,i-2,i-1}^{\begin{bmatrix} \gamma \\ \beta \\ \alpha \end{bmatrix} \times k} \right] \\
& + 4 \cdot \left[\sigma_{i-2,i-2,i-2,i-1}^{\begin{bmatrix} \beta \\ \gamma \\ \alpha \end{bmatrix} \times k} + 2 \cdot \left[\sigma_{i-3,i-3,i-2,i-2}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} + \sigma_{i-3,i-3,i-2,i-1}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} \right] \right. \\
& + 8 \cdot \left[\sigma_{i-2,i-2,i-2,i-1}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} + \sigma_{i-2,i-2,i-1,i-1}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} + \sigma_{i-2,i-2,i-1,i}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} \right] \\
& \left. \left. - 8 \cdot \left[\sigma_{i-3,i-3,i-3,i-2}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} + \sigma_{i-3,i-3,i-2,i-2}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} + \sigma_{i-3,i-3,i-2,i-1}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} \right] \right] = 0 \\
& \iff \\
& \left(\sigma_{i-1,i,i,i}^{\begin{bmatrix} \gamma \\ \beta \\ \alpha \end{bmatrix} \times k} - 4 \cdot \sigma_{i-1,i-1,i-1,i}^{\begin{bmatrix} \beta \\ \alpha \\ \gamma \end{bmatrix} \times k} \right) - \left(\sigma_{i-2,i-1,i-1,i-1}^{\begin{bmatrix} \gamma \\ \beta \\ \alpha \end{bmatrix} \times k} - 4 \cdot \sigma_{i-2,i-2,i-2,i-1}^{\begin{bmatrix} \beta \\ \alpha \\ \gamma \end{bmatrix} \times k} \right) \\
& - 2 \cdot \left(\sigma_{i-2,i-1,i-1,i-1}^{\begin{bmatrix} \gamma \\ \alpha \\ \beta \end{bmatrix} \times k} - 4 \cdot \sigma_{i-2,i-2,i-2,i-1}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} \right) + 2 \cdot \left(\sigma_{i-3,i-2,i-2,i-2}^{\begin{bmatrix} \gamma \\ \alpha \\ \beta \end{bmatrix} \times k} - 4 \cdot \sigma_{i-3,i-3,i-3,i-2}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k} \right) = 0
\end{aligned}$$

□

3. Rank properties of a partition of triple persymmetric matrices

3.1. Notation.

Definition 3.1. We introduce the following definitions in the three - dimensional $\mathbb{K}$ - vectorspace.

- Let k, s, m and l denote rational integers such that $k \geq 1$, $s \geq 2$ and $m \geq 0$, $l \geq 0$
- We denote by $\mathbb{P}/\mathbb{P}_i \times \mathbb{P}/\mathbb{P}_j \times \mathbb{P}/\mathbb{P}_r$ a complete set of coset representatives of $\mathbb{P}_i \times \mathbb{P}_j \times \mathbb{P}_r$ in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$, for instance $\mathbb{P}/\mathbb{P}_{s+k-1} \times \mathbb{P}/\mathbb{P}_{s+m+k-1} \times \mathbb{P}/\mathbb{P}_{s+m+l+k-1}$ denotes a complete set of coset representatives of $\mathbb{P}_{s+k-1} \times \mathbb{P}_{s+m+k-1} \times \mathbb{P}_{s+m+l+k-1}$ in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$.

- $Set(t, \eta, \xi) = (\sum_{i \geq 1} \alpha_i T^{-i}, \sum_{i \geq 1} \beta_i T^{-i}, \sum_{i \geq 1} \gamma_i T^{-i}) \in \mathbb{P}^3$

- We denote by $D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_s - \\ \beta_{s+m} - \\ \gamma_{s+m+l} - \end{bmatrix} \times k$ any $(3s+2m+l) \times k$ matrix, such that after a rearrangement of the rows, if necessary, we can obtain a matrix where the first $(3s+2m+l-3)$ rows form the following triple persymmetric matrix

$$\begin{bmatrix} \frac{D_{s-1 \times k}(t)}{D_{(s+m-1) \times k}(\eta)} \\ \frac{D_{(s+m+l-1) \times k}(\xi)}{D_{(s+m+l-1) \times k}(\xi)} \end{bmatrix} \text{ and the last three rows are equal to}$$

the following $3 \times k$ matrix $\begin{pmatrix} \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-1} \\ \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-1} \\ \gamma_{s+m+l} & \gamma_{s+m+l+1} & \dots & \gamma_{s+m+l+k-1} \end{pmatrix}$, that is we get a matrix of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\ \hline \beta_1 & \beta_2 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \dots & \beta_{k+m-1} & \beta_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \hline \gamma_1 & \gamma_2 & \dots & \gamma_{k-1} & \gamma_k \\ \gamma_2 & \gamma_3 & \dots & \gamma_k & \gamma_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{m+l+1} & \gamma_{m+l+2} & \dots & \gamma_{k+m+l-1} & \gamma_{k+m+l} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{s+m+l-1} & \gamma_{s+m+l} & \dots & \gamma_{s+m+l+k-3} & \gamma_{s+m+l+k-2} \\ \hline \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-2} & \alpha_{s+k-1} \\ \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \\ \gamma_{s+m+l} & \gamma_{s+m+l+1} & \dots & \gamma_{s+m+l+k-2} & \gamma_{s+m+l+k-1} \end{pmatrix}.$$

- Let j be a rational integer such that $1 \leq j \leq k-1$.

$$Set(t, \eta, \xi) = (\sum_{i \geq 1} \alpha_i T^{-i}, \sum_{i \geq 1} \beta_i T^{-i}, \sum_{i \geq 1} \gamma_i T^{-i}) \in \mathbb{P}^3$$

We denote by $D_j \begin{bmatrix} s \\ s+m \\ s+m+l-1 \end{bmatrix} \times (k-j+1)$ any $(3s+2m+l) \times (k-j+1)$ matrix, such that after a rearrangement of the rows, if necessary, we can obtain the following

$$\text{triple persymmetric matrix } \begin{bmatrix} \frac{D_{s \times (k-j+1)}^j(t)}{D_{(s+m) \times (k-j+1)}^j(\eta)} \\ \frac{D_{(s+m+l) \times (k-j+1)}^j(\xi)}{D_{(s+m+l) \times (k-j+1)}^j(\xi)} \end{bmatrix}$$

$$\left(\begin{array}{ccccc|c}
\alpha_j & \alpha_{j+1} & \alpha_{j+2} & \dots & \alpha_{k-1} & \alpha_k \\
\alpha_{j+1} & \alpha_{j+2} & \alpha_{j+3} & \dots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{j+s-1} & \alpha_{j+s} & \alpha_{j+s+1} & \dots & \alpha_{k+s-2} & \alpha_{k+s-1} \\
\hline
\beta_j & \beta_{j+1} & \beta_{j+2} & \dots & \beta_{k-1} & \beta_k \\
\beta_{j+1} & \beta_{j+2} & \beta_{j+3} & \dots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+j} & \beta_{m+j+1} & \beta_{m+j+2} & \dots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m+j-1} & \beta_{s+m+j} & \beta_{s+m+j+1} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \\
\hline
\gamma_j & \gamma_{j+1} & \gamma_{j+2} & \dots & \gamma_{k-1} & \gamma_k \\
\gamma_{j+1} & \gamma_{j+2} & \gamma_{j+3} & \dots & \gamma_k & \gamma_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\gamma_{m+l+j} & \gamma_{m+l+j+1} & \gamma_{m+l+j+2} & \dots & \gamma_{k+m+l-1} & \gamma_{k+m+l} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\gamma_{s+m+l+j-1} & \gamma_{s+m+l+j} & \gamma_{s+m+l+j+1} & \dots & \gamma_{s+m+l+k-2} & \gamma_{s+m+l+k-1}
\end{array} \right).$$

- To simplify the notations concerning the exponential sums used in the proofs, we introduce the following definitions.

- Let $\psi_1(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$ defined by

$$(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P} \mapsto \sum_{\deg Y = k-1} \sum_{\deg Z = s-1} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \in \mathbb{Z}$$

- Let $\psi_2(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$ defined by

$$(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P} \mapsto \sum_{\deg Y \leq k-2} \sum_{\deg Z \leq s-1} E(tYZ) \sum_{\deg U = s+m-1} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \in \mathbb{Z}$$

- Let $\psi_3(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$ defined by

$$(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P} \mapsto \sum_{\deg Y \leq k-2} \sum_{\deg Z \leq s-1} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V = s+m+l-1} E(\xi YV) \in \mathbb{Z}$$

- Let $\chi_1(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$ defined by

$$(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P} \mapsto \sum_{\deg Y = k-1} \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \in \mathbb{Z}$$

- Let $\chi_2(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$ defined by

$$(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P} \mapsto \sum_{\deg Y \leq k-2} \sum_{\deg Z = s-1} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \in \mathbb{Z}$$

- We recall the definition of $h_1(t, \eta, \xi)$ (see Definition 2.1)

Let $h_1(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$ defined by

$$(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P} \mapsto \sum_{\deg Y \leq k-1} \sum_{\deg Z = s-1} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \in \mathbb{Z}$$

- Let $\theta_1(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$ defined by

$$(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P} \mapsto \sum_{\deg Y = k-1} \sum_{\deg Z \leq s-1} E(tYZ) \sum_{\deg U = s+m-1} E(\eta YU) \sum_{\deg V = s+m+l-2} E(\xi YV) \in \mathbb{Z}$$

- Let $\theta_2(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P} \times \mathbb{P} \times \mathbb{P}$ defined by

$$(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P} \mapsto \sum_{\deg Y = k-1} \sum_{\deg Z \leq s-1} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V = s+m+l-1} E(\xi YV) \in \mathbb{Z}$$

- Let $(j_1, j_2, j_3, j_4, j_5, j_6, j_7, j_8) \in \mathbb{N}^8$, we define

$$\# \left(\begin{array}{c|c} j_1 & j_2 \\ \hline j_3 & j_4 \\ \hline j_5 & j_6 \\ \hline j_7 & j_8 \end{array} \right)_{\frac{\frac{\alpha}{2}}{\gamma}}^{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}}$$

to be the cardinality of the following set

$$\left\{ (t, \eta, \xi) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1} \mid r(D \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right]^{\times(k-1)} (t, \eta, \xi)) = j_1, \right. \\ r(D \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right]^{\times k} (t, \eta, \xi)) = j_2, r(D \left[\begin{smallmatrix} s \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right]^{\times(k-1)} (t, \eta, \xi)) = j_3, \\ r(D \left[\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix} \right]^{\times k} (t, \eta, \xi)) = j_4, \quad r(D \left[\begin{smallmatrix} s \\ s+m \\ s+m+l-1 \end{smallmatrix} \right]^{\times(k-1)} (t, \eta, \xi)) = j_5, \\ \left. r(D \left[\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix} \right]^{\times k} (t, \eta, \xi)) = j_6, \quad r(D \left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right]^{\times(k-1)} (t, \eta, \xi)) = j_7, \quad r(D \left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right]^{\times k} (t, \eta, \xi)) = j_8. \right\}$$

- We define

$k-1$	k	
.	.	$D \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right]^{\times \cdot}$
j	j	$\alpha_s -$
j	$j+1$	$\beta_{s+m} -$
.	.	$\gamma_{s+m+l} -$

to be the following subset of $\mathbb{P}^3$

$$\left\{ (t, \eta, \xi) = (\sum_{i \geq 1} \alpha_i T^{-i}, \sum_{i \geq 1} \beta_i T^{-i}, \sum_{i \geq 1} \gamma_i T^{-i}) \in \mathbb{P}^3 \mid r(D \left[\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix} \right]^{\times(k-1)} (t, \eta, \xi)) = j, \right. \\ \left. r(D \left[\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix} \right]^{\times k} (t, \eta, \xi)) = j, \quad r(D \left[\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix} \right]^{\times(k-1)} (t, \eta, \xi)) = j, \quad r(D \left[\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix} \right]^{\times k} (t, \eta, \xi)) = j+1 \right\}$$

- Similar expressions are defined in a similar way.

3.2. Introduction. We study rank properties of a partition of triple persymmetric matrices by integrating some appropriate exponential sums on the unit interval of $\mathbf{K}^3$. We adapt the method used in Section 6 and Section 7 of [2]

3.3. Computation of exponential sums in $\mathbb{K}^3$.

Lemma 3.2. *Let $(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P}$ and*

$$\begin{aligned}
 \psi_1(t, \eta, \xi) &= \sum_{\deg Y = k-1} \sum_{\deg Z = s-1} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \\
 \psi_2(t, \eta, \xi) &= \sum_{\deg Y \leq k-2} \sum_{\deg Z \leq s-1} E(tYZ) \sum_{\deg U = s+m-1} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \\
 \psi_3(t, \eta, \xi) &= \sum_{\deg Y \leq k-2} \sum_{\deg Z \leq s-1} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V = s+m+l-1} E(\xi YV) \\
 \chi_1(t, \eta, \xi) &= \sum_{\deg Y = k-1} \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \\
 \chi_2(t, \eta, \xi) &= \sum_{\deg Y \leq k-2} \sum_{\deg Z = s-1} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \\
 h_1(t, \eta, \xi) &= \sum_{\deg Y \leq k-1} \sum_{\deg Z = s-1} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV)
 \end{aligned}$$

Then

$$(3.1) \quad \psi_1(t, \eta, \xi) = 2^{3s+2m+l-3} \cdot \sum_{\substack{\deg Y = k-1 \\ \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k \\ Y \in \ker D}} E(tYT^{s-1})_{(t, \eta, \xi)}$$

(3.2)

$$\begin{aligned}
 \psi_2(t, \eta, \xi) &= 2^{3s+2m+l-2} \cdot \sum_{\substack{\deg Y \leq k-2 \\ \begin{bmatrix} s \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1) \\ Y \in \ker D}} E(\eta YT^{s+m-1})_{(t, \eta, \xi)} \\
 &= \begin{cases} 2^{k+3s+2m+l-3-r(D \begin{bmatrix} s+m-1 \\ s+m+l-1 \end{bmatrix} \times k)}_{(t, \eta, \xi)} & \text{if } r(D \begin{bmatrix} s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1))_{(t, \eta, \xi)} = r(D \begin{bmatrix} s \\ s+m+l-1 \end{bmatrix} \times (k-1))_{(t, \eta, \xi)} \\ 0 & \text{otherwise} \end{cases}
 \end{aligned}$$

(3.3)

$$\begin{aligned}
 \psi_3(t, \eta, \xi) &= 2^{3s+2m+l-1} \cdot \sum_{\substack{\deg Y \leq k-2 \\ \begin{bmatrix} s \\ s+m \\ s+m+l-1 \end{bmatrix} \times (k-1) \\ Y \in \ker D}} E(\xi YT^{s+m+l-1})_{(t, \eta, \xi)} \\
 &= \begin{cases} 2^{k+3s+2m+l-2-r(D \begin{bmatrix} s+m \\ s+m+l-1 \end{bmatrix} \times (k-1))_{(t, \eta, \xi)}} & \text{if } r(D \begin{bmatrix} s+m \\ s+m+l-1 \end{bmatrix} \times (k-1))_{(t, \eta, \xi)} = r(D \begin{bmatrix} s \\ s+m+l \end{bmatrix} \times (k-1))_{(t, \eta, \xi)} \\ 0 & \text{otherwise} \end{cases}
 \end{aligned}$$

(3.4)

$$\chi_1(t, \eta, \xi) = \begin{cases} 2^{k+3s+2m+l-4-r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1))_{(t, \eta, \xi)}} & \text{if } r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1))_{(t, \eta, \xi)} = r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k)_{(t, \eta, \xi)} \\ 0 & \text{otherwise} \end{cases}$$

(3.5)

$$\chi_2(t, \eta, \xi) = \begin{cases} 2^{k+3s+2m+l-4-r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1))} & (t, \eta, \xi) \\ 0 & \text{otherwise} \end{cases} \quad \text{if } r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1)) = r(D \begin{bmatrix} s \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1)) \quad (t, \eta, \xi)$$

We recall that see (2.2)

(3.6)

$$h_1(t, \eta, \xi) = 2^{3s+2m+l-3} \cdot \sum_{\substack{\deg Y \leq k-1 \\ Y \in \ker D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k}} E(tYT^{s-1}) \quad (t, \eta, \xi)$$

$$= \begin{cases} 2^{k+3s+2m+l-3-r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k)} & (t, \eta, \xi) \\ 0 & \text{otherwise} \end{cases} \quad \text{if } r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k) = r(D \begin{bmatrix} s \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k) \quad (t, \eta, \xi)$$

Proof. The proof of 3.2 is somewhat similar to the proofs of the results obtained in [2, see section 4] $\square$

Lemma 3.3. *Let $(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P}$ then we have*

$$(3.7) \quad \psi_1^2(t, \eta, \xi) = \chi_1(t, \eta, \xi) \cdot \chi_2(t, \eta, \xi)$$

where

$$\begin{aligned} \psi_1(t, \eta, \xi) &= \sum_{\deg Y = k-1} \sum_{\deg Z = s-1} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \\ \chi_1(t, \eta, \xi) &= \sum_{\deg Y = k-1} \sum_{\deg Z \leq s-2} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \\ \chi_2(t, \eta, \xi) &= \sum_{\deg Y \leq k-2} \sum_{\deg Z = s-1} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \end{aligned}$$

Proof. We have

$$\psi_1^2(t, \eta, \xi) = [2^{3s+2m+l-3} \cdot \sum_{\substack{\deg Y_1 = k-1 \\ \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k \\ Y_1 \in \ker D}} E(tY_1 T^{s-1})] [2^{3s+2m+l-3} \cdot \sum_{\substack{\deg Y_2 = k-1 \\ \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k \\ Y_2 \in \ker D}} E(tY_2 T^{s-1})]$$

We set

$$\begin{cases} Y_1 + Y_2 = Y_3, & \deg Y_1 = k-1, & Y_1 \in \ker D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k \\ Y_2 = Y_4, & \deg Y_2 = k-1, & Y_2 \in \ker D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k \end{cases} (t, \eta, \xi)$$

$$\iff \begin{cases} \deg Y_3 \leq k-2, & Y_3 \in \ker D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1) \\ \deg Y_4 = k-1, & Y_4 \in \ker D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k \end{cases} (t, \eta, \xi)$$

Then we obtain

$$\begin{aligned} \psi_1^2(t, \eta, \xi) &= [2^{3s+2m+l-3} \cdot \sum_{\substack{\deg Y_4 = k-1 \\ \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k \\ Y_4 \in \ker D}} 1] [2^{3s+2m+l-3} \cdot \sum_{\substack{\deg Y_3 \leq k-2 \\ \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k \\ Y_3 \in \ker D}} E(tY_3 T^{s-1})] \\ &= \chi_1(t, \eta, \xi) \cdot \chi_2(t, \eta, \xi) \end{aligned}$$

□

Lemma 3.4. *We have the following equivalences :*

(3.8)

$$\psi_1(t, \eta, \xi) \neq 0 \iff \psi_1^2(t, \eta, \xi) \neq 0 \iff \chi_1(t, \eta, \xi) \cdot \chi_2(t, \eta, \xi) \neq 0$$

$$\iff \chi_1(t, \eta, \xi) \neq 0 \quad \text{and} \quad \chi_2(t, \eta, \xi) \neq 0$$

$$\iff r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1) (t, \eta, \xi)) = r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k (t, \eta, \xi)) = r(D \begin{bmatrix} s \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1) (t, \eta, \xi))$$

Proof. (3.8) follows from (3.4) and (3.5).

□

Lemma 3.5. *We have*

(3.9)

$$\psi_1(t, \eta, \xi) = \begin{cases} 2^{k+3s+2m+l-4-j} & \text{if } (t, \eta, \xi) \in \begin{array}{|c|c|c|} \hline k-1 & k & \\ \hline j & j & D \begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \end{array} \times \cdot \\ \hline j & j & \alpha_s - \\ \hline \cdot & \cdot & \beta_{s+m} - \\ \hline \cdot & \cdot & \gamma_{s+m+l} - \\ \hline \end{array} \\ \\ -2^{k+3s+2m+l-4-j} & \text{if } (t, \eta, \xi) \in \begin{array}{|c|c|c|} \hline k-1 & k & \\ \hline j & j & D \begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \end{array} \times \cdot \\ \hline j & j+1 & \alpha_s - \\ \hline \cdot & \cdot & \beta_{s+m} - \\ \hline \cdot & \cdot & \gamma_{s+m+l} - \\ \hline \end{array} \\ \\ 0 & \text{otherwise} \end{cases}$$

$$(3.10) \quad \psi_2(t, \eta, \xi) = \begin{cases} 2^{k+3s+2m+l-3-j} & \text{if } (t, \eta, \xi) \in \begin{array}{|c|c|c|} \hline k-1 & k & \\ \hline \cdot & \cdot & D \begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \end{array} \times k \\ \hline j & \cdot & \alpha_s - \\ \hline j & \cdot & \beta_{s+m} - \\ \hline \cdot & \cdot & \gamma_{s+m+l} - \\ \hline \end{array} \\ \\ 0 & \text{otherwise} \end{cases}$$

$$(3.11) \quad \psi_3(t, \eta, \xi) = \begin{cases} 2^{k+3s+2m+l-2-j} & \text{if } (t, \eta, \xi) \in \begin{array}{|c|c|c|} \hline k-1 & k & \\ \hline \cdot & \cdot & D \begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \end{array} \times \cdot \\ \hline \cdot & \cdot & \alpha_s - \\ \hline j & \cdot & \beta_{s+m} - \\ \hline j & \cdot & \gamma_{s+m+l} - \\ \hline \end{array} \\ \\ 0 & \text{otherwise} \end{cases}$$

Proof. We deduce respectively from (3.6) and (3.6) with $k \rightarrow k-1$

(3.12)

$$\begin{aligned} \psi_1(t, \eta, \xi) &= \sum_{\deg Y = k-1} \sum_{\deg Z = s-1} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \\ &= \sum_{\deg Y \leq k-1} \sum_{\deg Z = s-1} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \\ &\quad - \sum_{\deg Y \leq k-2} \sum_{\deg Z = s-1} E(tYZ) \sum_{\deg U \leq s+m-2} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \end{aligned}$$

$$\begin{aligned}
&= \begin{cases} 2^{k+3s+2m+l-3-r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k)} (t, \eta, \xi) & \text{if } r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k) (t, \eta, \xi) = r(D \begin{bmatrix} s \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k) (t, \eta, \xi) \\ 0 & \text{otherwise .} \end{cases} \\
&- \begin{cases} 2^{k+3s+2m+l-4-r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1))} (t, \eta, \xi) & \text{if } r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1)) (t, \eta, \xi) = r(D \begin{bmatrix} s \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1)) (t, \eta, \xi) \\ 0 & \text{otherwise .} \end{cases}
\end{aligned}$$

By Lemma 3.4

$$\psi_1(t, \eta, \xi) \neq 0 \iff r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1)) (t, \eta, \xi) = r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k) (t, \eta, \xi) = r(D \begin{bmatrix} s \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1)) (t, \eta, \xi)$$

We consider now the following two cases in which ψ_1 are different from zero :
first case :

$$\begin{aligned}
&r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1)) (t, \eta, \xi) = r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k) (t, \eta, \xi) \\
&= r(D \begin{bmatrix} s \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1)) (t, \eta, \xi) = r(D \begin{bmatrix} s \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k) (t, \eta, \xi) = j
\end{aligned}$$

second case :

$$\begin{aligned}
&r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1)) (t, \eta, \xi) = r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k) (t, \eta, \xi) \\
&= r(D \begin{bmatrix} s \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1)) (t, \eta, \xi) = j, \quad r(D \begin{bmatrix} s \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k) (t, \eta, \xi) = j + 1
\end{aligned}$$

In the first case we obtain from (3.12)

$$\psi_1(t, \eta, \xi) = 2^{k+3s+2m+l-3-j} - 2^{k+3s+2m+l-4-j} = 2^{k+3s+2m+l-4-j}$$

In the second case we obtain from (3.12)

$$\psi_1(t, \eta, \xi) = 0 - 2^{k+3s+2m+l-4-j} = -2^{k+3s+2m+l-4-j}$$

This completes the proof of (3.9).

(3.10) and (3.11) follow respectively from (3.2) and (3.3). $\square$

Lemma 3.6. *We have*

(3.13)

$$\psi_1(t, \eta, \xi) \cdot \psi_2(t, \eta, \xi) \cdot \psi_3^{q-2}(t, \eta, \xi) = \begin{cases} 2^{(k+3s+2m+l-3-j)q} \cdot 2^{q-3} & \text{if } (t, \eta, \xi) \in \begin{array}{|c|c|c|} \hline k-1 & k & \\ \hline j & j & D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times \cdot \\ \hline j & j & \alpha_s - \\ \hline j & \cdot & \beta_{s+m} - \\ \hline j & \cdot & \gamma_{s+m+l} - \\ \hline \end{array} \\ \\ -2^{(k+3s+2m+l-3-j)q} \cdot 2^{q-3} & \text{if } (t, \eta, \xi) \in \begin{array}{|c|c|c|} \hline k-1 & k & \\ \hline j & j & D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times \cdot \\ \hline j & j+1 & \alpha_s - \\ \hline j & j+1 & \beta_{s+m} - \\ \hline j & j+1 & \gamma_{s+m+l} - \\ \hline \end{array} \\ \\ 0 & \text{otherwise} \end{cases}$$

Proof. Combining (3.9), (3.10) and (3.11) we deduce (3.13). $\square$

Lemma 3.7. *Let $(t, \eta, \xi) \in \mathbb{P} \times \mathbb{P} \times \mathbb{P}$ and*

$$\begin{aligned} \theta_1(t, \eta, \xi) &= \sum_{\deg Y = k-1} \sum_{\deg Z \leq s-1} E(tYZ) \sum_{\deg U = s+m-1} E(\eta YU) \sum_{\deg V \leq s+m+l-2} E(\xi YV) \\ \theta_2(t, \eta, \xi) &= \sum_{\deg Y = k-1} \sum_{\deg Z \leq s-1} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V = s+m+l-1} E(\xi YV) \end{aligned}$$

Then

(3.14)

$$\theta_1(t, \eta, \xi) = \begin{cases} 2^{k+3s+2m+l-3-j} & \text{if } (t, \eta, \xi) \in \begin{array}{|c|c|c|} \hline k-1 & k & \\ \hline \cdot & \cdot & D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times \cdot \\ \hline j & j & \alpha_s - \\ \hline j & j & \beta_{s+m} - \\ \hline \cdot & \cdot & \gamma_{s+m+l} - \\ \hline \end{array} \\ \\ -2^{k+3s+2m+l-3-j} & \text{if } (t, \eta, \xi) \in \begin{array}{|c|c|c|} \hline k-1 & k & \\ \hline \cdot & \cdot & D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times \cdot \\ \hline j & j & \alpha_s - \\ \hline j & j+1 & \beta_{s+m} - \\ \hline \cdot & \cdot & \gamma_{s+m+l} - \\ \hline \end{array} \\ \\ 0 & \text{otherwise} \end{cases}$$

$$(3.15) \quad \theta_2(t, \eta, \xi) = \begin{cases} 2^{k+3s+2m+l-3-j} & \text{if } (t, \eta, \xi) \in \begin{array}{|c|c|c|} \hline k-1 & k & \\ \hline \cdot & \cdot & D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times \cdot \\ \hline \cdot & \cdot & \alpha_s - \\ \hline j & j & \beta_{s+m} - \\ \hline j & j & \gamma_{s+m+l} - \\ \hline \end{array} \\ \\ -2^{k+3s+2m+l-3-j} & \text{if } (t, \eta, \xi) \in \begin{array}{|c|c|c|} \hline k-1 & k & \\ \hline \cdot & \cdot & D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times \cdot \\ \hline \cdot & \cdot & \alpha_s - \\ \hline j & j & \beta_{s+m} - \\ \hline j & j+1 & \gamma_{s+m+l} - \\ \hline \end{array} \\ \\ 0 & \text{otherwise} \end{cases}$$

Proof. Similarly to the proof of (3.9). $\square$

Lemma 3.8. *We have*

$$(3.16) \quad h_1(t, \eta, \xi) \cdot \theta_1(t, \eta, \xi) \cdot \psi_3^{q-2}(t, \eta, \xi) = \begin{cases} 2^{(k+3s+2m+l-3-j)q} \cdot 2^{q-2} & \text{if } (t, \eta, \xi) \in \begin{array}{|c|c|c|} \hline k-1 & k & \\ \hline j & j & D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times \cdot \\ \hline j & j & \alpha_s - \\ \hline j & j & \beta_{s+m} - \\ \hline j & \cdot & \gamma_{s+m+l} - \\ \hline \end{array} \\ \\ -2^{(k+3s+2m+l-3-j)q} \cdot 2^{q-2} & \text{if } (t, \eta, \xi) \in \begin{array}{|c|c|c|} \hline k-1 & k & \\ \hline j & j & D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times \cdot \\ \hline j & j & \alpha_s - \\ \hline j & j+1 & \beta_{s+m} - \\ \hline j & \cdot & \gamma_{s+m+l} - \\ \hline \end{array} \\ \\ 0 & \text{otherwise} \end{cases}$$

Proof. Combining (3.6), (3.14) and (3.11) we deduce (3.16). $\square$

Lemma 3.9. *We have*

(3.17)

$$h_1^{q-2}(t, \eta, \xi) \cdot \psi_2(t, \eta, \xi) \cdot \theta_2(t, \eta, \xi) = \begin{cases} 2^{(k+3s+2m+l-3-j)q+1} & \text{if } (t, \eta, \xi) \in \begin{array}{|c|c|c|} \hline k-1 & k & \\ \hline j & j & D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times \cdot \\ \hline j & j & \alpha_s - \\ \hline j & j & \beta_{s+m} - \\ \hline j & j & \gamma_{s+m+l} - \\ \hline \end{array} \\ \\ -2^{(k+3s+2m+l-3-j)q+1} & \text{if } (t, \eta, \xi) \in \begin{array}{|c|c|c|} \hline k-1 & k & \\ \hline j & j & D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times \cdot \\ \hline j & j & \alpha_s - \\ \hline j & j & \beta_{s+m} - \\ \hline j & j+1 & \gamma_{s+m+l} - \\ \hline \end{array} \\ \\ 0 & \text{otherwise} \end{cases}$$

Proof. Combining (3.6), (3.10) and (3.15) we deduce (3.17). $\square$

3.4. Some rank formulas for partitions of triple persymmetric matrices related

to $\sigma_{i,i,i,i}^{\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times k}$.

Lemma 3.10. *We have for $q \geq 3$*

$$\int_{\mathbb{P} \times \mathbb{P} \times \mathbb{P}} \psi_1(t, \eta, \xi) \cdot \psi_2(t, \eta, \xi) \cdot \psi_3^{q-2}(t, \eta, \xi) dt d\eta d\xi = 0.$$

Proof. The integral above is equal to the number of solutions

$(Y_1, Z_1, U_1, V_1, Y_2, Z_2, U_2, V_2, \dots, Y_q, Z_q, U_q, V_q)$ of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0, \\ Y_1 U_1 + Y_2 U_2 + \dots + Y_q U_q = 0, \\ Y_1 V_1 + Y_2 V_2 + \dots + Y_q V_q = 0. \end{cases}$$

satisfying the degree conditions

$$\begin{cases} \deg Y_1 = k-1, & \deg Z_1 = s-1 & \deg U_1 \leq s+m-2 & \deg V_1 \leq s+m+l-2 \\ \deg Y_2 \leq k-2, & \deg Z_2 \leq s-1, & \deg U_2 = s+m-1, & \deg V_2 \leq s+m+l-2 \\ \deg Y_i \leq k-2, & \deg Z_i \leq s-1, & \deg U_i \leq s+m-1, & \deg V_i = s+m+l-1 \quad \text{for } 3 \leq i \leq q \end{cases}$$

By degree considerations we have that $\deg[Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q]$ is equal to $s+k-2$.

Lemma 3.10 follows. $\square$

Lemma 3.11. *We have for $q \geq 3$*

$$\int_{\mathbb{P} \times \mathbb{P} \times \mathbb{P}} h_1(t, \eta, \xi) \cdot \theta_1(t, \eta, \xi) \cdot \psi_3^{q-2}(t, \eta, \xi) dt d\eta d\xi = 0.$$

Proof. The integral above is equal to the number of solutions

$(Y_1, Z_1, U_1, V_1, Y_2, Z_2, U_2, V_2, \dots, Y_q, Z_q, U_q, V_q)$ of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0, \\ Y_1 U_1 + Y_2 U_2 + \dots + Y_q U_q = 0, \\ Y_1 V_1 + Y_2 V_2 + \dots + Y_q V_q = 0. \end{cases}$$

satisfying the degree conditions

$$\begin{cases} \deg Y_1 \leq k-1, & \deg Z_1 = s-1 & \deg U_1 \leq s+m-2 & \deg V_1 \leq s+m+l-2 \\ \deg Y_2 = k-1, & \deg Z_2 \leq s-1, & \deg U_2 = s+m-1, & \deg V_2 \leq s+m+l-2 \\ \deg Y_i \leq k-2, & \deg Z_i \leq s-1, & \deg U_i \leq s+m-1, & \deg V_i = s+m+l-1 \end{cases} \text{ for } 3 \leq i \leq q$$

By degree considerations we have that $\deg[Y_1U_1 + Y_2U_2 + \dots + Y_qU_q]$ is equal to $s+k+m-2$. Lemma 3.11 follows. $\square$

Lemma 3.12. *We have for $q \geq 3$*

$$\int_{\mathbb{P} \times \mathbb{P} \times \mathbb{P}} h_1^{q-2}(t, \eta, \xi) \cdot \psi_2(t, \eta, \xi) \cdot \theta_2(t, \eta, \xi) dt d\eta d\xi = 0.$$

Proof. The integral above is equal to the number of solutions $(Y_1, Z_1, U_1, V_1, Y_2, Z_2, U_2, V_2, \dots, Y_q, Z_q, U_q, V_q)$ of the polynomial equations

$$\begin{cases} Y_1Z_1 + Y_2Z_2 + \dots + Y_qZ_q = 0, \\ Y_1U_1 + Y_2U_2 + \dots + Y_qU_q = 0, \\ Y_1V_1 + Y_2V_2 + \dots + Y_qV_q = 0. \end{cases}$$

satisfying the degree conditions

$$\begin{cases} \deg Y_1 \leq k-2, & \deg Z_1 \leq s-1 & \deg U_1 = s+m-1 & \deg V_1 \leq s+m+l-2 \\ \deg Y_2 = k-1, & \deg Z_2 \leq s-1, & \deg U_2 \leq s+m-1, & \deg V_2 = s+m+l-1 \\ \deg Y_i \leq k-1, & \deg Z_i = s-1, & \deg U_i \leq s+m-2, & \deg V_i \leq s+m+l-2 \end{cases} \text{ for } 3 \leq i \leq q$$

By degree considerations we have that $\deg[Y_1V_1 + Y_2V_2 + \dots + Y_qV_q]$ is equal to $k+s+m+l-2$. Lemma 3.12 follows. $\square$

Lemma 3.13. *We have the following rank identity for $0 \leq j \leq \inf(3s+2m+l-3, k-1)$:*

$$(3.18) \quad \begin{aligned} & \# \left(\begin{array}{c|c} j & j \\ \hline j & j+1 \\ \hline j & j+1 \\ \hline j & j+1 \end{array} \right)_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}}^{\frac{\alpha}{\gamma}} \\ & + \# \left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j \\ \hline j & j+1 \end{array} \right)_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}}^{\frac{\alpha}{\gamma}} \\ & = \# \left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j \\ \hline j & j \end{array} \right)_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}}^{\frac{\alpha}{\gamma}} \\ & + \# \left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j+1 \\ \hline j & j+1 \end{array} \right)_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}}^{\frac{\alpha}{\gamma}} \end{aligned}$$

Proof. We define

$$\begin{aligned} \mathbb{A} &= \left\{ (t, \eta, \xi) \in \mathbb{P}^3 \mid r\left(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1) \right) (t, \eta, \xi) = r\left(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k \right) (t, \eta, \xi) = r\left(D \begin{bmatrix} s \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1) \right) (t, \eta, \xi) \right. \\ &= \left. r\left(D \begin{bmatrix} s+m-1 \\ s+m+l-1 \end{bmatrix} \times k \right) (t, \eta, \xi) = r\left(D \begin{bmatrix} s+m \\ s+m+l-1 \end{bmatrix} \times (k-1) \right) (t, \eta, \xi) = r\left(D \begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times (k-1) \right) (t, \eta, \xi) \right\} \end{aligned}$$

and

$$\begin{aligned}
\mathbb{B} &= \left\{ (t, \eta, \xi) \in \mathbb{P}^3 \mid r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1)) (t, \eta, \xi) = r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times k) (t, \eta, \xi) = r(D \begin{bmatrix} s \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1)) (t, \eta, \xi) \right. \\
&= r(D \begin{bmatrix} s \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1)) (t, \eta, \xi) = r(D \begin{bmatrix} s \\ s+m \\ s+m+l \end{bmatrix} \times (k-1)) (t, \eta, \xi) \quad \text{and} \\
&\left. r(D \begin{bmatrix} s+m-1 \\ s+m+l-1 \end{bmatrix} \times k) (t, \eta, \xi) = r(D \begin{bmatrix} s+m \\ s+m+l-1 \end{bmatrix} \times k) (t, \eta, \xi) = r(D \begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k) (t, \eta, \xi) = r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1)) (t, \eta, \xi) + 1 \right\}
\end{aligned}$$

By (3.13) we have, observing that $\psi_1(t, \eta, \xi) \cdot \psi_2(t, \eta, \xi) \cdot \psi_3^{q-2}(t, \eta, \xi)$ is constant on cosets of $\mathbb{P}_{k+s-1} \times \mathbb{P}_{k+s+m-1} \times \mathbb{P}_{k+s+m+l-1}$

$$\begin{aligned}
&\int_{\mathbb{P}^3} \psi_1(t, \eta, \xi) \cdot \psi_2(t, \eta, \xi) \cdot \psi_3^{q-2}(t, \eta, \xi) dt d\eta d\xi \\
&= \int_{\mathbb{A}} \psi_1(t, \eta, \xi) \cdot \psi_2(t, \eta, \xi) \cdot \psi_3^{q-2}(t, \eta, \xi) dt d\eta d\xi + \int_{\mathbb{B}} \psi_1(t, \eta, \xi) \cdot \psi_2(t, \eta, \xi) \cdot \psi_3^{q-2}(t, \eta, \xi) dt d\eta d\xi \\
&= \sum_{(t, \eta, \xi) \in \mathbb{A} \cap \left(\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1} \right)} 2^{q-3} \cdot 2^{(k+3s+2m+l-3-r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1)) q)} \\
&\int_{\mathbb{P}_{s+k-1}} dt \int_{\mathbb{P}_{s+k+m-1}} d\eta \int_{\mathbb{P}_{s+k+m+l-1}} d\xi \\
&+ \sum_{(t, \eta, \xi) \in \mathbb{B} \cap \left(\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1} \right)} -2^{q-3} \cdot 2^{(k+3s+2m+l-3-r(D \begin{bmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{bmatrix} \times (k-1)) q)} \\
&\int_{\mathbb{P}_{s+k-1}} dt \int_{\mathbb{P}_{s+k+m-1}} d\eta \int_{\mathbb{P}_{s+k+m+l-1}} d\xi \\
&= \sum_{j=0}^{\inf(3s+2m+l-3, k-1)} 2^{q-3} \cdot 2^{(k+3s+2m+l-3-j)q} \cdot \# \left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j \end{array} \right)^{\frac{\frac{\alpha}{\beta}}{\gamma}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}} \cdot 2^{-(3k+3s+2m+l-3)} \\
&+ \sum_{j=0}^{\inf(3s+2m+l-3, k-1)} 2^{q-3} \cdot 2^{(k+3s+2m+l-3-j)q} \cdot \# \left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j+1 \end{array} \right)^{\frac{\frac{\alpha}{\beta}}{\gamma}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}} \cdot 2^{-(3k+3s+2m+l-3)} \\
&+ \sum_{j=0}^{\inf(3s+2m+l-3, k-1)} 2^{q-3} \cdot 2^{(k+3s+2m+l-3-j)q} \cdot \# \left(\begin{array}{c|c} j & j \\ \hline j & j+1 \\ \hline j & j+1 \end{array} \right)^{\frac{\frac{\alpha}{\beta}}{\gamma}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}} \cdot 2^{-(3k+3s+2m+l-3)} \\
&- \sum_{j=0}^{\inf(3s+2m+l-3, k-1)} 2^{q-3} \cdot 2^{(k+3s+2m+l-3-j)q} \cdot \# \left(\begin{array}{c|c} j & j \\ \hline j & j+1 \\ \hline j & j+1 \end{array} \right)^{\frac{\frac{\alpha}{\beta}}{\gamma}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}} \cdot 2^{-(3k+3s+2m+l-3)}
\end{aligned}$$

Now by Lemma 3.10 we get for all $q \geq 3$

$$\begin{aligned}
& \sum_{j=0}^{\inf(3s+2m+l-3, k-1)} 2^{-jq} \cdot \left[\# \left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j \end{array} \right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}} \right. \\
& + \# \left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j+1 \end{array} \right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}} + \# \left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j+1 \end{array} \right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}} \\
& \left. - \# \left(\begin{array}{c|c} j & j \\ \hline j & j+1 \\ \hline j & j+1 \end{array} \right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}} \right] = 0.
\end{aligned}$$

□

Lemma 3.14. *We have the following rank identities for $0 \leq j \leq \inf(3s+2m+l-3, k-1)$:*

$$\begin{aligned}
(3.19) \quad & \# \left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j+1 \end{array} \right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}} = \# \left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j \end{array} \right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}} \\
& + \# \left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j+1 \end{array} \right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}}
\end{aligned}$$

and

$$\begin{aligned}
(3.20) \quad & \# \left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j \end{array} \right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}} = \# \left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j+1 \end{array} \right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}}
\end{aligned}$$

Proof. The proofs of (3.19) and (3.20) are somewhat similar to the proof of (3.18) applying respectively (3.16), Lemma 3.11 and (3.17), Lemma 3.12

□

Lemma 3.15. *We have the following rank identities for all integers j for which $0 \leq j \leq \inf(3s+2m+l-3, k-2)$:*

$$(3.21) \quad \# \left(\begin{array}{c|c} j & j+1 \\ \hline j & j+1 \\ \hline j & j+1 \end{array} \right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}} = 0$$

Proof. Proof by contradiction. Assume on the contrary that there exists $j_0 \in [0, \inf(3s + 2m + l - 3, k - 2)]$ such that

$$(3.22) \quad \# \left(\begin{array}{c|c} j_0 & j_0 + 1 \\ \hline j_0 & j_0 + 1 \\ \hline j_0 & j_0 + 1 \\ \hline j_0 & j_0 + 1 \end{array} \right)_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}}^{\frac{\alpha}{\beta} \gamma} > 0.$$

We are going to show that

$$\begin{aligned} & \# \left(\begin{array}{c|c} j_0 & j_0 + 1 \\ \hline j_0 & j_0 + 1 \\ \hline j_0 & j_0 + 1 \\ \hline j_0 & j_0 + 1 \end{array} \right)_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}}^{\frac{\alpha}{\beta} \gamma} > 0 \\ \implies & \# \left(\begin{array}{c|c} j_0 - 1 & j_0 \\ \hline j_0 - 1 & j_0 \\ \hline j_0 - 1 & j_0 \\ \hline j_0 - 1 & j_0 \end{array} \right)_{\mathbb{P}_1/\mathbb{P}_{k+s-1} \times \mathbb{P}_1/\mathbb{P}_{k+s+m-1} \times \mathbb{P}_1/\mathbb{P}_{k+s+m+l-1}}^{\frac{\alpha}{\beta} \gamma} > 0 \\ \implies \dots \implies & \# \left(\begin{array}{c|c} 0 & 1 \\ \hline 0 & 1 \\ \hline 0 & 1 \\ \hline 0 & 1 \end{array} \right)_{\mathbb{P}_{j_0}/\mathbb{P}_{k+s-1} \times \mathbb{P}_{j_0}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}_{j_0}/\mathbb{P}_{k+s+m+l-1}}^{\frac{\alpha}{\beta} \gamma} > 0 \end{aligned}$$

which obviously contradicts

$$\# \left(\begin{array}{c|c} 0 & 1 \\ \hline 0 & 1 \\ \hline 0 & 1 \\ \hline 0 & 1 \end{array} \right)_{\mathbb{P}_{j_0}/\mathbb{P}_{k+s-1} \times \mathbb{P}_{j_0}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}_{j_0}/\mathbb{P}_{k+s+m+l-1}}^{\frac{\alpha}{\beta} \gamma} = 0$$

By (3.22) there exists $(t_0, \eta_0, \xi_0) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}$ such that

$$(3.23) \quad \begin{cases} r(D_{\begin{smallmatrix} s-1 \\ s+m+l-1 \end{smallmatrix}}^{\begin{smallmatrix} s-1 \\ s+m-1 \end{smallmatrix}} \times (k-1)) (t_0, \eta_0, \xi_0) = r(D_{\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix}}^{\begin{smallmatrix} s \\ s+m-1 \end{smallmatrix}} \times (k-1)) (t_0, \eta_0, \xi_0) \\ = r(D_{\begin{smallmatrix} s \\ s+m+l-1 \end{smallmatrix}}^{\begin{smallmatrix} s \\ s+m-1 \end{smallmatrix}} \times (k-1)) (t_0, \eta_0, \xi_0) = r(D_{\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix}}^{\begin{smallmatrix} s \\ s+m-1 \end{smallmatrix}} \times (k-1)) (t_0, \eta_0, \xi_0) = j_0, \\ r(D_{\begin{smallmatrix} s-1 \\ s+m+l-1 \end{smallmatrix}}^{\begin{smallmatrix} s-1 \\ s+m-1 \end{smallmatrix}} \times k) (t_0, \eta_0, \xi_0) = r(D_{\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix}}^{\begin{smallmatrix} s \\ s+m-1 \end{smallmatrix}} \times k) (t_0, \eta_0, \xi_0) \\ = r(D_{\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix}}^{\begin{smallmatrix} s \\ s+m-1 \end{smallmatrix}} \times k) (t_0, \eta_0, \xi_0) = r(D_{\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix}}^{\begin{smallmatrix} s \\ s+m-1 \end{smallmatrix}} \times k) (t_0, \eta_0, \xi_0) = j_0 + 1 \end{cases}$$

Consider the following partition of the matrix

$$D \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{array} \right] \times k \quad (t_0, \eta_0, \xi_0)$$

$$\left(\begin{array}{c|c|c|c|c|c} \alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\ \beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\ \gamma_1 & \gamma_2 & \gamma_3 & \dots & \gamma_{k-1} & \gamma_k \\ \hline \alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\ \beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\ \gamma_2 & \gamma_3 & \gamma_4 & \dots & \gamma_k & \gamma_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\ \beta_{s-1} & \beta_s & \beta_{s+1} & \dots & \beta_{s+k-3} & \beta_{s+k-2} \\ \gamma_{s-1} & \gamma_s & \gamma_{s+1} & \dots & \gamma_{s+k-3} & \gamma_{s+k-2} \\ \beta_s & \beta_{s+1} & \beta_{s+2} & \dots & \beta_{s+k-2} & \beta_{s+k-1} \\ \beta_{s+1} & \beta_{s+2} & \beta_{s+3} & \dots & \beta_{s+k-1} & \beta_{s+k} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \gamma_s & \gamma_{s+1} & \gamma_{s+2} & \dots & \gamma_{s+k-2} & \gamma_{s+k-1} \\ \gamma_{s+1} & \gamma_{s+2} & \gamma_{s+3} & \dots & \gamma_{s+k-1} & \gamma_{s+k} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{s+m+l-1} & \gamma_{s+m+l} & \gamma_{s+m+l+1} & \dots & \gamma_{s+m+l+k-3} & \gamma_{s+m+l+k-2} \\ \hline \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \dots & \alpha_{s+k-2} & \alpha_{s+k-1} \\ \beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \\ \gamma_{s+m+l} & \gamma_{s+m+l+1} & \gamma_{s+m+l+2} & \dots & \gamma_{s+m+l+k-2} & \gamma_{s+m+l+k-1} \end{array} \right).$$

Let j be a rational integer such that $1 \leq j \leq k-1$.

$$\text{Set } (t, \eta, \xi) = (\sum_{i \geq 1} \alpha_i T^{-i}, \sum_{i \geq 1} \beta_i T^{-i}, \sum_{i \geq 1} \gamma_i T^{-i}) \in \mathbb{P}^3$$

Recall that we denote by $D_j^{\begin{smallmatrix} s \\ s+m \end{smallmatrix}} \times (k-j+1)$ any $(3s+2m+l) \times (k-j+1)$ matrix, such that after a rearrangement of the rows, if necessary, we can obtain the following triple

$$\text{persymmetric matrix } \left[\begin{array}{c} D_{s \times (k-j+1)}^j(t) \\ \frac{D_{(s+m) \times (k-j+1)}^j(\eta)}{D_{(s+m+l) \times (k-j+1)}^j(\xi)} \end{array} \right]$$

$$\left(\begin{array}{ccccc|c} \alpha_j & \alpha_{j+1} & \alpha_{j+2} & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_{j+1} & \alpha_{j+2} & \alpha_{j+3} & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{j+s-1} & \alpha_{j+s} & \alpha_{j+s+1} & \dots & \alpha_{k+s-2} & \alpha_{k+s-1} \\ \hline \beta_j & \beta_{j+1} & \beta_{j+2} & \dots & \beta_{k-1} & \beta_k \\ \beta_{j+1} & \beta_{j+2} & \beta_{j+3} & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+j} & \beta_{m+j+1} & \beta_{m+j+2} & \dots & \beta_{k+m-1} & \beta_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m+j-1} & \beta_{s+m+j} & \beta_{s+m+j+1} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \\ \hline \gamma_j & \gamma_{j+1} & \gamma_{j+2} & \dots & \gamma_{k-1} & \gamma_k \\ \gamma_{j+1} & \gamma_{j+2} & \gamma_{j+3} & \dots & \gamma_k & \gamma_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{m+l+j} & \gamma_{m+l+j+1} & \gamma_{m+l+j+2} & \dots & \gamma_{k+m+l-1} & \gamma_{k+m+l} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{s+m+l+j-1} & \gamma_{s+m+l+j} & \gamma_{s+m+l+j+1} & \dots & \gamma_{s+m+l+k-2} & \gamma_{s+m+l+k-1} \end{array} \right).$$

By (3.23) we have $r(D_2^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \end{smallmatrix} \right] \times (k-1)}(t_0, \eta_0, \xi_0)) \leq j_0$, $r(D^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \end{smallmatrix} \right] \times k}(t_0, \eta_0, \xi_0)) = j_0 + 1$.

It follows that $r(D_2^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \end{smallmatrix} \right] \times (k-1)}(t_0, \eta_0, \xi_0)) = j_0$. Let $a_1, a_2, \dots, a_k$ denote the columns of $D^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \end{smallmatrix} \right] \times k}(t_0, \eta_0, \xi_0)$ that is,

$$a_i = \begin{pmatrix} \alpha_i \\ \beta_i \\ \gamma_i \\ \alpha_{i+1} \\ \beta_{i+1} \\ \gamma_{i+1} \\ \vdots \\ \alpha_{i+s-2} \\ \beta_{i+s-2} \\ \gamma_{i+s-2} \\ \beta_{i+s-1} \\ \beta_{i+s} \\ \vdots \\ \beta_{i+s+m-3} \\ \beta_{i+s+m-2} \\ \gamma_{i+s-1} \\ \gamma_{i+s} \\ \vdots \\ \gamma_{i+s+m+l-3} \\ \gamma_{i+s+m+l-2} \end{pmatrix}.$$

Since $r(D_2^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \end{smallmatrix} \right] \times (k-1)}(t_0, \eta_0, \xi_0)) = j_0$ and $r(D^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \end{smallmatrix} \right] \times k}(t_0, \eta_0, \xi_0)) = j_0 + 1$,

we have $a_1 \notin \text{span}\{a_2, a_3, \dots, a_k\}$, therefore $r(D_2^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times (k-2)}(t_0, \eta_0, \xi_0)) = j_0 - 1$
 We have then

$$(3.24) \quad \begin{cases} r(D_2^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times (k-2)}(t_0, \eta_0, \xi_0)) = j_0 - 1, & r(D_2^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times (k-1)}(t_0, \eta_0, \xi_0)) = j_0 \\ r(D^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times (k-1)}(t_0, \eta_0, \xi_0)) = r(D^{\left[\begin{smallmatrix} s \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times (k-1)}(t_0, \eta_0, \xi_0)) \\ = r(D^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l-1 \end{smallmatrix} \right] \times (k-1)}(t_0, \eta_0, \xi_0)) = r(D^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times (k-1)}(t_0, \eta_0, \xi_0)) = j_0. \end{cases}$$

Now we consider the matrix obtained from the matrix

$$D^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \frac{\alpha_{s-1}}{\beta_{s+m-1}} \\ \gamma_{s+m+l-1} \end{smallmatrix} \right] \times k}(t_0, \eta_0, \xi_0)$$

by delating the first column and replacing the last column by the first one :

$$\left(\begin{array}{cccc|c} \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_1 \\ \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_1 \\ \gamma_2 & \gamma_3 & \dots & \gamma_{k-1} & \gamma_1 \\ \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_2 \\ \beta_3 & \beta_4 & \dots & \beta_k & \beta_2 \\ \gamma_3 & \gamma_4 & \dots & \gamma_k & \gamma_2 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-3} & \alpha_{s-1} \\ \beta_s & \beta_{s+1} & \dots & \beta_{s+k-3} & \beta_{s-1} \\ \gamma_s & \gamma_{s+1} & \dots & \gamma_{s+k-3} & \gamma_{s-1} \\ \beta_{s+1} & \beta_{s+2} & \dots & \beta_{s+k-2} & \beta_s \\ \beta_{s+2} & \beta_{s+3} & \dots & \beta_{s+k-1} & \beta_{s+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m-1} \\ \gamma_{s+1} & \gamma_{s+2} & \dots & \gamma_{s+k-2} & \gamma_s \\ \gamma_{s+2} & \gamma_{s+3} & \dots & \gamma_{s+k-1} & \gamma_{s+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{s+m+l} & \gamma_{s+m+l+1} & \dots & \gamma_{s+m+l+k-3} & \gamma_{s+m+l-1} \\ \hline \alpha_{s+1} & \alpha_{s+2} & \dots & \alpha_{s+k-2} & \alpha_s \\ \beta_{s+m+1} & \beta_{s+m+2} & \dots & \beta_{s+m+k-2} & \beta_{s+m} \\ \gamma_{s+m+l+1} & \gamma_{s+m+l+2} & \dots & \gamma_{s+m+l+k-2} & \gamma_{s+m+l} \end{array} \right).$$

From (3.24) we obtain by elementary rank considerations

$$(3.25) \quad \begin{cases} r(D_2^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times (k-2)}(t_0, \eta_0, \xi_0)) = r(D_2^{\left[\begin{smallmatrix} s \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times (k-2)}(t_0, \eta_0, \xi_0)) = \\ r(D^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l-1 \end{smallmatrix} \right] \times (k-2)}(t_0, \eta_0, \xi_0)) = r(D^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times (k-2)}(t_0, \eta_0, \xi_0)) = j_0 - 1 \\ r(D_2^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times (k-1)}(t_0, \eta_0, \xi_0)) = j_0 \end{cases}$$

Consider now the matrix

$$D_2 \begin{bmatrix} \frac{s-1}{s+m-1} \\ \frac{s+m-1}{s+m+l-1} \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{bmatrix} \times (k-1) \quad (t_0, \eta_0, \xi_0)$$

obtained by the matrix

$$D \begin{bmatrix} \frac{s-1}{s+m-1} \\ \frac{s+m-1}{s+m+l-1} \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{bmatrix} \times k \quad (t_0, \eta_0, \xi_0)$$

by delating the first column, that is

$$\left(\begin{array}{ccccc} \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\ \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\ \gamma_2 & \gamma_3 & \dots & \gamma_{k-1} & \gamma_k \\ \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\ \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\ \gamma_3 & \gamma_4 & \dots & \gamma_k & \gamma_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\ \beta_s & \beta_{s+1} & \dots & \beta_{s+k-3} & \beta_{s+k-2} \\ \gamma_s & \gamma_{s+1} & \dots & \gamma_{s+k-3} & \gamma_{s+k-2} \\ \beta_{s+1} & \beta_{s+2} & \dots & \beta_{s+k-2} & \beta_{s+k-1} \\ \beta_{s+2} & \beta_{s+3} & \dots & \beta_{s+k-1} & \beta_{s+k} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \gamma_{s+1} & \gamma_{s+2} & \dots & \gamma_{s+k-2} & \gamma_{s+k-1} \\ \gamma_{s+2} & \gamma_{s+3} & \dots & \gamma_{s+k-1} & \gamma_{s+k} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \hline \gamma_{s+m+l} & \gamma_{s+m+l+1} & \dots & \gamma_{s+m+l+k-3} & \gamma_{s+m+l+k-2} \\ \hline \alpha_{s+1} & \alpha_{s+2} & \dots & \alpha_{s+k-2} & \alpha_{s+k-1} \\ \hline \beta_{s+m+1} & \beta_{s+m+2} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \\ \hline \gamma_{s+m+l+1} & \gamma_{s+m+l+2} & \dots & \gamma_{s+m+l+k-2} & \gamma_{s+m+l+k-1} \end{array} \right).$$

From (3.25) we then deduce

$$\left\{ \begin{array}{l} r(D_2 \begin{bmatrix} \frac{s-1}{s+m-1} \\ \frac{s+m-1}{s+m+l-1} \end{bmatrix} \times (k-1)) (t_0, \eta_0, \xi_0) = r(D_2 \begin{bmatrix} \frac{s-1}{s+m-1} \\ \frac{s+m-1}{s+m+l-1} \end{bmatrix} \times (k-1)) (t_0, \eta_0, \xi_0) = \\ r(D_2 \begin{bmatrix} \frac{s-1}{s+m-1} \\ \frac{s+m-1}{s+m+l-1} \end{bmatrix} \times (k-1)) (t_0, \eta_0, \xi_0) = r(D_2 \begin{bmatrix} \frac{s-1}{s+m-1} \\ \frac{s+m-1}{s+m+l-1} \end{bmatrix} \times (k-1)) (t_0, \eta_0, \xi_0) = j_0 \end{array} \right.$$

We get

$$(3.26) \quad \# \left(\begin{array}{c|c} j_0 - 1 & j_0 \\ \hline j_0 - 1 & j_0 \\ \hline j_0 - 1 & j_0 \\ \hline j_0 - 1 & j_0 \end{array} \right)^{\frac{\alpha}{\gamma}} > 0$$

$\mathbb{P}_1 / \mathbb{P}_{k+s-1} \times \mathbb{P}_1 / \mathbb{P}_{k+s+m-1} \times \mathbb{P}_1 / \mathbb{P}_{k+s+m+l-1}$

We have now proved that

$$\begin{aligned} & \# \left(\begin{array}{c|c} j_0 & j_0 + 1 \\ \hline j_0 & j_0 + 1 \\ \hline j_0 & j_0 + 1 \\ \hline j_0 & j_0 + 1 \end{array} \right)^{\frac{\alpha}{\gamma}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}} > 0 \\ \Rightarrow & \# \left(\begin{array}{c|c} j_0 - 1 & j_0 \\ \hline j_0 - 1 & j_0 \\ \hline j_0 - 1 & j_0 \\ \hline j_0 - 1 & j_0 \end{array} \right)^{\frac{\alpha}{\gamma}}_{\mathbb{P}_1/\mathbb{P}_{k+s-1} \times \mathbb{P}_1/\mathbb{P}_{k+s+m-1} \times \mathbb{P}_1/\mathbb{P}_{k+s+m+l-1}} > 0 \end{aligned}$$

We repeat this procedure and obtain after finitely many steps

$$\# \left(\begin{array}{c|c} 0 & 1 \\ \hline 0 & 1 \\ \hline 0 & 1 \\ \hline 0 & 1 \end{array} \right)^{\frac{\alpha}{\gamma}}_{\mathbb{P}_{j_0}/\mathbb{P}_{k+s-1} \times \mathbb{P}_{j_0}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}_{j_0}/\mathbb{P}_{k+s+m+l-1}} > 0$$

From

$$D_{j_0+1}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{array} \right]} \times (k-j_0) (t_0, \eta_0, \xi_0) = \left(\begin{array}{cccc|c} \alpha_{j_0+1} & \alpha_{j_0+2} & \dots & \alpha_{k-1} & \alpha_k \\ \beta_{j_0+1} & \beta_{j_0+2} & \dots & \beta_{k-1} & \beta_k \\ \gamma_{j_0+1} & \gamma_{j_0+2} & \dots & \gamma_{k-1} & \gamma_k \\ \alpha_{j_0+2} & \alpha_{j_0+3} & \dots & \alpha_k & \alpha_{k+1} \\ \beta_{j_0+2} & \beta_{j_0+3} & \dots & \beta_k & \beta_{k+1} \\ \gamma_{j_0+2} & \gamma_{j_0+3} & \dots & \gamma_k & \gamma_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s+j_0-1} & \alpha_{s+j_0} & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\ \beta_{s+j_0-1} & \beta_{s+j_0} & \dots & \beta_{s+k-3} & \beta_{s+k-2} \\ \gamma_{s+j_0-1} & \gamma_{s+j_0} & \dots & \gamma_{s+k-3} & \gamma_{s+k-2} \\ \beta_{s+j_0} & \beta_{s+j_0+1} & \dots & \beta_{s+k-2} & \beta_{s+k-1} \\ \beta_{s+j_0} & \beta_{s+j_0+1} & \dots & \beta_{k+s-1} & \beta_{k+s} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+j_0+m-1} & \beta_{s+j_0+m} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \gamma_{s+j_0} & \gamma_{s+j_0+1} & \dots & \gamma_{s+k-2} & \gamma_{s+k-1} \\ \gamma_{s+j_0+1} & \gamma_{s+j_0+2} & \dots & \gamma_{s+k-1} & \gamma_{s+k} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \hline \gamma_{s+j_0+m+l-1} & \gamma_{s+j_0+m+l} & \dots & \gamma_{s+m+l+k-3} & \gamma_{s+m+l+k-2} \\ \hline \alpha_{s+j_0} & \alpha_{s+j_0+1} & \dots & \alpha_{s+k-2} & \alpha_{s+k-1} \\ \hline \beta_{s+j_0+m} & \beta_{s+j_0+m+1} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \\ \hline \gamma_{s+j_0+m+l} & \gamma_{s+j_0+m+l+1} & \dots & \gamma_{s+m+l+k-2} & \gamma_{s+m+l+k-1} \end{array} \right),$$

we obviously get

$$\# \left(\begin{array}{c|c} 0 & 1 \\ \hline 0 & 1 \\ \hline 0 & 1 \\ \hline 0 & 1 \end{array} \right)^{\frac{\alpha}{\gamma}}_{\mathbb{P}_{j_0}/\mathbb{P}_{k+s-1} \times \mathbb{P}_{j_0}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}_{j_0}/\mathbb{P}_{k+s+m+l-1}} = 0,$$

which obviously contradicts (3.26) $\square$

4. Study of the remainder $\Delta_i^{\left[\begin{array}{c} s \\ s+m \\ s+m+l \end{array} \right]} \times k$ in the recurrent formula

4.1. **Notation.** We define $\Delta_i \left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k$ to be the sum

$$\sigma_{i,i,i,i} \left[\begin{smallmatrix} \alpha \\ \beta \\ \gamma \end{smallmatrix} \right] \times k - 7 \cdot \sigma_{i-1,i-1,i-1,i-1} \left[\begin{smallmatrix} \alpha \\ \beta \\ \gamma \end{smallmatrix} \right] \times k + 14 \cdot \sigma_{i-2,i-2,i-2,i-2} \left[\begin{smallmatrix} \alpha \\ \beta \\ \gamma \end{smallmatrix} \right] \times k - 8 \cdot \sigma_{i-3,i-3,i-3,i-3} \left[\begin{smallmatrix} \alpha \\ \beta \\ \gamma \end{smallmatrix} \right] \times k$$

Recall that

$$\sigma_{i,i,i,i} \left[\begin{smallmatrix} \alpha \\ \beta \\ \gamma \end{smallmatrix} \right] \times k$$

denotes the cardinality of the following set

$$\left\{ (t, \eta, \xi) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1} \mid r(D \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times k (t, \eta, \xi)) = i \right. \\ \left. r(D \left[\begin{smallmatrix} s \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times k (t, \eta, \xi)) = i, \quad r(D \left[\begin{smallmatrix} s \\ s+m \\ s+m+l-1 \end{smallmatrix} \right] \times k (t, \eta, \xi)) = i, \quad r(D \left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k (t, \eta, \xi)) = i \right\}$$

4.2. **Introduction.** Adapting the methods used in Section 8 of [2] and using the rank formulas established in Section 3, we deduce by elementary rank considerations the following formula for $3 \leq i \leq 3s + 2m + l$, $k \geq i + 1$

$$\Delta_i \left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k = \sum_{j=i-3}^{i+1} a_j \cdot \Gamma_j \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times j$$

where the $a_j \in \mathbf{Z}$ are explicitly determined.

4.3. **A rank formula for** $\sigma_{i,i,i,i} \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{smallmatrix} \right] \times k$.

Lemma 4.1. For $1 \leq i \leq 2s + m + l - 4$, $k \geq i + 1$ we have

$$(4.1) \quad \sigma_{i,i,i,i} \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{smallmatrix} \right] \times k = \sigma_{i,i,i,i} \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{smallmatrix} \right] \times (i+1)$$

Proof. The formula (4.1) is obvious for $k=i+1$.

We consider the following partition of the matrix

$$D \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{smallmatrix} \right] \times k (t, \eta, \xi)$$

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \dots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \dots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\beta_1 & \beta_2 & \dots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \dots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \dots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\gamma_1 & \gamma_2 & \dots & \gamma_{k-1} & \gamma_k \\
\gamma_2 & \gamma_3 & \dots & \gamma_k & \gamma_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\gamma_{m+l+1} & \gamma_{m+l+2} & \dots & \gamma_{k+m+l-1} & \gamma_{k+m+l} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\hline
\gamma_{s+m+l-1} & \gamma_{s+m+l} & \dots & \gamma_{s+m+l+k-3} & \gamma_{s+m+l+k-2} \\
\hline
\alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-2} & \alpha_{s+k-1} \\
\hline
\beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \\
\hline
\gamma_{s+m+l} & \gamma_{s+m+l+1} & \dots & \gamma_{s+m+l+k-2} & \gamma_{s+m+l+k-1}
\end{pmatrix}.$$

Let $k \geq i + 2$. Using Lemma 3.15 we obtain by elementary rank considerations

$$\begin{aligned}
(4.2) \quad & \sigma_{i,i,i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_s \\ \beta_{s+m} \\ \gamma_{s+m+l} \end{array} \right] \times k \\
&= \# \left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i \\ \hline i & i \end{array} \right)_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}}^{\frac{\alpha}{\beta} \gamma} + \# \left(\begin{array}{c|c} i-1 & i \\ \hline i & i \\ \hline i & i \\ \hline i & i \end{array} \right)_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}}^{\frac{\alpha}{\beta} \gamma} \\
&+ \# \left(\begin{array}{c|c} i-1 & i \\ \hline i-1 & i \\ \hline i & i \\ \hline i & i \end{array} \right)_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}}^{\frac{\alpha}{\beta} \gamma} + \# \left(\begin{array}{c|c} i-1 & i \\ \hline i-1 & i \\ \hline i-1 & i \\ \hline i & i \end{array} \right)_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}}^{\frac{\alpha}{\beta} \gamma} \\
&+ \# \left(\begin{array}{c|c} i-1 & i \\ \hline i-1 & i \\ \hline i-1 & i \\ \hline i-1 & i \end{array} \right)_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}}^{\frac{\alpha}{\beta} \gamma} \\
&= \# \left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i \\ \hline i & i \end{array} \right)_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}}^{\frac{\alpha}{\beta} \gamma}
\end{aligned}$$

Further, by Lemmas 3.13, 3.14 and 3.15 we have

$$\begin{aligned}
(4.3) \quad & 8 \cdot \sigma_{i,i,i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{array} \right] \times (k-1) \\
&= \# \left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i \\ \hline i & i \end{array} \right)^{\frac{\alpha}{\gamma}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}} + \# \left(\begin{array}{c|c} i & i \\ \hline i & i+1 \\ \hline i & i+1 \\ \hline i & i+1 \end{array} \right)^{\frac{\alpha}{\gamma}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}} \\
&+ \# \left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i+1 \\ \hline i & i+1 \end{array} \right)^{\frac{\alpha}{\gamma}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}} + \# \left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i \\ \hline i & i+1 \end{array} \right)^{\frac{\alpha}{\gamma}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}} \\
&+ \# \left(\begin{array}{c|c} i & i+1 \\ \hline i & i+1 \\ \hline i & i+1 \\ \hline i & i+1 \end{array} \right)^{\frac{\alpha}{\gamma}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}} \\
&= 8 \cdot \# \left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i \\ \hline i & i \end{array} \right)^{\frac{\alpha}{\gamma}}_{\mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m+l-1}}
\end{aligned}$$

By (4.2),(4.3) we deduce

$$(4.4) \quad \sigma_{i,i,i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{array} \right] \times (k-1) = \sigma_{i,i,i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{array} \right] \times k \quad \text{for all } k \geq i+2$$

Hence by (4.4) we obtain successively

$$\sigma_{i,i,i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{array} \right] \times k = \sigma_{i,i,i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{array} \right] \times (k-1) = \dots = \sigma_{i,i,i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{array} \right] \times (i+1).$$

□

Lemma 4.2. *For all i such that $1 \leq i \leq 3s + 2m + l - 4$ we have*

$$(4.5) \quad \sigma_{i,i,i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{array} \right] \times (i+1) = 8 \cdot \Gamma_i \left[\begin{array}{c} s-1 \\ s-1+m \\ s-1+m+l \end{array} \right] \times i - \Gamma_{i+1} \left[\begin{array}{c} s-1 \\ s-1+m \\ s-1+m+l \end{array} \right] \times (i+1)$$

Proof. We consider the following partition of the matrix

$$D \begin{bmatrix} \begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \alpha_s - \\ \beta_{s+m} - \\ \gamma_{s+m+l} - \end{array} \end{bmatrix} \times (i+1) \quad (t, \eta, \xi)$$

$$\left(\begin{array}{ccccc} \alpha_1 & \alpha_2 & \dots & \alpha_i & \alpha_{i+1} \\ \alpha_2 & \alpha_3 & \dots & \alpha_{i+1} & \alpha_{i+2} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \dots & \alpha_{s+i-2} & \alpha_{s+i-1} \\ \beta_1 & \beta_2 & \dots & \beta_i & \beta_{i+1} \\ \beta_2 & \beta_3 & \dots & \beta_{i+1} & \beta_{i+2} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \dots & \beta_{i+m} & \beta_{i+m+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \dots & \beta_{s+m+i-2} & \beta_{s+m+i-1} \\ \gamma_1 & \gamma_2 & \dots & \gamma_i & \gamma_{i+1} \\ \gamma_2 & \gamma_3 & \dots & \gamma_{i+1} & \gamma_{i+2} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{m+l+1} & \gamma_{m+l+2} & \dots & \gamma_{i+m+l} & \gamma_{i+m+l+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \hline \gamma_{s+m+l-1} & \gamma_{s+m+l} & \dots & \gamma_{s+m+l+i-2} & \gamma_{s+m+l+i-1} \\ \hline \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+i-1} & \alpha_{s+i} \\ \hline \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+i-1} & \beta_{s+m+i} \\ \hline \gamma_{s+m+l} & \gamma_{s+m+l+1} & \dots & \gamma_{s+m+l+i-1} & \gamma_{s+m+l+i} \end{array} \right).$$

By elementary rank considerations and using Lemma 3.15 we obtain

$$(4.6) \quad \sigma_{i,i,i,i} \begin{bmatrix} \begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \alpha_s - \\ \beta_{s+m} - \\ \gamma_{s+m+l} - \end{array} \end{bmatrix} \times (i+1) \\ = \# \left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i \\ \hline i & i \end{array} \right)_{\mathbb{P}/\mathbb{P}_{i+s} \times \mathbb{P}/\mathbb{P}_{i+s+m} \times \mathbb{P}/\mathbb{P}_{i+s+m+l}}^{\frac{\alpha}{\beta}} + \# \left(\begin{array}{c|c} i-1 & i \\ \hline i & i \\ \hline i & i \\ \hline i & i \end{array} \right)_{\mathbb{P}/\mathbb{P}_{i+s} \times \mathbb{P}/\mathbb{P}_{i+s+m} \times \mathbb{P}/\mathbb{P}_{i+s+m+l}}^{\frac{\alpha}{\beta}} \\ + \# \left(\begin{array}{c|c} i-1 & i \\ \hline i-1 & i \\ \hline i & i \\ \hline i & i \end{array} \right)_{\mathbb{P}/\mathbb{P}_{i+s} \times \mathbb{P}/\mathbb{P}_{i+s+m} \times \mathbb{P}/\mathbb{P}_{i+s+m+l}}^{\frac{\alpha}{\beta}} + \# \left(\begin{array}{c|c} i-1 & i \\ \hline i-1 & i \\ \hline i-1 & i \\ \hline i & i \end{array} \right)_{\mathbb{P}/\mathbb{P}_{i+s} \times \mathbb{P}/\mathbb{P}_{i+s+m} \times \mathbb{P}/\mathbb{P}_{i+s+m+l}}^{\frac{\alpha}{\beta}} \\ + \# \left(\begin{array}{c|c} i-1 & i \\ \hline i-1 & i \\ \hline i-1 & i \\ \hline i-1 & i \end{array} \right)_{\mathbb{P}/\mathbb{P}_{i+s} \times \mathbb{P}/\mathbb{P}_{i+s+m} \times \mathbb{P}/\mathbb{P}_{i+s+m+l}}^{\frac{\alpha}{\beta}}$$

$$= \# \left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i \\ \hline i & i \end{array} \right)^{\frac{\alpha}{\beta}}_{\gamma} \mathbb{P}/\mathbb{P}_{i+s} \times \mathbb{P}/\mathbb{P}_{i+s+m} \times \mathbb{P}/\mathbb{P}_{i+s+m+l}$$

Further by by Lemmas 3.13, 3.14 and 3.15 we have

$$\begin{aligned}
(4.7) \quad & 8 \cdot \sigma_{i,i,i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{array} \right] \times i = 8^2 \cdot \Gamma_i \left[\begin{array}{c} s-1 \\ s-1+m \\ s-1+m+l \end{array} \right] \times i \\
& = \# \left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i \\ \hline i & i \end{array} \right)^{\frac{\alpha}{\beta}}_{\gamma} \mathbb{P}/\mathbb{P}_{i+s} \times \mathbb{P}/\mathbb{P}_{i+s+m} \times \mathbb{P}/\mathbb{P}_{i+s+m+l} \\
& \quad + \# \left(\begin{array}{c|c} i & i \\ \hline i & i+1 \\ \hline i & i+1 \\ \hline i & i+1 \end{array} \right)^{\frac{\alpha}{\beta}}_{\gamma} \mathbb{P}/\mathbb{P}_{i+s} \times \mathbb{P}/\mathbb{P}_{i+s+m} \times \mathbb{P}/\mathbb{P}_{i+s+m+l} \\
& \quad + \# \left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i+1 \\ \hline i & i+1 \end{array} \right)^{\frac{\alpha}{\beta}}_{\gamma} \mathbb{P}/\mathbb{P}_{i+s} \times \mathbb{P}/\mathbb{P}_{i+s+m} \times \mathbb{P}/\mathbb{P}_{i+s+m+l} \\
& \quad + \# \left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i+1 \\ \hline i & i+1 \end{array} \right)^{\frac{\alpha}{\beta}}_{\gamma} \mathbb{P}/\mathbb{P}_{i+s} \times \mathbb{P}/\mathbb{P}_{i+s+m} \times \mathbb{P}/\mathbb{P}_{i+s+m+l} \\
& = 8 \cdot \# \left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i \\ \hline i & i \end{array} \right)^{\frac{\alpha}{\beta}}_{\gamma} \mathbb{P}/\mathbb{P}_{i+s} \times \mathbb{P}/\mathbb{P}_{i+s+m} \times \mathbb{P}/\mathbb{P}_{i+s+m+l} \\
& \quad + 8 \cdot \Gamma_{i+1} \left[\begin{array}{c} s-1 \\ s-1+m \\ s-1+m+l \end{array} \right] \times (i+1)
\end{aligned}$$

Combining (4.6), (4.7) we deduce

$$8^2 \cdot \Gamma_i \left[\begin{array}{c} s-1 \\ s-1+m \\ s-1+m+l \end{array} \right] \times i = 8 \cdot \sigma_{i,i,i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{array} \right] \times (i+1) + 8 \cdot \Gamma_{i+1} \left[\begin{array}{c} s-1 \\ s-1+m \\ s-1+m+l \end{array} \right] \times (i+1)$$

and (4.5) is established. $\square$

Lemma 4.3. *Let $s \geq 2$ and $m \geq 0$, $l \geq 0$ we have in the following two cases :*

The case $1 \leq k \leq 3s + 2m + l - 3$

$$(4.8) \quad \sigma_{i,i,i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{array} \right] \times k = \begin{cases} 1 & \text{if } i = 0, \quad k \geq 1, \\ 8 \cdot \Gamma_i \left[\begin{array}{c} s-1 \\ s-1+m \\ s-1+m+l \end{array} \right] \times i - \Gamma_{i+1} \left[\begin{array}{c} s-1 \\ s-1+m \\ s-1+m+l \end{array} \right] \times (i+1) & \text{if } 1 \leq i \leq k-1, \\ 8 \cdot \Gamma_k \left[\begin{array}{c} s-1 \\ s-1+m \\ s-1+m+l \end{array} \right] \times k & \text{if } i = k. \end{cases}$$

The case $k \geq 3s + 2m + l - 3$

$$(4.9) \quad \sigma_{i,i,i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{array} \right] \times k = \begin{cases} 1 & \text{if } i = 0, \\ 8 \cdot \Gamma_i \left[\begin{array}{c} s-1 \\ s-1+m \\ s-1+m+l \end{array} \right] \times i - \Gamma_{i+1} \left[\begin{array}{c} s-1 \\ s-1+m \\ s-1+m+l \end{array} \right] \times (i+1) & \text{if } 1 \leq i \leq 3s + 2m + l - 4, \\ 8 \cdot \Gamma_{3s+2m+l-3} \left[\begin{array}{c} s-1 \\ s-1+m \\ s-1+m+l \end{array} \right] \times (3s+2m+l-3) & \text{if } i = 3s + 2m + l - 3. \end{cases}$$

Proof. Combining (4.1) and (4.5) and recalling the definition of $\sigma_{i,i,i,i} \left[\begin{array}{c} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{array} \right] \times k$ we deduce easily Lemma 4.3 $\square$

4.4. A rank formula for $\Delta_i \left[\begin{array}{c} s \\ s+m \\ s+m+l \end{array} \right] \times k$.

Lemma 4.4. The remainder $\Delta_i \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] \times k$ in the recurrent formula is equal to

$$\begin{aligned}
 & \left. \begin{aligned}
 & 1 \\
 & 8 \cdot \Gamma_1 \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times 1 - \Gamma_2 \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times 2 - 7 \\
 & 8 \cdot \Gamma_1 \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times 1 - 7 \\
 & 15 \cdot \Gamma_2 \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times 2 - 56 \cdot \Gamma_1 \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times 1 - \Gamma_3 \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times 3 + 14 \\
 & 15 \cdot \Gamma_2 \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times 2 - 56 \cdot \Gamma_1 \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times 1 + 14 \\
 & 15 \cdot \Gamma_3 \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times 3 - 80 \cdot \Gamma_2 \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times 2 + 112 \cdot \Gamma_1 \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times 1 \\
 & - \Gamma_4 \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times 4 - 8 \\
 & 15 \cdot \Gamma_3 \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times 3 - 80 \cdot \Gamma_2 \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times 2 + 112 \cdot \Gamma_1 \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times 1 - 8 \\
 & 15 \cdot \Gamma_i \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times i - 80 \cdot \Gamma_{i-1} \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times (i-1) + 120 \cdot \Gamma_{i-2} \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times (i-2) \\
 & - 64 \cdot \Gamma_{i-3} \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times (i-3) - \Gamma_{i+1} \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times (i+1) \\
 & 15 \cdot \Gamma_i \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times i - 80 \cdot \Gamma_{i-1} \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times (i-1) + 120 \cdot \Gamma_{i-2} \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times (i-2) \\
 & - 64 \cdot \Gamma_{i-3} \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times (i-3) \\
 & 15 \cdot \Gamma_{3s+2m+l-3} \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times (3s+2m+l-3) - 80 \cdot \Gamma_{3s+2m+l-4} \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times (3s+2m+l-4) \\
 & + 120 \cdot \Gamma_{3s+2m+l-5} \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times (3s+2m+l-5) - 64 \cdot \Gamma_{3s+2m+l-6} \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times (3s+2m+l-6) \\
 & - 80 \cdot \Gamma_{3s+2m+l-3} \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times (3s+2m+l-3) \\
 & + 120 \cdot \Gamma_{3s+2m+l-4} \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times (3s+2m+l-4) - 64 \cdot \Gamma_{3s+2m+l-5} \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times (3s+2m+l-5) \\
 & 120 \cdot \Gamma_{3s+2m+l-3} \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times (3s+2m+l-3) - 64 \cdot \Gamma_{3s+2m+l-4} \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times (3s+2m+l-4) \\
 & - 64 \cdot \Gamma_{3s+2m+l-3} \left[\begin{smallmatrix} s-1 \\ s-1+m+l \end{smallmatrix} \right] \times (3s+2m+l-3)
 \end{aligned} \right\}
 \end{aligned}$$

if $i = 0, k \geq 1$,
 if $i = 1, k \geq 2$,
 if $i = 1, k = 1$,
 if $i = 2, k \geq 3$,
 if $i = 2, k = 2$,
 if $i = 3, k \geq 4$,
 if $i = 3, k = 3$,
 if $4 \leq i \leq 3s + 2m + l - 4, k \geq i + 1$,
 if $4 \leq i \leq 3s + 2m + l - 4, k = i$,
 if $i = 3s + 2m + l - 3, k \geq i$,
 if $i = 3s + 2m + l - 2, k \geq i$,
 if $i = 3s + 2m + l - 1, k \geq i$,
 if $i = 3s + 2m + l, k \geq i$.

Proof. From (4.8) and (4.9) and recalling that $\Delta_i \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] \times k$ is equal to

$$\begin{aligned}
 & \sigma_{i,i,i,i} \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{smallmatrix} \right] \times k - 7 \cdot \sigma_{i-1,i-1,i-1,i-1} \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{smallmatrix} \right] \times k \\
 & + 14 \cdot \sigma_{i-2,i-2,i-2,i-2} \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{smallmatrix} \right] \times k - 8 \cdot \sigma_{i-3,i-3,i-3,i-3} \left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \\ \alpha_{s-} \\ \beta_{s+m-} \\ \gamma_{s+m+l-} \end{smallmatrix} \right] \times k
 \end{aligned}$$

we deduce easily Lemma 4.4 □

Lemma 4.5. *We have*

$$(4.11) \quad \Delta_i \begin{bmatrix} s+m \\ s+m+l \end{bmatrix}^{\times k} = \Delta_i \begin{bmatrix} s+m \\ s+m+l \end{bmatrix}^{\times (i+1)} \quad \text{for } 0 \leq i \leq \inf(3s+2m+l-4, k-1)$$

$$(4.12) \quad \Delta_i \begin{bmatrix} s+m \\ s+m+l \end{bmatrix}^{\times k} = \Delta_i \begin{bmatrix} s+m \\ s+m+l \end{bmatrix}^{\times i} \quad \text{for } i \in \{3s+2m+l-3, 3s+2m+l-2, 3s+2m+l-1, 3s+2m+l\}, k \geq i$$

Proof. Follows immediately from Lemma 4.4. $\square$

5. A recurrent formula for $\omega_i(s, m, l, k)$

5.1. Notation.

Definition 5.1. Set for $0 \leq i \leq \inf(3s+2m+l, k)$

$$\omega_i(s, m, l, k) = \Gamma_i \begin{bmatrix} s+m \\ s+m+l \end{bmatrix}^{\times k} - 4 \cdot \Gamma_{i-1} \begin{bmatrix} s+m-1 \\ s+m+l \end{bmatrix}^{\times k} - 8 \cdot \Gamma_{i-1} \begin{bmatrix} s+m \\ s+m+l-1 \end{bmatrix}^{\times k} + 32 \cdot \Gamma_{i-2} \begin{bmatrix} s+m-1 \\ s+m+l-1 \end{bmatrix}^{\times k}$$

5.2. Introduction. In this section we prove that $\omega_i(s, m, l, k)$ can be partly obtained from $\omega_i(s, m, l, k)$ with $i \rightarrow i - (s-1)$, $s \rightarrow 1$, $m \rightarrow m + s - 1$, $l \rightarrow l$, $k \rightarrow k$, that is from the sum

$$\Gamma_{i-s+1} \begin{bmatrix} s+m \\ s+m+l \end{bmatrix}^{\times k} - 4 \cdot \Gamma_{i-s} \begin{bmatrix} s+m-1 \\ s+m+l \end{bmatrix}^{\times k} - 8 \cdot \Gamma_{i-s} \begin{bmatrix} s+m \\ s+m+l-1 \end{bmatrix}^{\times k} + 32 \cdot \Gamma_{i-s-1} \begin{bmatrix} s+m-1 \\ s+m+l-1 \end{bmatrix}^{\times k}.$$

These sums were studied in the first section.

5.3. The recursive formula for $\omega_i(s, m, l, k)$.

Lemma 5.2. *Let $s \geq 4, m \geq 0, l \geq 0$ and $k \geq 1$, we have in the following two cases:*

$$(5.1) \quad \omega_i(s, m, l, k) = \begin{cases} 2^{i-3} \cdot \omega_3(s - (i-3), m + (i-3), l, k) + \sum_{j=0}^{i-4} 2^j \Delta_{i-j} \begin{bmatrix} s-j \\ s+m \\ s+m+l \end{bmatrix}^{\times k} & \text{if } 4 \leq i \leq \inf(s-1, k) \\ 2^{s-1} \cdot \omega_{i-(s-1)}(1, m + s - 1, l, k) + \sum_{j=0}^{s-2} 2^j \Delta_{i-j} \begin{bmatrix} s-j \\ s+m \\ s+m+l \end{bmatrix}^{\times k} & \text{if } s \leq i \leq \inf(3s+2m+l, k) \end{cases}$$

Proof. From the recurrent formula (2.83) we obtain the following equivalence

$$(5.2) \quad \omega_i(s, m, l, k) = 2 \cdot \omega_{i-1}(s-1, m+1, l, k) + \Delta_i \begin{bmatrix} s+m \\ s+m+l \end{bmatrix}^{\times k}$$

The first case $4 \leq i \leq \inf(s-1, k)$

Using successively (5.2) we get

$$\begin{aligned} \omega_i(s, m, l, k) &= 2 \cdot \omega_{i-1}(s-1, m+1, l, k) + \Delta_i \begin{bmatrix} s+m \\ s+m+l \end{bmatrix}^{\times k} \\ 2 \cdot \omega_{i-1}(s-1, m+1, l, k) &= 2 \cdot \left(2 \cdot \omega_{i-2}(s-2, m+2, l, k) + \Delta_{i-1} \begin{bmatrix} s-1 \\ s+m \\ s+m+l \end{bmatrix}^{\times k} \right) \\ &\vdots \\ 2^j \cdot \omega_{i-j}(s-j, m+j, l, k) &= 2^j \cdot \left(2 \cdot \omega_{i-(j+1)}(s-(j+1), m+(j+1), l, k) + \Delta_{i-j} \begin{bmatrix} s-j \\ s+m \\ s+m+l \end{bmatrix}^{\times k} \right) \end{aligned}$$

$$\begin{aligned} & \vdots \\ 2^{i-4} \cdot \omega_{i-(i-4)}(s-(i-4), m+(i-4), l, k) &= 2^{i-4} \cdot \left(2 \cdot \omega_{i-(i-3)}(s-(i-3), m+(i-3), l, k) + \Delta_{i-(i-4)}^{\left[\begin{smallmatrix} s-(i-4) \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k} \right) \end{aligned}$$

By summing the above equations we obtain:

$$\begin{aligned} \sum_{j=0}^{i-4} 2^j \omega_{i-j}(s-j, m+j, l, k) &= \sum_{j=0}^{i-4} 2^j \cdot \left(2 \cdot \omega_{i-(j+1)}(s-(j+1), m+(j+1), l, k) + \Delta_{i-j}^{\left[\begin{smallmatrix} s-j \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k} \right) \\ \iff \omega_i(s, m, l, k) &= 2^{i-3} \cdot \omega_3(s-(i-3), m+(i-3), l, k) + \sum_{j=0}^{i-4} 2^j \Delta_{i-j}^{\left[\begin{smallmatrix} s-j \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k} \end{aligned}$$

The second case $s \leq i \leq \inf(3s+2m+l, k)$

$$\begin{aligned} \omega_i(s, m, l, k) &= 2 \cdot \omega_{i-1}(s-1, m+1, l, k) + \Delta_i^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k} \\ 2 \cdot \omega_{i-1}(s-1, m+1, l, k) &= 2 \cdot \left(2 \cdot \omega_{i-2}(s-2, m+2, l, k) + \Delta_{i-1}^{\left[\begin{smallmatrix} s-1 \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k} \right) \\ 2^2 \cdot \omega_{i-2}(s-2, m+2, l, k) &= 2^2 \cdot \left(2 \cdot \omega_{i-3}(s-3, m+3, l, k) + \Delta_{i-2}^{\left[\begin{smallmatrix} s-2 \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k} \right) \\ &\vdots \\ 2^{s-3} \cdot \omega_{i-(s-3)}(s-(s-3), m+(s-3), l, k) &= 2^{s-3} \cdot \left(2 \cdot \omega_{i-(s-2)}(s-(s-2), m+(s-2), l, k) + \Delta_{i-(s-3)}^{\left[\begin{smallmatrix} s-(s-3) \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k} \right) \\ &\vdots \\ 2^{s-2} \cdot \omega_{i-(s-2)}(s-(s-2), m+(s-2), l, k) &= 2^{s-2} \cdot \left(2 \cdot \omega_{i-(s-1)}(s-(s-1), m+(s-1), l, k) + \Delta_{i-(s-2)}^{\left[\begin{smallmatrix} s-(s-2) \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k} \right) \end{aligned}$$

By summing the above equations we obtain:

$$\begin{aligned} \sum_{j=0}^{s-2} 2^j \omega_{i-j}(s-j, m+j, l, k) &= \sum_{j=0}^{s-2} 2^j \cdot \left(2 \cdot \omega_{i-(j+1)}(s-(j+1), m+(j+1), l, k) + \Delta_{i-j}^{\left[\begin{smallmatrix} s-j \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k} \right) \\ \iff \omega_i(s, m, l, k) &= 2^{s-1} \cdot \omega_{i-(s-1)}(1, m+s-1, l, k) + \sum_{j=0}^{s-2} 2^j \Delta_{i-j}^{\left[\begin{smallmatrix} s-j \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k} \end{aligned}$$

□

6. **COMPUTATION OF $\Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k}$ for $i \leq s-1$, $i \leq k$, $m \geq 0$, $l \geq 0$**

6.1. Notation. We define γ_j to be equal to the number $\Gamma_j^{\begin{bmatrix} j \\ j \\ j \end{bmatrix} \times (j+1)}$ of rank j matrices of the form $\begin{bmatrix} A \\ B \\ C \end{bmatrix}$, where A, B and C are $j \times (j+1)$ persymmetric matrices.

6.2. Introduction. In this section we adapt the method used in Section 11 of [2] to compute by induction on i the number $\Gamma_i^{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k}$ of rank i ($\leq s-1$) matrices of the form $\begin{bmatrix} A \\ B \\ C \end{bmatrix}$, where A, B and C are persymmetric.

6.3. Computation of $\Gamma_i^{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k}$ for $i \leq s-1$, $i \leq k$, $m \geq 0$, $l \geq 0$.

Lemma 6.1. Consider the matrix

$$D_{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k}(t, \eta, \xi) = \begin{pmatrix} \alpha_1 & \alpha_2 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\ \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-2} & \alpha_{s+k-1} \\ \beta_1 & \beta_2 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \dots & \beta_{k+m-1} & \beta_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \\ \gamma_1 & \gamma_2 & \dots & \gamma_{k-1} & \gamma_k \\ \gamma_2 & \gamma_3 & \dots & \gamma_k & \gamma_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{m+l+1} & \gamma_{m+l+2} & \dots & \gamma_{k+m+l-1} & \gamma_{k+m+l} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{s+m+l} & \gamma_{s+m+l+1} & \dots & \gamma_{s+m+l+k-2} & \gamma_{s+m+l+k-1} \end{pmatrix}.$$

Let $g_{k,s,m,l}(t, \eta, \xi)$ be the quadratic exponential sum in $\mathbb{P}^3$ defined by

$$(t, \eta, \xi) \in \mathbb{P}^3 \longmapsto \sum_{\deg Y \leq k-1} \sum_{\deg Z \leq s-1} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V \leq s+m+l-1} E(\xi YV) \in \mathbb{Z}.$$

We have

$$(6.1) \quad \sum_{i=0}^{\inf(3s+2m+l, k)} \Gamma_i^{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k} = 2^{3k+3s+2m+l-3}$$

and

$$(6.2) \quad \sum_{i=0}^{\inf(3s+2m+l, k)} \Gamma_i^{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k} \cdot 2^{-i} = 2^{2k+3s+2m+l-3} + 2^{3k-3} - 2^{2k-3}.$$

Proof. The proof of (6.1) is obvious.

By (2.6) with $s-2 \rightarrow s-1$, $s+m-2 \rightarrow s+m-1$, $s+m+l-2 \rightarrow s+m+l-1$ we obtain, observing that $g_{k,s,m,l}(t, \eta, \xi)$ is constant on cosets of $\mathbb{P}_{k+s-1} \times \mathbb{P}_{k+s+m-1} \times \mathbb{P}_{k+s+m+l-1}$

$$\begin{aligned}
& \int_{\mathbb{P}^3} g_{k,s,m,l}(t, \eta, \xi) dt d\eta d\xi \\
&= \sum_{(t, \eta, \xi) \in \mathbb{P}_{k+s-1} \times \mathbb{P}_{k+s+m-1} \times \mathbb{P}_{k+s+m+l-1}} 2^{3s+2m+l+k-r(D \begin{bmatrix} s \\ s+m+l \end{bmatrix}) \times k} (t, \eta, \xi) \int_{\mathbb{P}_{k+s-1}} dt \int_{\mathbb{P}_{k+s+m-1}} d\eta \int_{\mathbb{P}_{k+s+m+l-1}} d\xi \\
&= \sum_{i=0}^{\inf(3s+2m+l, k)} \Gamma_i \begin{bmatrix} s \\ s+m+l \end{bmatrix} \times k \cdot 2^{3s+2m+l+k-i} \int_{\mathbb{P}_{k+s-1}} dt \int_{\mathbb{P}_{k+s+m-1}} d\eta \int_{\mathbb{P}_{k+s+m+l-1}} d\xi \\
&= 2^{-2k+3} \cdot \sum_{i=0}^{\inf(3s+2m+l, k)} \Gamma_i \begin{bmatrix} s \\ s+m+l \end{bmatrix} \times k \cdot 2^{-i}.
\end{aligned}$$

On the other hand

$$\begin{aligned}
& \int_{\mathbb{P}^3} g_{k,s,m,l}(t, \eta, \xi) dt d\eta d\xi \\
&= \text{Card} \{ (Y, Z, U), \deg Y \leq k-1, \deg Z \leq s-1, \deg U \leq s+m-1, \deg V \leq s+m+l-1 \mid Y \cdot Z = Y \cdot U = Y \cdot V = 0 \} \\
&= 2^{3s+2m+l} + 2^k - 1.
\end{aligned}$$

The above equations imply (6.2). $\square$

Lemma 6.2. *We have for all $j \geq 1$*

$$(6.3) \quad (H_j) \quad \Gamma_j \begin{bmatrix} s \\ s+m+l \end{bmatrix} \times k = \Gamma_j \begin{bmatrix} j \\ j \end{bmatrix} \times (j+1) \stackrel{\text{def}}{=} \gamma_j \quad \text{for } s \geq j+1, k \geq j+1, m \geq 0, l \geq 0$$

Proof. The proof is by strong induction, that is:

If

- (H_1) is true, and
- for all $j \geq 1$, $(H_1) \wedge (H_2) \wedge \dots \wedge (H_j)$ implies (H_{j+1})

then (H_j) is true for all $j \geq 1$.

(H_1) is true

$$\begin{aligned}
& \text{Indeed from (2.83) with } i = 1 \text{ and (4.11)} \\
& \Gamma_1 \begin{bmatrix} s \\ s+m+l \end{bmatrix} \times (k+1) - \Gamma_1 \begin{bmatrix} s \\ s+m+l \end{bmatrix} \times k = 14 + \Delta_1 \begin{bmatrix} s \\ s+m+l \end{bmatrix} \times (k+1) - (14 + \Delta_1 \begin{bmatrix} s \\ s+m+l \end{bmatrix} \times k) = 0 \text{ for all} \\
& s \geq 2, k \geq 2, m \geq 0, l \geq 0
\end{aligned}$$

which implies that

$$(6.4) \quad \Gamma_1 \begin{bmatrix} s \\ s+m+l \end{bmatrix} \times k = \Gamma_1 \begin{bmatrix} s \\ s+m+l \end{bmatrix} \times 2 \quad \text{for all } s \geq 2, k \geq 2, m \geq 0, l \geq 0.$$

Consider the matrix

$$\begin{pmatrix} \alpha_1 & \alpha_2 \\ \alpha_2 & \alpha_3 \\ \vdots & \vdots \\ \alpha_s & \alpha_{s+1} \\ \beta_1 & \beta_2 \\ \beta_2 & \beta_3 \\ \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} \\ \vdots & \vdots \\ \beta_{s+m} & \beta_{s+m+1} \\ \gamma_1 & \gamma_2 \\ \gamma_2 & \gamma_3 \\ \vdots & \vdots \\ \gamma_{m+l+1} & \gamma_{m+l+2} \\ \vdots & \vdots \\ \gamma_{s+m+l} & \gamma_{s+m+l+1} \end{pmatrix}.$$

We have respectively by (6.1), (6.2) with $k = 2$

$$(6.5) \quad \sum_{i=0}^2 \Gamma_i \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right]^{\times 2} = 2^{3s+2m+l+3},$$

$$(6.6) \quad \sum_{i=0}^2 \Gamma_i \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right]^{\times 2} \cdot 2^{-i} = 2^{3s+2m+l+1} + 2^3 - 2.$$

From (6.5) and (6.6) using (6.4) we deduce

$$(6.7) \quad \Gamma_1 \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right]^{\times k} = \Gamma_1 \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right]^{\times 2} = 21 \quad \text{for all } s \geq 2, k \geq 2, m \geq 0, l \geq 0$$

$$(6.8) \quad \Gamma_2 \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right]^{\times 2} = 2^{3s+2m+l+3} - 22 \quad \text{for all } s \geq 2, k = 2, m \geq 0, l \geq 0$$

$(H_1) \Rightarrow (H_2)$

From (2.83) with $i = 2$ we obtain

$$\Gamma_2 \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right]^{\times k} = 2 \cdot \Gamma_1 \left[\begin{smallmatrix} s-1 \\ s+m+l \end{smallmatrix} \right]^{\times k} + 4 \cdot \Gamma_1 \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right]^{\times k} + 8 \cdot \Gamma_1 \left[\begin{smallmatrix} s \\ s+m+l-1 \end{smallmatrix} \right]^{\times k} - 56 + \Delta_2 \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right]^{\times k}$$

Applying (4.11) with $i=2$ and $k \geq 3$ we get using (6.7) with $s \geq 3, m \geq 0, l \geq 0$

$$\begin{aligned} & \Gamma_2 \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right]^{\times (k+1)} - \Gamma_2 \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right]^{\times k} = 2 \cdot \left[\Gamma_1 \left[\begin{smallmatrix} s-1 \\ s+m+l \end{smallmatrix} \right]^{\times (k+1)} - \Gamma_1 \left[\begin{smallmatrix} s-1 \\ s+m+l \end{smallmatrix} \right]^{\times k} \right] \\ & + 4 \cdot \left[\Gamma_1 \left[\begin{smallmatrix} s+m-1 \\ s+m+l \end{smallmatrix} \right]^{\times (k+1)} - \Gamma_1 \left[\begin{smallmatrix} s+m-1 \\ s+m+l \end{smallmatrix} \right]^{\times k} \right] + 8 \cdot \left[\Gamma_1 \left[\begin{smallmatrix} s \\ s+m+l-1 \end{smallmatrix} \right]^{\times (k+1)} - \Gamma_1 \left[\begin{smallmatrix} s \\ s+m+l-1 \end{smallmatrix} \right]^{\times k} \right] = 0 \end{aligned}$$

which implies

$$(6.9) \quad \Gamma_2^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k} = \Gamma_2^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times 3} \quad \text{for all } k \geq 3, s \geq 3, m \geq 0, l \geq 0$$

Consider the matrix

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 \\ \alpha_2 & \alpha_3 & \alpha_4 \\ \vdots & \vdots & \vdots \\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} \\ \beta_1 & \beta_2 & \beta_3 \\ \beta_2 & \beta_3 & \beta_4 \\ \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \beta_{m+3} \\ \vdots & \vdots & \vdots \\ \beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} \\ \gamma_1 & \gamma_2 & \gamma_3 \\ \gamma_2 & \gamma_3 & \gamma_4 \\ \vdots & \vdots & \vdots \\ \gamma_{m+l+1} & \gamma_{m+l+2} & \gamma_{m+l+3} \\ \vdots & \vdots & \vdots \\ \gamma_{s+m+l} & \gamma_{s+m+l+1} & \gamma_{s+m+l+2} \end{pmatrix}.$$

We have respectively by (6.1), (6.2) with $k = 3$

$$(6.10) \quad \sum_{i=0}^3 \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times 3} = 2^{3s+2m+l+6},$$

$$(6.11) \quad \sum_{i=0}^3 \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times 3} \cdot 2^{-i} = 2^{3s+2m+l+3} + 2^6 - 2^3.$$

From (6.10) and (6.11) using (6.7) and (6.9) we deduce

$$(6.12) \quad \Gamma_2^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k} = \Gamma_2^{\left[\begin{smallmatrix} 2 \\ 2 \end{smallmatrix} \right] \times 3} = 378 \quad \text{for all } s \geq 3, k \geq 3, m \geq 0, l \geq 0$$

$$(6.13) \quad \Gamma_3^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times 3} = 2^{3s+2m+l+6} - 400 \quad \text{for all } s \geq 3, k = 3, m \geq 0, l \geq 0$$

$(H_1) \wedge (H_2) \text{ implies } (H_3)$
From (2.83) with $i = 3$ we obtain

$$\begin{aligned} & \Gamma_3^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k} \\ &= \left[2 \cdot \Gamma_2^{\left[\begin{smallmatrix} s-1 \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k} + 4 \cdot \Gamma_2^{\left[\begin{smallmatrix} s \\ s+m-1 \\ s+m+l \end{smallmatrix} \right] \times k} + 8 \cdot \Gamma_2^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l-1 \end{smallmatrix} \right] \times k} \right] - \left[8 \cdot \Gamma_1^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l \end{smallmatrix} \right] \times k} + 16 \cdot \Gamma_1^{\left[\begin{smallmatrix} s-1 \\ s+m \\ s+m+l-1 \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_1^{\left[\begin{smallmatrix} s \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times k} \right] \end{aligned}$$

$$+ 64 \cdot \Gamma_0^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times k} + \Delta_3^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k}$$

Applying (4.11) with $i=3$ and $k \geq 4$ we get using (6.7) and (6.12) with $s \geq 4$, $m \geq 0$, $l \geq 0$

$$\begin{aligned} & \Gamma_3^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times (k+1)} - \Gamma_3^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k} = 2 \cdot \left[\Gamma_2^{\left[\begin{smallmatrix} s-1 \\ s+m \\ s+m+l \end{smallmatrix} \right] \times (k+1)} - \Gamma_2^{\left[\begin{smallmatrix} s-1 \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k} \right] \\ & + 4 \cdot \left[\Gamma_2^{\left[\begin{smallmatrix} s+m-1 \\ s+m+l \end{smallmatrix} \right] \times (k+1)} - \Gamma_2^{\left[\begin{smallmatrix} s+m-1 \\ s+m+l \end{smallmatrix} \right] \times k} \right] + 8 \cdot \left[\Gamma_2^{\left[\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix} \right] \times (k+1)} - \Gamma_2^{\left[\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix} \right] \times k} \right] \\ & - 8 \cdot \left[\Gamma_1^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l \end{smallmatrix} \right] \times (k+1)} - \Gamma_1^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \\ s+m+l \end{smallmatrix} \right] \times k} \right] - 16 \cdot \left[\Gamma_1^{\left[\begin{smallmatrix} s-1 \\ s+m \\ s+m+l-1 \end{smallmatrix} \right] \times (k+1)} - \Gamma_1^{\left[\begin{smallmatrix} s-1 \\ s+m \\ s+m+l-1 \end{smallmatrix} \right] \times k} \right] \\ & - 32 \cdot \left[\Gamma_1^{\left[\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times (k+1)} - \Gamma_1^{\left[\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times k} \right] = 0 \end{aligned} \quad \text{which implies}$$

$$(6.14) \quad \Gamma_3^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k} = \Gamma_3^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times 4} \quad \text{for all } k \geq 4, s \geq 4, m \geq 0, l \geq 0$$

Consider the matrix

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 \\ \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \alpha_{s+3} \\ \beta_1 & \beta_2 & \beta_3 & \beta_4 \\ \beta_2 & \beta_3 & \beta_4 & \beta_5 \\ \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \beta_{m+4} \\ \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \beta_{s+m+3} \\ \gamma_1 & \gamma_2 & \gamma_3 & \gamma_4 \\ \gamma_2 & \gamma_3 & \gamma_4 & \gamma_5 \\ \vdots & \vdots & \vdots & \vdots \\ \gamma_{m+l+1} & \gamma_{m+l+2} & \gamma_{m+l+3} & \gamma_{m+l+4} \\ \vdots & \vdots & \vdots & \vdots \\ \gamma_{s+m+l} & \gamma_{s+m+l+1} & \gamma_{s+m+l+2} & \gamma_{s+m+l+3} \end{pmatrix}.$$

We have respectively by (6.1), (6.2) with $k = 4$

$$(6.15) \quad \sum_{i=0}^4 \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times 4} = 2^{3s+2m+l+9},$$

$$(6.16) \quad \sum_{i=0}^4 \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times 4} \cdot 2^{-i} = 2^{3s+2m+l+5} + 2^9 - 2^5.$$

From (6.15) and (6.16) using (6.14), (6.7) (6.12) we deduce

$$(6.17) \quad \Gamma_3 \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] \times k = \Gamma_3 \left[\begin{smallmatrix} 3 \\ 3 \end{smallmatrix} \right] \times 4 = 6384 \quad \text{for all } s \geq 4, k \geq 4, m \geq 0, l \geq 0$$

$$(6.18) \quad \Gamma_4 \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] \times 4 = 2^{3s+2m+l+9} - 6784 \quad \text{for all } s \geq 4, k = 4, m \geq 0, l \geq 0$$

Computation of $\omega_3(s - (i - 3), m + (i - 3), l, k)$ in the formula (5.1) for $4 \leq i \leq \inf(s - 1, k)$

By Definition 5.1

$$\omega_3(s - (i - 3), m + (i - 3), l, k) = \Gamma_3 \left[\begin{smallmatrix} s - (i - 3) \\ s + m \end{smallmatrix} \right] \times k - 4 \cdot \Gamma_2 \left[\begin{smallmatrix} s - (i - 3) \\ s + m - 1 \end{smallmatrix} \right] \times k - 8 \cdot \Gamma_2 \left[\begin{smallmatrix} s - (i - 3) \\ s + m + l - 1 \end{smallmatrix} \right] \times k + 32 \cdot \Gamma_1 \left[\begin{smallmatrix} s - (i - 3) \\ s + m + l - 1 \end{smallmatrix} \right] \times k$$

from (6.7), (6.12) and (6.17) we obtain

$$(6.19) \quad \omega_3(s - (i - 3), m + (i - 3), l, k) = 6384 - 4 \cdot 378 - 8 \cdot 378 + 32 \cdot 21 = 2520 \quad \text{for } 4 \leq i \leq \inf(s - 1, k)$$

Computation of $\omega_i(s, m, l, k + 1) - \omega_i(s, m, l, k)$ for $4 \leq i \leq \inf(s - 1, k - 1)$

By (5.1) and (6.19) we get for $4 \leq i \leq \inf(s - 1, k)$

$$\begin{aligned} \omega_i(s, m, l, k) &= 2^{i-3} \cdot \omega_3(s - (i - 3), m + (i - 3), l, k) + \sum_{j=0}^{i-4} 2^j \Delta_{i-j} \left[\begin{smallmatrix} s-j \\ s+m \end{smallmatrix} \right] \times k \\ &= 315 \cdot 2^i + \sum_{j=0}^{i-4} 2^j \Delta_{i-j} \left[\begin{smallmatrix} s-j \\ s+m+l \end{smallmatrix} \right] \times k \end{aligned}$$

From (4.11) we then deduce

$$\omega_i(s, m, l, k + 1) - \omega_i(s, m, l, k) = 0 \quad \text{for } 4 \leq i \leq \inf(s - 1, k - 1),$$

that is

$$\begin{aligned} (6.20) \quad & \Gamma_i \left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right] \times k+1 - \Gamma_i \left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right] \times k \\ &= 4 \cdot \left(\Gamma_{i-1} \left[\begin{smallmatrix} s+m-1 \\ s+m+l \end{smallmatrix} \right] \times k+1 - \Gamma_{i-1} \left[\begin{smallmatrix} s+m-1 \\ s+m+l \end{smallmatrix} \right] \times k \right) + 8 \cdot \left(\Gamma_{i-1} \left[\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix} \right] \times k+1 - \Gamma_{i-1} \left[\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix} \right] \times k \right) \\ &\quad - 32 \cdot \left(\Gamma_{i-2} \left[\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times k+1 - \Gamma_{i-2} \left[\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times k \right) \quad \text{for } 4 \leq i \leq \inf(s - 1, k - 1) \end{aligned}$$

$(H_1) \wedge (H_2) \wedge \dots \wedge (H_j)$ implies (H_{j+1})

From (6.20) with $i \rightarrow j+1$ we get

$$\begin{aligned} (6.21) \quad & \Gamma_{j+1} \left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right] \times k+1 - \Gamma_{j+1} \left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right] \times k \\ &= 4 \cdot \left(\Gamma_j \left[\begin{smallmatrix} s+m-1 \\ s+m+l \end{smallmatrix} \right] \times k+1 - \Gamma_j \left[\begin{smallmatrix} s+m-1 \\ s+m+l \end{smallmatrix} \right] \times k \right) + 8 \cdot \left(\Gamma_j \left[\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix} \right] \times k+1 - \Gamma_j \left[\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix} \right] \times k \right) \\ &\quad - 32 \cdot \left(\Gamma_{j-1} \left[\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times k+1 - \Gamma_{j-1} \left[\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times k \right) \quad \text{for } s \geq j+2, k \geq j+2, m \geq 0, l \geq 0 \end{aligned}$$

By (H_j) , (H_{j-1}) we obtain respectively

(6.22)

$$\Gamma_j \begin{bmatrix} s+m-1 \\ s+m+l \end{bmatrix} \times k = \Gamma_j \begin{bmatrix} s+m \\ s+m+l-1 \end{bmatrix} \times k = \Gamma_j \begin{bmatrix} j \\ j \end{bmatrix} \times (j+1) = \gamma_j \quad \text{for } s \geq j+2, k \geq j+2, m \geq 0, l \geq 0$$

(6.23)

$$\Gamma_{j-1} \begin{bmatrix} s+m-1 \\ s+m+l-1 \end{bmatrix} \times k = \Gamma_{j-1} \begin{bmatrix} j-1 \\ j-1 \end{bmatrix} \times j = \gamma_{j-1} \quad \text{for } s \geq j+1, k \geq j+1, m \geq 0, l \geq 0$$

we deduce from (6.21), (6.22) and (6.23)

(6.24)

$$\Gamma_{j+1} \begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k = \Gamma_{j+1} \begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times (j+2) \quad \text{for } k \geq j+2, s \geq j+2, m \geq 0, l \geq 0$$

Consider now the matrix

$$D \begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times (j+2) (t, \eta, \xi) = \begin{pmatrix} \alpha_1 & \alpha_2 & \dots & \alpha_{j+1} & \alpha_{j+2} \\ \alpha_2 & \alpha_3 & \dots & \alpha_{j+2} & \alpha_{j+3} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \dots & \alpha_{s+j-1} & \alpha_{s+j} \\ \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+j} & \alpha_{s+j+1} \\ \beta_1 & \beta_2 & \dots & \beta_{j+1} & \beta_{j+2} \\ \beta_2 & \beta_3 & \dots & \beta_{j+2} & \beta_{j+3} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \dots & \beta_{j+m+1} & \beta_{j+m+2} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \dots & \beta_{s+m+j-1} & \beta_{s+m+j} \\ \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+j} & \beta_{s+m+j+1} \\ \gamma_1 & \gamma_2 & \dots & \gamma_{j+1} & \gamma_{j+2} \\ \gamma_2 & \gamma_3 & \dots & \gamma_{j+2} & \gamma_{j+3} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{m+l+1} & \gamma_{m+l+2} & \dots & \gamma_{j+m+l+1} & \gamma_{j+m+l+2} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{s+m+l} & \gamma_{s+m+l+1} & \dots & \gamma_{s+m+l+j} & \gamma_{s+m+l+j+1} \end{pmatrix}.$$

We have respectively by (6.1), (6.2) with $k = j+2$

$$(6.25) \quad \sum_{i=0}^{j+2} \Gamma_i \begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times (j+2) = 2^{3j+3s+2m+l+3},$$

$$(6.26) \quad \sum_{i=0}^{j+2} \Gamma_i \begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times (j+2) \cdot 2^{-i} = 2^{2j+3s+2m+l+1} + 2^{3j+3} - 2^{2j+1}.$$

Setting $s = j+1$, $m = 0$, $l = 0$ in (6.25), (6.26) we have

$$(6.27) \quad \sum_{i=0}^{j+2} \Gamma_i \begin{bmatrix} j+1 \\ j+1 \end{bmatrix} \times (j+2) = 2^{6j+6},$$

$$(6.28) \quad \sum_{i=0}^{j+2} \Gamma_i^{\left[\begin{smallmatrix} j+1 \\ j+1 \\ j+1 \end{smallmatrix} \right] \times (j+2)} \cdot 2^{-i} = 2^{5j+4} + 2^{3j+3} - 2^{2j+1}.$$

From $(H_1) \wedge (H_2) \wedge \dots \wedge (H_j)$ it follows that

$$(6.29) \quad \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times (j+2)} = \Gamma_i^{\left[\begin{smallmatrix} j+1 \\ j+1 \\ j+1 \end{smallmatrix} \right] \times (j+2)} = \Gamma_i^{\left[\begin{smallmatrix} i \\ i \\ i \end{smallmatrix} \right] \times (i+1)} \quad \text{for } 1 \leq i \leq j$$

We then deduce from (6.25), (6.26) and (6.29)

$$(6.30) \quad \Gamma_{j+1}^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times (j+2)} = 2^{4j+5} - 2^{3j+3} + \sum_{i=0}^j \gamma_i (1 - 2^{j+2-i})$$

and from (6.27), (6.28) and (6.29)

$$(6.31) \quad \Gamma_{j+1}^{\left[\begin{smallmatrix} j+1 \\ j+1 \\ j+1 \end{smallmatrix} \right] \times (j+2)} = 2^{4j+5} - 2^{3j+3} + \sum_{i=0}^j \gamma_i (1 - 2^{j+2-i})$$

By (6.24), (6.30) and (6.31) it follows that

$$(6.32) \quad (H_{j+1}) \quad \Gamma_{j+1}^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k} = \Gamma_{j+1}^{\left[\begin{smallmatrix} j+1 \\ j+1 \\ j+1 \end{smallmatrix} \right] \times (j+2)} = \gamma_{j+1} \quad \text{for all } s \geq j+2, k \geq j+2, m \geq 0, l \geq 0$$

□

Lemma 6.3. *We have*

$$(6.33) \quad \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \\ s+m+l \end{smallmatrix} \right] \times k} = \Gamma_i^{\left[\begin{smallmatrix} i \\ i \\ i \end{smallmatrix} \right] \times (i+1)} = \begin{cases} 1 & \text{if } i = 0, \\ 105 \cdot 2^{4i-6} - 21 \cdot 2^{3i-5} & \text{if } 1 \leq i \leq s-1, k \geq i+1, m \geq 0, l \geq 0. \end{cases}$$

Proof. We have respectively by (6.1), (6.2) with $k = i+1$, $s = i$ and $m=0, l=0$ using (6.3)

$$(6.34) \quad \sum_{j=0}^{i+1} \Gamma_j^{\left[\begin{smallmatrix} i \\ i \\ i \end{smallmatrix} \right] \times (i+1)} = \sum_{j=0}^i \gamma_j + \Gamma_{i+1}^{\left[\begin{smallmatrix} i \\ i \\ i \end{smallmatrix} \right] \times (i+1)} = 2^{6i}$$

and

$$(6.35) \quad \sum_{j=0}^{i+1} \Gamma_j^{\left[\begin{smallmatrix} i \\ i \\ i \end{smallmatrix} \right] \times (i+1)} \cdot 2^{-j} = \sum_{j=0}^i \gamma_j \cdot 2^{-j} + \Gamma_{i+1}^{\left[\begin{smallmatrix} i \\ i \\ i \end{smallmatrix} \right] \times (i+1)} \cdot 2^{-(i+1)} = 2^{5i-1} + 2^{3i} - 2^{2i-1}.$$

Subtracting (6.34) from (6.35) multiplied with 2^{i+1} we deduce

$$(6.36) \quad \gamma_i = 2^{4i+1} - 2^{3i} + \sum_{j=0}^{i-1} \gamma_j - \sum_{j=0}^{i-1} \gamma_j \cdot 2^{i+1-j}$$

Substituting i for $i-1$ in (6.36) we get

$$(6.37) \quad \gamma_{i-1} = 2^{4i-3} - 2^{3i-3} + \sum_{j=0}^{i-2} \gamma_j - \sum_{j=0}^{i-2} \gamma_j \cdot 2^{i-j}$$

From (6.36), (6.37) we obtain

$$\gamma_i - 2 \cdot \gamma_{i-1} = 2^{4i+1} - 2^{3i} + \sum_{j=0}^{i-1} \gamma_j - \sum_{j=0}^{i-1} \gamma_j \cdot 2^{i+1-j} - 2 \cdot (2^{4i-3} - 2^{3i-3} + \sum_{j=0}^{i-2} \gamma_j - \sum_{j=0}^{i-2} \gamma_j \cdot 2^{i-j})$$

$$\iff$$

$$\gamma_i - 2 \cdot \gamma_{i-1} = 7 \cdot 2^{4i-2} - 3 \cdot 2^{3i-2} - \sum_{j=0}^{i-2} \gamma_j - 3 \cdot \gamma_{i-1}$$

$$\iff$$

(6.38)

$$\sum_{j=0}^i \gamma_j = 7 \cdot 2^{4i-2} - 3 \cdot 2^{3i-2}$$

Finally from (6.38) we deduce

$$\gamma_i = \sum_{j=0}^i \gamma_j - \sum_{j=0}^{i-1} \gamma_j = 7 \cdot 2^{4i-2} - 3 \cdot 2^{3i-2} - (7 \cdot 2^{4i-6} - 3 \cdot 2^{3i-5}) = 105 \cdot 2^{4i-6} - 21 \cdot 2^{3i-5}$$

□

Lemma 6.4. *We have*

(6.39)

$$\sum_{j=0}^{s-1} \gamma_j = 7 \cdot 2^{4s-6} - 3 \cdot 2^{3s-5}$$

(6.40)

$$\sum_{j=0}^{s-1} \gamma_j \cdot 2^{-j} = 15 \cdot 2^{3s-6} - 7 \cdot 2^{2s-5}$$

(6.41)

$$\Gamma_i \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right]^{\times i} = 2^{3s+2m+l+3i-3} - 7 \cdot 2^{4i-6} + 3 \cdot 2^{3i-5} \quad \text{for } 1 \leq i \leq s+1, m \geq 0, l \geq 0$$

Proof.

Proof of (6.39)

From (6.33) we obtain

$$\sum_{j=0}^{s-1} \gamma_j = \sum_{j=0}^{s-1} (105 \cdot 2^{4j-6} - 21 \cdot 2^{3j-5}) = 7 \cdot 2^{4s-6} - 3 \cdot 2^{3s-5}$$

Proof of (6.40)

$$\sum_{j=0}^{s-1} \gamma_j \cdot 2^{-j} = \sum_{j=0}^{s-1} (105 \cdot 2^{4j-6} - 21 \cdot 2^{3j-5}) \cdot 2^{-j} = \sum_{j=0}^{s-1} (105 \cdot 2^{3j-6} - 21 \cdot 2^{2j-5}) = 15 \cdot 2^{3s-6} - 7 \cdot 2^{2s-5}$$

Proof of (6.41) in the case $1 \leq i \leq s$.From (6.1) with $k = i$ using (6.33) we obtain

$$\Gamma_i \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right]^{\times i} = 2^{3s+2m+l+3i-3} - \sum_{j=0}^{i-1} \gamma_j = 2^{3s+2m+l+3i-3} - \sum_{j=0}^{i-1} (105 \cdot 2^{4j-6} - 21 \cdot 2^{3j-5})$$

$$= 2^{3s+2m+l+3i-3} - 7 \cdot 2^{4i-6} + 3 \cdot 2^{3i-5}$$

Proof of (6.41) in the case $i = s+1$

From (6.1) with $k = s+1$ using (6.39) we obtain

$$\Gamma_s \left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right]^{\times(s+1)} + \Gamma_{s+1} \left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right]^{\times(s+1)} = 2^{6s+2m+l} - \sum_{j=0}^{s-1} \gamma_j = 2^{6s+2m+l} - 7 \cdot 2^{4s-6} + 3 \cdot 2^{3s-5}$$

From (6.2) with $k = s+1$ using (6.40) we obtain

$$\begin{aligned} 2 \cdot \Gamma_s \left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right]^{\times(s+1)} + \Gamma_{s+1} \left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right]^{\times(s+1)} &= 2^{6s+2m+l} + 2^{4s+1} - 2^{3s} - 2^{s+1} \cdot \sum_{j=0}^{s-1} \gamma_j \cdot 2^{-j} \\ &= 2^{6s+2m+l} + 2^{4s+1} - 2^{3s} - 2^{s+1} \cdot (15 \cdot 2^{3s-6} - 7 \cdot 2^{2s-5}) \\ &= 2^{6s+2m+l} + 2^{4s+1} - 2^{3s} - 15 \cdot 2^{4s-5} + 7 \cdot 2^{3s-4} = 2^{6s+2m+l} + 49 \cdot 2^{4s-5} - 9 \cdot 2^{3s-4} \end{aligned}$$

We then deduce from the above equations

(6.42)

$$\Gamma_s \left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right]^{\times(s+1)} = 105 \cdot 2^{4s-6} - 21 \cdot 2^{3s-5}, \quad \text{remark that (6.33) holds for } i = s, \quad k = s+1$$

(6.43)

$$\Gamma_{s+1} \left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right]^{\times(s+1)} = 2^{6s+2m+l} - 7 \cdot 2^{4s-2} + 3 \cdot 2^{3s-2}$$

□

7. Computation of $\omega_i(s, m, l, k+1) - \omega_i(s, m, l, k)$ in the case $s \leq i \leq \inf(3s+2m+l, k)$, $k > i$

7.1. Notation.

Definition 7.1.

We define $A_i^{[s, m, l, k]}$ to be equal to $\Gamma_{i-s} \left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right]^{\times k} - 4 \cdot \Gamma_{i-s-1} \left[\begin{smallmatrix} s+m-1 \\ s+m+l \end{smallmatrix} \right]^{\times k} - 8 \cdot \Gamma_{i-s-1} \left[\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix} \right]^{\times k} + 32 \cdot \Gamma_{i-s-2} \left[\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix} \right]^{\times k}$ for $s \leq i \leq \inf(k, 3s+2m+l)$

We recall that (see 5.1)

$$\begin{aligned} \omega_i(s, m, l, k+1) - \omega_i(s, m, l, k) &= \left(\Gamma_i \left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right]^{\times(k+1)} - \Gamma_i \left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right]^{\times k} \right) - 4 \cdot \left(\Gamma_{i-1} \left[\begin{smallmatrix} s+m-1 \\ s+m+l \end{smallmatrix} \right]^{\times(k+1)} - \Gamma_{i-1} \left[\begin{smallmatrix} s+m-1 \\ s+m+l \end{smallmatrix} \right]^{\times k} \right) \\ &\quad - 8 \cdot \left(\Gamma_{i-1} \left[\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix} \right]^{\times(k+1)} - \Gamma_{i-1} \left[\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix} \right]^{\times k} \right) + 32 \cdot \left(\Gamma_{i-2} \left[\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix} \right]^{\times(k+1)} - \Gamma_{i-2} \left[\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix} \right]^{\times k} \right) \end{aligned}$$

7.2. Introduction. We adapt the method used in Section 10 of [2] to compute explicitly

$\omega_i(s, m, l, k+1) - \omega_i(s, m, l, k)$ for $s \leq i \leq \inf(k, 3s+2m+l)$, $k > i$

7.3. An equivalent to the formula for $\omega_i(s, m, l, k)$ in the case $s \leq i \leq \inf(3s+2m+l, k)$.

Lemma 7.2. *Let $s \geq 2, m \geq 0, l \geq 0$ and $k \geq 1$, then we have*

(7.1)

$$\omega_i(s, m, l, k) = 2^{k+s-1} A_i^{[s, m, l, k]} + 2^i \Delta_{i-s+1} \left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right]^{\times k} - 2^{i-1} \Delta_{i-s} \left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right]^{\times k} + \sum_{j=0}^{s-2} 2^j \Delta_{i-j} \left[\begin{smallmatrix} s-j \\ s+m+l \end{smallmatrix} \right]^{\times k} \quad \text{if } s \leq i \leq \inf(3s+2m+l, k)$$

(7.2)

$$\omega_i(s, m, l, k+1) - \omega_i(s, m, l, k) = 2^{k+s} A_i^{[s, m, l, k+1]} - 2^{k+s-1} A_i^{[s, m, l, k]} \quad \text{if } s \leq i \leq \inf(3s+2m+l, k), \quad k > i$$

where

$$A_i^{[s, m, l, k]} = \Gamma_{i-s}^{\left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_{i-s-1}^{\left[\begin{smallmatrix} s+m-1 \\ s+m+l \end{smallmatrix} \right] \times k} - 8 \cdot \Gamma_{i-s-1}^{\left[\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_{i-s-2}^{\left[\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times k}$$

Proof.

proof of (7.1)

From Definition 5.1 and Theorem 1.4 we obtain

$$\begin{aligned} \omega_{i-(s-1)}(1, m+s-1, l, k) &= \Gamma_{i-(s-1)}^{\left[\begin{smallmatrix} 1 \\ s+m+l \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_{i-s}^{\left[\begin{smallmatrix} 1 \\ s+m+l \end{smallmatrix} \right] \times k} - 8 \cdot \Gamma_{i-s}^{\left[\begin{smallmatrix} 1 \\ s+m+l-1 \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_{i-s-1}^{\left[\begin{smallmatrix} 1 \\ s+m+l-1 \end{smallmatrix} \right] \times k} \\ &= (2^k - 2^{i-s}) \Gamma_{i-s}^{\left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right] \times k} + 2^{i-s+1} \Gamma_{i-s+1}^{\left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right] \times k} \\ &\quad - 4 \cdot [(2^k - 2^{i-s-1}) \Gamma_{i-s-1}^{\left[\begin{smallmatrix} s+m-1 \\ s+m+l \end{smallmatrix} \right] \times k} + 2^{i-s} \Gamma_{i-s}^{\left[\begin{smallmatrix} s+m-1 \\ s+m+l \end{smallmatrix} \right] \times k}] \\ &\quad - 8 \cdot [(2^k - 2^{i-s-1}) \Gamma_{i-s-1}^{\left[\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix} \right] \times k} + 2^{i-s} \Gamma_{i-s}^{\left[\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix} \right] \times k}] \\ &\quad + 32 \cdot [(2^k - 2^{i-s-2}) \Gamma_{i-s-2}^{\left[\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times k} + 2^{i-s-1} \Gamma_{i-s-1}^{\left[\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times k}] \\ &= 2^k \cdot [\Gamma_{i-s}^{\left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_{i-s-1}^{\left[\begin{smallmatrix} s+m-1 \\ s+m+l \end{smallmatrix} \right] \times k} - 8 \cdot \Gamma_{i-s-1}^{\left[\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_{i-s-2}^{\left[\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times k}] \\ &\quad + 2^{i-s+1} \cdot [\Gamma_{i-s+1}^{\left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right] \times k} - 2 \cdot \Gamma_{i-s}^{\left[\begin{smallmatrix} s+m-1 \\ s+m+l \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_{i-s}^{\left[\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix} \right] \times k} + 8 \cdot \Gamma_{i-s-1}^{\left[\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times k}] \\ &\quad - 2^{i-s} \cdot [\Gamma_{i-s}^{\left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right] \times k} - 2 \cdot \Gamma_{i-s-1}^{\left[\begin{smallmatrix} s+m-1 \\ s+m+l \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_{i-s-1}^{\left[\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix} \right] \times k} + 8 \cdot \Gamma_{i-s-2}^{\left[\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] \times k}] \\ &= 2^k A_i^{[s, m, l, k]} + 2^{i-s+1} \Delta_{i-s+1}^{\left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right] \times k} - 2^{i-s} \Delta_{i-s}^{\left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right] \times k} \end{aligned}$$

Hence from (5.1) we get easily (7.1)

proof of (7.2)

From Lemma 4.5 we obtain

$$\sum_{j=0}^{s-2} 2^j \Delta_{i-j}^{\left[\begin{smallmatrix} s-j \\ s+m+l \end{smallmatrix} \right] \times (k+1)} - \sum_{j=0}^{s-2} 2^j \Delta_{i-j}^{\left[\begin{smallmatrix} s-j \\ s+m+l \end{smallmatrix} \right] \times k} = 0 \quad \text{for } k > i,$$

and from Lemma 8.5 (see [2] section 8) we get

$$\Delta_{i-s+1}^{\left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right] \times (k+1)} - \Delta_{i-s+1}^{\left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right] \times k} = 0, \quad \Delta_{i-s}^{\left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right] \times (k+1)} - \Delta_{i-s}^{\left[\begin{smallmatrix} s+m \\ s+m+l \end{smallmatrix} \right] \times k} = 0 \quad \text{for } k > i$$

Hence from the above equations we obtain easily (7.2)

□

7.4. Computation of $A_i^{[s, m, l, k]}$.

Lemma 7.3. Set for $s \leq i \leq \inf(k, 3s+2m+l)$

$$A_i^{[s, m, l, k]} = \Gamma_{i-s}^{\left[\begin{smallmatrix} m+s \\ m+s+l \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_{i-s-1}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l \end{smallmatrix} \right] \times k} - 8 \cdot \Gamma_{i-s-1}^{\left[\begin{smallmatrix} m+s \\ m+s+l-1 \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_{i-s-2}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l-1 \end{smallmatrix} \right] \times k}$$

We have in the case $l=0, m=0, s=2$

$$A_i^{[2, 0, 0, k]} = \Gamma_{i-2}^{\left[\begin{smallmatrix} 2 \\ 2 \end{smallmatrix} \right] \times k} - 12 \cdot \Gamma_{i-3}^{\left[\begin{smallmatrix} 1 \\ 2 \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_{i-4}^{\left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right] \times k}$$

$$(7.3) \quad A_i^{[2,0,0,k]} = \begin{cases} 1 & \text{if } i = 2 \\ -3 & \text{if } i = 3 \\ -3 \cdot 2^{k+1} + 2 & \text{if } i = 4 \\ 3 \cdot 2^{k+1} & \text{if } i = 5 \\ 3 \cdot 2^{2k+2} & \text{if } i = 6 \end{cases}$$

We have in the case $l = 0, m = 1, s = 2$

$$(7.4) \quad A_i^{[2,1,0,k]} = \Gamma_{i-2}^{\begin{bmatrix} 3 \\ 3 \end{bmatrix} \times k} - 12 \cdot \Gamma_{i-3}^{\begin{bmatrix} 2 \\ 3 \end{bmatrix} \times k} + 32 \cdot \Gamma_{i-4}^{\begin{bmatrix} 2 \\ 2 \end{bmatrix} \times k}$$

$$A_i^{[2,1,0,k]} = \begin{cases} 1 & \text{if } i = 2 \\ -3 & \text{if } i = 3 \\ 2 & \text{if } i = 4 \\ -3 \cdot 2^{k+2} & \text{if } i = 5 \\ 3 \cdot 2^{k+2} & \text{if } i = 6 \\ 0 & \text{if } i = 7 \\ 3 \cdot 2^{2k+4} & \text{if } i = 8 \end{cases}$$

We have in the case $l = 0, m \geq 2, s = 2$

$$(7.5) \quad A_i^{[2,m,0,k]} = \Gamma_{i-2}^{\begin{bmatrix} m+2 \\ m+2 \end{bmatrix} \times k} - 12 \cdot \Gamma_{i-3}^{\begin{bmatrix} m+1 \\ m+2 \end{bmatrix} \times k} + 32 \cdot \Gamma_{i-4}^{\begin{bmatrix} m+1 \\ m+1 \end{bmatrix} \times k}$$

$$A_i^{[2,m,0,k]} = \begin{cases} 1 & \text{if } i = 2 \\ -3 & \text{if } i = 3 \\ 2 & \text{if } i = 4 \\ 0 & \text{if } 5 \leq i \leq m+3 \\ -3 \cdot 2^{k+m+1} & \text{if } i = m+4 \\ 3 \cdot 2^{k+m+1} & \text{if } i = m+5 \\ 0 & \text{if } m+7 \leq i \leq 2m+5 \\ 3 \cdot 2^{2k+2m+2} & \text{if } i = 2m+6 \end{cases}$$

We have in the case $l = 1, m = 0, s = 2$

$$(7.6) \quad A_i^{[2,0,1,k]} = \Gamma_{i-2}^{\begin{bmatrix} 2 \\ 3 \end{bmatrix} \times k} - 4 \cdot \Gamma_{i-3}^{\begin{bmatrix} 1 \\ 3 \end{bmatrix} \times k} - 8 \cdot \Gamma_{i-3}^{\begin{bmatrix} 2 \\ 2 \end{bmatrix} \times k} + 32 \cdot \Gamma_{i-4}^{\begin{bmatrix} 2 \\ 2 \end{bmatrix} \times k}$$

$$A_i^{[2,0,1,k]} = \begin{cases} 1 & \text{if } i = 2 \\ -3 & \text{if } i = 3 \\ -2^{k+1} + 2 & \text{if } i = 4 \\ -3 \cdot 2^{k+1} & \text{if } i = 5 \\ 2^{k+3} & \text{if } i = 6 \\ 3 \cdot 2^{2k+3} & \text{if } i = 7 \end{cases}$$

We have in the case $l = 1, m = 1, s = 2$

$$A_i^{[2,1,1,k]} = \Gamma_{i-2}^{\begin{bmatrix} 3 \\ 4 \end{bmatrix} \times k} - 4 \cdot \Gamma_{i-3}^{\begin{bmatrix} 2 \\ 4 \end{bmatrix} \times k} - 8 \cdot \Gamma_{i-3}^{\begin{bmatrix} 3 \\ 3 \end{bmatrix} \times k} + 32 \cdot \Gamma_{i-4}^{\begin{bmatrix} 2 \\ 3 \end{bmatrix} \times k}$$

$$(7.7) \quad A_i^{[2,1,1,k]} = \begin{cases} 1 & \text{if } i = 2 \\ -3 & \text{if } i = 3 \\ 2 & \text{if } i = 4 \\ -2^{k+2} & \text{if } i = 5 \\ -3 \cdot 2^{k+2} & \text{if } i = 6 \\ 2^{k+4} & \text{if } i = 7 \\ 0 & \text{if } i = 8 \\ 3 \cdot 2^{2k+5} & \text{if } i = 9 \end{cases}$$

We have in the case $\underline{l = 1, m \geq 2, s = 2}$

$$(7.8) \quad A_i^{[2,m,1,k]} = \Gamma_{i-2}^{\begin{bmatrix} m+2 \\ m+3 \end{bmatrix} \times k} - 4 \cdot \Gamma_{i-3}^{\begin{bmatrix} m+1 \\ m+3 \end{bmatrix} \times k} - 8 \cdot \Gamma_{i-3}^{\begin{bmatrix} m+2 \\ m+2 \end{bmatrix} \times k} + 32 \cdot \Gamma_{i-4}^{\begin{bmatrix} m+1 \\ m+2 \end{bmatrix} \times k}$$

$$A_i^{[2,m,1,k]} = \begin{cases} 1 & \text{if } i = 2 \\ -3 & \text{if } i = 3 \\ 2 & \text{if } i = 4 \\ 0 & \text{if } 5 \leq i \leq m+3 \\ -2^{k+m+1} & \text{if } i = m+4 \\ -3 \cdot 2^{k+m+1} & \text{if } i = m+5 \\ 2^{k+m+3} & \text{if } i = m+6 \\ 0 & \text{if } m+7 \leq i \leq 2m+6 \\ 3 \cdot 2^{2k+2m+3} & \text{if } i = 2m+7 \end{cases}$$

We have in the case $\underline{l = 2, m = 0, s = 2}$

$$(7.9) \quad A_i^{[2,0,2,k]} = \Gamma_{i-2}^{\begin{bmatrix} 2 \\ 4 \end{bmatrix} \times k} - 4 \cdot \Gamma_{i-3}^{\begin{bmatrix} 1 \\ 4 \end{bmatrix} \times k} - 8 \cdot \Gamma_{i-3}^{\begin{bmatrix} 2 \\ 3 \end{bmatrix} \times k} + 32 \cdot \Gamma_{i-4}^{\begin{bmatrix} 1 \\ 3 \end{bmatrix} \times k}$$

$$A_i^{[2,0,2,k]} = \begin{cases} 1 & \text{if } i = 2 \\ -3 & \text{if } i = 3 \\ -2^{k+1} + 2 & \text{if } i = 4 \\ 5 \cdot 2^{k+1} & \text{if } i = 5 \\ -5 \cdot 2^{k+3} & \text{if } i = 6 \\ 2^{k+5} & \text{if } i = 7 \\ 3 \cdot 2^{2k+4} & \text{if } i = 8 \end{cases}$$

We have in the case $\underline{l = 2, m = 1, s = 2}$

$$(7.10) \quad A_i^{[2,1,2,k]} = \Gamma_{i-2}^{\begin{bmatrix} 3 \\ 5 \end{bmatrix} \times k} - 4 \cdot \Gamma_{i-3}^{\begin{bmatrix} 2 \\ 5 \end{bmatrix} \times k} - 8 \cdot \Gamma_{i-3}^{\begin{bmatrix} 3 \\ 4 \end{bmatrix} \times k} + 32 \cdot \Gamma_{i-4}^{\begin{bmatrix} 2 \\ 4 \end{bmatrix} \times k}$$

$$A_i^{[2,1,2,k]} = \begin{cases} 1 & \text{if } i = 2 \\ -3 & \text{if } i = 3 \\ 2 & \text{if } i = 4 \\ -2^{k+2} & \text{if } i = 5 \\ 5 \cdot 2^{k+2} & \text{if } i = 6 \\ -5 \cdot 2^{k+4} & \text{if } i = 7 \\ 2^{k+6} & \text{if } i = 8 \\ 0 & \text{if } i = 9 \\ 3 \cdot 2^{2k+6} & \text{if } i = 10 \end{cases}$$

We have in the case $\underline{l = 2, m \geq 2, s = 2}$

$$A_i^{[2,m,2,k]} = \Gamma_{i-2}^{\left[\begin{smallmatrix} m+2 \\ m+4 \end{smallmatrix}\right] \times k} - 4 \cdot \Gamma_{i-3}^{\left[\begin{smallmatrix} m+1 \\ m+4 \end{smallmatrix}\right] \times k} - 8 \cdot \Gamma_{i-3}^{\left[\begin{smallmatrix} m+2 \\ m+3 \end{smallmatrix}\right] \times k} + 32 \cdot \Gamma_{i-4}^{\left[\begin{smallmatrix} m+1 \\ m+3 \end{smallmatrix}\right] \times k}$$

$$(7.11) \quad A_i^{[2,m,2,k]} = \begin{cases} 1 & \text{if } i = 2 \\ -3 & \text{if } i = 3 \\ 2 & \text{if } i = 4 \\ 0 & \text{if } 5 \leq i \leq m+3 \\ -2^{k+m+1} & \text{if } i = m+4 \\ 5 \cdot 2^{k+m+1} & \text{if } i = m+5 \\ -5 \cdot 2^{k+m+3} & \text{if } i = m+6 \\ 2^{k+m+5} & \text{if } i = m+7 \\ 0 & \text{if } m+8 \leq i \leq 2m+7 \\ 3 \cdot 2^{2k+2m+4} & \text{if } i = 2m+8 \end{cases}$$

We have in the case $\underline{l \geq 3, m = 0, s = 2}$

$$A_i^{[2,0,l,k]} = \Gamma_{i-2}^{\left[\begin{smallmatrix} 2 \\ 2+l \end{smallmatrix}\right] \times k} - 4 \cdot \Gamma_{i-3}^{\left[\begin{smallmatrix} 1 \\ 2+l \end{smallmatrix}\right] \times k} - 8 \cdot \Gamma_{i-3}^{\left[\begin{smallmatrix} 2 \\ 1+l \end{smallmatrix}\right] \times k} + 32 \cdot \Gamma_{i-4}^{\left[\begin{smallmatrix} 1 \\ 1+l \end{smallmatrix}\right] \times k}$$

$$(7.12) \quad A_i^{[2,0,l,k]} = \begin{cases} 1 & \text{if } i = 2 \\ -3 & \text{if } i = 3 \\ -2^{k+1} + 2 & \text{if } i = 4 \\ 5 \cdot 2^{k+1} & \text{if } i = 5 \\ 3 \cdot 2^{k+2i-9} & \text{if } 6 \leq i \leq l+3 \\ -5 \cdot 2^{k+2l-1} & \text{if } i = l+4 \\ 2^{k+2l+1} & \text{if } i = l+5 \\ 3 \cdot 2^{2k+l+2} & \text{if } i = l+6 \end{cases}$$

We have in the case $\underline{l \geq 3, m = 1, s = 2}$

$$A_i^{[2,1,l,k]} = \Gamma_{i-2}^{\left[\begin{smallmatrix} 3 \\ 3+l \end{smallmatrix}\right] \times k} - 4 \cdot \Gamma_{i-3}^{\left[\begin{smallmatrix} 2 \\ 3+l \end{smallmatrix}\right] \times k} - 8 \cdot \Gamma_{i-3}^{\left[\begin{smallmatrix} 3 \\ 2+l \end{smallmatrix}\right] \times k} + 32 \cdot \Gamma_{i-4}^{\left[\begin{smallmatrix} 2 \\ 2+l \end{smallmatrix}\right] \times k}$$

$$(7.13) \quad A_i^{[2,1,l,k]} = \begin{cases} 1 & \text{if } i = 2 \\ -3 & \text{if } i = 3 \\ 2 & \text{if } i = 4 \\ -2^{k+2} & \text{if } i = 5 \\ 5 \cdot 2^{k+2} & \text{if } i = 6 \\ 3 \cdot 2^{k+2i-10} & \text{if } 7 \leq i \leq l+4 \\ -5 \cdot 2^{k+2l} & \text{if } i = l+5 \\ 2^{k+2l+2} & \text{if } i = l+6 \\ 0 & \text{if } i = l+7 \\ 3 \cdot 2^{2k+l+4} & \text{if } i = l+8 \end{cases}$$

We have in the case $\underline{l \geq 3, m \geq 2, s = 2}$

$$A_i^{[2,m,l,k]} = \Gamma_{i-2}^{\left[\begin{smallmatrix} m+2 \\ m+2+l \end{smallmatrix}\right] \times k} - 4 \cdot \Gamma_{i-3}^{\left[\begin{smallmatrix} m+1 \\ m+2+l \end{smallmatrix}\right] \times k} - 8 \cdot \Gamma_{i-3}^{\left[\begin{smallmatrix} m+2 \\ m+1+l \end{smallmatrix}\right] \times k} + 32 \cdot \Gamma_{i-4}^{\left[\begin{smallmatrix} m+1 \\ m+1+l \end{smallmatrix}\right] \times k}$$

$$(7.14) \quad A_i^{[2,m,l,k]} = \begin{cases} 1 & \text{if } i = 2 \\ -3 & \text{if } i = 3 \\ 2 & \text{if } i = 4 \\ 0 & \text{if } 5 \leq i \leq m+3 \\ -2^{k+m+1} & \text{if } i = m+4 \\ 5 \cdot 2^{k+m+1} & \text{if } i = m+5 \\ 3 \cdot 2^{k-m+2i-9} & \text{if } m+6 \leq i \leq m+l+3 \\ -5 \cdot 2^{k+m+2l-1} & \text{if } i = m+l+4 \\ 2^{k+m+2l+1} & \text{if } i = m+l+5 \\ 0 & \text{if } m+l+6 \leq i \leq 2m+l+5 \\ 3 \cdot 2^{2k+2m+l+2} & \text{if } i = 2m+l+6 \end{cases}$$

We have in the case $\underline{l=0, m=0, s=3}$

$$(7.15) \quad A_i^{[3,0,0,k]} = \Gamma_{i-3}^{\begin{bmatrix} 3 \\ 3 \end{bmatrix} \times k} - 12 \cdot \Gamma_{i-4}^{\begin{bmatrix} 2 \\ 3 \end{bmatrix} \times k} + 32 \cdot \Gamma_{i-5}^{\begin{bmatrix} 2 \\ 2 \end{bmatrix} \times k}$$

$$A_i^{[3,0,0,k]} = \begin{cases} 1 & \text{if } i = 3 \\ -3 & \text{if } i = 4 \\ 2 & \text{if } i = 5 \\ -3 \cdot 2^{k+2} & \text{if } i = 6 \\ 3 \cdot 2^{k+2} & \text{if } i = 7 \\ 0 & \text{if } i = 8 \\ 3 \cdot 2^{2k+4} & \text{if } i = 9 \end{cases}$$

We have in the case $\underline{l=0, m \geq 1, s=3}$

$$(7.16) \quad A_i^{[s,m,0,k]} = \Gamma_{i-3}^{\begin{bmatrix} m+3 \\ m+3 \end{bmatrix} \times k} - 12 \cdot \Gamma_{i-4}^{\begin{bmatrix} m+2 \\ m+3 \end{bmatrix} \times k} + 32 \cdot \Gamma_{i-5}^{\begin{bmatrix} m+2 \\ m+2 \end{bmatrix} \times k}$$

$$A_i^{[3,m,0,k]} = \begin{cases} 1 & \text{if } i = 3 \\ -3 & \text{if } i = 4 \\ 2 & \text{if } i = 5 \\ 0 & \text{if } 6 \leq i \leq m+5 \\ -3 \cdot 2^{k+m+2} & \text{if } i = m+6 \\ 3 \cdot 2^{k+m+2} & \text{if } i = m+7 \\ 0 & \text{if } m+8 \leq i \leq 2m+8 \\ 3 \cdot 2^{2k+2m+4} & \text{if } i = 2m+9 \end{cases}$$

We have in the case $\underline{l=1, m=0, s=3}$

$$(7.17) \quad A_i^{[3,0,1,k]} = \Gamma_{i-3}^{\begin{bmatrix} 3 \\ 4 \end{bmatrix} \times k} - 4 \cdot \Gamma_{i-4}^{\begin{bmatrix} 2 \\ 4 \end{bmatrix} \times k} - 8 \cdot \Gamma_{i-4}^{\begin{bmatrix} 3 \\ 3 \end{bmatrix} \times k} + 32 \cdot \Gamma_{i-5}^{\begin{bmatrix} 2 \\ 3 \end{bmatrix} \times k}$$

$$A_i^{[3,0,1,k]} = \begin{cases} 1 & \text{if } i = 3 \\ -3 & \text{if } i = 4 \\ 2 & \text{if } i = 5 \\ -2^{k+2} & \text{if } i = 6 \\ -3 \cdot 2^{k+2} & \text{if } i = 7 \\ 2^{k+4} & \text{if } i = 8 \\ 0 & \text{if } i = 9 \\ 3 \cdot 2^{2k+5} & \text{if } i = 10 \end{cases}$$

We have in the case $\underline{l = 1, m \geq 1, s = 3}$

$$(7.18) \quad A_i^{[3,m,1,k]} = \Gamma_{i-3}^{\left[\begin{smallmatrix} m+3 \\ m+4 \end{smallmatrix}\right] \times k} - 4 \cdot \Gamma_{i-4}^{\left[\begin{smallmatrix} m+2 \\ m+4 \end{smallmatrix}\right] \times k} - 8 \cdot \Gamma_{i-4}^{\left[\begin{smallmatrix} m+3 \\ m+3 \end{smallmatrix}\right] \times k} + 32 \cdot \Gamma_{i-5}^{\left[\begin{smallmatrix} m+2 \\ m+3 \end{smallmatrix}\right] \times k}$$

$$A_i^{[3,m,1,k]} = \begin{cases} 1 & \text{if } i = 3 \\ -3 & \text{if } i = 4 \\ 2 & \text{if } i = 5 \\ 0 & \text{if } 6 \leq i \leq m+5 \\ -2^{k+m+2} & \text{if } i = m+6 \\ -3 \cdot 2^{k+m+2} & \text{if } i = m+7 \\ 2^{k+m+4} & \text{if } i = m+8 \\ 0 & \text{if } m+9 \leq i \leq 2m+9 \\ 3 \cdot 2^{2k+2m+5} & \text{if } i = 2m+10 \end{cases}$$

We have in the case $\underline{l = 2, m = 0, s = 3}$

$$(7.19) \quad A_i^{[3,0,2,k]} = \Gamma_{i-3}^{\left[\begin{smallmatrix} 3 \\ 5 \end{smallmatrix}\right] \times k} - 4 \cdot \Gamma_{i-4}^{\left[\begin{smallmatrix} 2 \\ 5 \end{smallmatrix}\right] \times k} - 8 \cdot \Gamma_{i-4}^{\left[\begin{smallmatrix} 3 \\ 4 \end{smallmatrix}\right] \times k} + 32 \cdot \Gamma_{i-5}^{\left[\begin{smallmatrix} 2 \\ 4 \end{smallmatrix}\right] \times k}$$

$$A_i^{[3,0,2,k]} = \begin{cases} 1 & \text{if } i = 3 \\ -3 & \text{if } i = 4 \\ 2 & \text{if } i = 5 \\ -2^{k+2} & \text{if } i = 6 \\ 5 \cdot 2^{k+2} & \text{if } i = 7 \\ -5 \cdot 2^{k+4} & \text{if } i = 8 \\ 2^{k+6} & \text{if } i = 9 \\ 0 & \text{if } i = 10 \\ 3 \cdot 2^{2k+6} & \text{if } i = 11 \end{cases}$$

We have in the case $\underline{l = 2, m \geq 1, s = 3}$

$$(7.20) \quad A_i^{[3,m,2,k]} = \Gamma_{i-3}^{\left[\begin{smallmatrix} m+3 \\ m+5 \end{smallmatrix}\right] \times k} - 4 \cdot \Gamma_{i-4}^{\left[\begin{smallmatrix} m+2 \\ m+5 \end{smallmatrix}\right] \times k} - 8 \cdot \Gamma_{i-4}^{\left[\begin{smallmatrix} m+3 \\ m+4 \end{smallmatrix}\right] \times k} + 32 \cdot \Gamma_{i-5}^{\left[\begin{smallmatrix} m+2 \\ m+4 \end{smallmatrix}\right] \times k}$$

$$A_i^{[3,m,2,k]} = \begin{cases} 1 & \text{if } i = 3 \\ -3 & \text{if } i = 4 \\ 2 & \text{if } i = 5 \\ 0 & \text{if } 6 \leq i \leq m+5 \\ -2^{k+m+2} & \text{if } i = m+6 \\ 5 \cdot 2^{k+m+2} & \text{if } i = m+7 \\ -5 \cdot 2^{k+m+4} & \text{if } i = m+8 \\ 2^{k+m+6} & \text{if } i = m+9 \\ 0 & \text{if } m+10 \leq i \leq 2m+10 \\ 3 \cdot 2^{2k+2m+6} & \text{if } i = 2m+11 \end{cases}$$

We have in the case $\underline{l \geq 3, m = 0, s = 3}$

$$A_i^{[3,0,l,k]} = \Gamma_{i-3}^{\left[\begin{smallmatrix} 3 \\ 3+l \end{smallmatrix}\right] \times k} - 4 \cdot \Gamma_{i-4}^{\left[\begin{smallmatrix} 2 \\ 3+l \end{smallmatrix}\right] \times k} - 8 \cdot \Gamma_{i-4}^{\left[\begin{smallmatrix} 3 \\ 2+l \end{smallmatrix}\right] \times k} + 32 \cdot \Gamma_{i-5}^{\left[\begin{smallmatrix} 2 \\ 2+l \end{smallmatrix}\right] \times k}$$

$$(7.21) \quad A_i^{[3,0,l,k]} = \begin{cases} 1 & \text{if } i = 3 \\ -3 & \text{if } i = 4 \\ 2 & \text{if } i = 5 \\ -2^{k+2} & \text{if } i = 6 \\ 5 \cdot 2^{k+2} & \text{if } i = 7 \\ 3 \cdot 2^{k+2i-9} & \text{if } 8 \leq i \leq l+5 \\ -5 \cdot 2^{k+2l} & \text{if } i = 6+l \\ 2^{k+2l+2} & \text{if } i = 7+l \\ 0 & \text{if } 8+l \\ 3 \cdot 2^{2k+l+4} & \text{if } i = 9+l \end{cases}$$

We have in the case $l \geq 3, m \geq 1, s = 3$

$$(7.22) \quad A_i^{[3,m,l,k]} = \Gamma_{i-3}^{\left[\begin{smallmatrix} m+3 \\ m+3+l \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_{i-4}^{\left[\begin{smallmatrix} m+2 \\ m+3+l \end{smallmatrix} \right] \times k} - 8 \cdot \Gamma_{i-4}^{\left[\begin{smallmatrix} m+3 \\ m+2+l \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_{i-5}^{\left[\begin{smallmatrix} m+2 \\ m+2+l \end{smallmatrix} \right] \times k}$$

$$A_i^{[3,m,l,k]} = \begin{cases} 1 & \text{if } i = 3 \\ -3 & \text{if } i = 4 \\ 2 & \text{if } i = 5 \\ 0 & \text{if } 6 \leq i \leq m+5 \\ -2^{k+m+2} & \text{if } i = m+6 \\ 5 \cdot 2^{k+m+2} & \text{if } i = m+7 \\ 3 \cdot 2^{k-m+2i-9} & \text{if } m+8 \leq i \leq m+l+5 \\ -5 \cdot 2^{k+m+2l} & \text{if } i = m+6+l \\ 2^{k+m+2l+2} & \text{if } i = m+7+l \\ 0 & \text{if } m+8+l \leq i \leq 2m+l+8 \\ 3 \cdot 2^{2k+2m+l+4} & \text{if } i = 2m+9+l \end{cases}$$

We have in the case $l = 0, m \geq 0, s \geq 4$

$$(7.23) \quad A_i^{[s,m,0,k]} = \Gamma_{i-s}^{\left[\begin{smallmatrix} m+s \\ m+s \end{smallmatrix} \right] \times k} - 12 \cdot \Gamma_{i-s-1}^{\left[\begin{smallmatrix} m+s-1 \\ m+s \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_{i-s-2}^{\left[\begin{smallmatrix} m+s-1 \\ m+s-1 \end{smallmatrix} \right] \times k}$$

$$A_i^{[s,m,0,k]} = \begin{cases} 1 & \text{if } i = s \\ -3 & \text{if } i = s+1 \\ 2 & \text{if } i = s+2 \\ 0 & \text{if } s+3 \leq i \leq m+2s-1 \\ -3 \cdot 2^{k+m+s-1} & \text{if } i = m+2s \\ 3 \cdot 2^{k+m+s-1} & \text{if } i = m+2s+1 \\ 0 & \text{if } m+2s+2 \leq i \leq 2m+3s-1 \\ 3 \cdot 2^{2k+2s+2m-2} & \text{if } i = 2m+3s \end{cases}$$

We have in the case $l = 1, m \geq 0, s \geq 4$

$$A_i^{[s,m,1,k]} = \Gamma_{i-s}^{\left[\begin{smallmatrix} m+s \\ m+s+1 \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_{i-s-1}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+1 \end{smallmatrix} \right] \times k} - 8 \cdot \Gamma_{i-s-1}^{\left[\begin{smallmatrix} m+s \\ m+s \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_{i-s-2}^{\left[\begin{smallmatrix} m+s-1 \\ m+s \end{smallmatrix} \right] \times k}$$

$$(7.24) \quad A_i^{[s,m,1,k]} = \begin{cases} 1 & \text{if } i = s \\ -3 & \text{if } i = s + 1 \\ 2 & \text{if } i = s + 2 \\ 0 & \text{if } s + 3 \leq i \leq m + 2s - 1 \\ -2^{k+m+s-1} & \text{if } i = m + 2s \\ -3 \cdot 2^{k+m+s-1} & \text{if } i = m + 2s + 1 \\ 2^{k+m+s+1} & \text{if } i = m + 2s + 2 \\ 0 & \text{if } m + 2s + 3 \leq i \leq 2m + 3s \\ 3 \cdot 2^{2k+2s+2m-1} & \text{if } i = 2m + 3s + 1 \end{cases}$$

We have in the case $l = 2, m \geq 0, s \geq 4$

$$(7.25) \quad A_i^{[s,m,2,k]} = \Gamma_{i-s}^{\left[\begin{smallmatrix} m+s \\ m+s+2 \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_{i-s-1}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+2 \end{smallmatrix} \right] \times k} - 8 \cdot \Gamma_{i-s-1}^{\left[\begin{smallmatrix} m+s \\ m+s+1 \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_{i-s-2}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+1 \end{smallmatrix} \right] \times k}$$

$$A_i^{[s,m,2,k]} = \begin{cases} 1 & \text{if } i = s \\ -3 & \text{if } i = s + 1 \\ 2 & \text{if } i = s + 2 \\ 0 & \text{if } s + 3 \leq i \leq m + 2s - 1 \\ -2^{k+m+s-1} & \text{if } i = m + 2s \\ 5 \cdot 2^{k+m+s-1} & \text{if } i = m + 2s + 1 \\ -5 \cdot 2^{k+m+s+1} & \text{if } i = m + 2s + 2 \\ 2^{k+m+s+3} & \text{if } i = m + 2s + 3 \\ 0 & \text{if } m + 2s + 4 \leq i \leq 2m + 3s + 1 \\ 3 \cdot 2^{2k+2s+2m} & \text{if } i = 2m + 3s + 2 \end{cases}$$

We have in the case $l \geq 3, m \geq 0, s \geq 4$

$$(7.26) \quad A_i^{[s,m,l,k]} = \Gamma_{i-s}^{\left[\begin{smallmatrix} m+s \\ m+s+l \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_{i-s-1}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l \end{smallmatrix} \right] \times k} - 8 \cdot \Gamma_{i-s-1}^{\left[\begin{smallmatrix} m+s \\ m+s+l-1 \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_{i-s-2}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l-1 \end{smallmatrix} \right] \times k}$$

$$A_i^{[s,m,l,k]} = \begin{cases} 1 & \text{if } i = s \\ -3 & \text{if } i = s + 1 \\ 2 & \text{if } i = s + 2 \\ 0 & \text{if } s + 3 \leq i \leq m + 2s - 1 \\ -2^{k+m+s-1} & \text{if } i = m + 2s \\ 5 \cdot 2^{k+m+s-1} & \text{if } i = m + 2s + 1 \\ 3 \cdot 2^{k-m-3s+2i-3} & \text{if } m + 2s + 2 \leq i \leq m + 2s + l - 1 \\ -5 \cdot 2^{k+m+s+2l-3} & \text{if } i = m + 2s + l \\ 2^{k+m+s+2l-1} & \text{if } i = m + 2s + l + 1 \\ 0 & \text{if } m + 2s + l + 2 \leq i \leq 2m + 3s + l - 1 \\ 3 \cdot 2^{2k+2s+2m+l-2} & \text{if } i = 2m + 3s + l \end{cases}$$

Proof. By way of example we prove (7.26)

Let $s \geq 4, m \geq 0, l \geq 4, i = s + j$, for $0 \leq j \leq 2m + 2s + l$

We can write

$$(7.27) \quad A_{s+j}^{[s,m,l,k]} = \Gamma_j^{\left[\begin{smallmatrix} m+s \\ m+s+l \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_{j-1}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l \end{smallmatrix} \right] \times k} - 8 \cdot \Gamma_{j-1}^{\left[\begin{smallmatrix} m+s \\ m+s+l-1 \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_{j-2}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l-1 \end{smallmatrix} \right] \times k} \quad \text{for } 0 \leq j \leq 2m + 2s + l$$

From Theorem 3.13 (see section 3 in [2]) we obtain

$$\begin{aligned}
\underline{j=0} \quad A_{s+0}^{[s,m,l,k]} &= \Gamma_0^{\left[\begin{smallmatrix} m+s \\ m+s+l \end{smallmatrix} \right] \times k} = 1 \\
\underline{j=1} \quad A_{s+1}^{[s,m,l,k]} &= \Gamma_1^{\left[\begin{smallmatrix} m+s \\ m+s+l \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_0^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l \end{smallmatrix} \right] \times k} - 8 \cdot \Gamma_0^{\left[\begin{smallmatrix} m+s \\ m+s+l-1 \end{smallmatrix} \right] \times k} = 9 - 12 = -3 \\
\underline{j=2} \quad A_{s+2}^{[s,m,l,k]} &= \Gamma_2^{\left[\begin{smallmatrix} m+s \\ m+s+l \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_1^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l \end{smallmatrix} \right] \times k} - 8 \cdot \Gamma_1^{\left[\begin{smallmatrix} m+s \\ m+s+l-1 \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_0^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l-1 \end{smallmatrix} \right] \times k} = 78 - 12 \cdot 9 + 32 = 2
\end{aligned}$$

$$\underline{3 \leq j \leq m+s-1}$$

$$\begin{aligned}
A_{s+j}^{[s,m,l,k]} &= \Gamma_j^{\left[\begin{smallmatrix} m+s \\ m+s+l \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_{j-1}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l \end{smallmatrix} \right] \times k} - 8 \cdot \Gamma_{j-1}^{\left[\begin{smallmatrix} m+s \\ m+s+l-1 \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_{j-2}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l-1 \end{smallmatrix} \right] \times k} \\
&= (21 \cdot 2^{3j-4} - 3 \cdot 2^{2j-3}) - 12 \cdot (21 \cdot 2^{3j-7} - 3 \cdot 2^{2j-5}) + 32 \cdot (21 \cdot 2^{3j-10} - 3 \cdot 2^{2j-7}) = 0
\end{aligned}$$

$$\underline{j = m+s}$$

$$\begin{aligned}
A_{2s+m}^{[s,m,l,k]} &= \Gamma_{m+s}^{\left[\begin{smallmatrix} m+s \\ m+s+l \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_{m+s-1}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l \end{smallmatrix} \right] \times k} - 8 \cdot \Gamma_{m+s-1}^{\left[\begin{smallmatrix} m+s \\ m+s+l-1 \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_{m+s-2}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l-1 \end{smallmatrix} \right] \times k} \\
&= (2^{k+m+s-1} + 21 \cdot 2^{3(m+s)-4} - 11 \cdot 2^{2(m+s)-3}) - 4 \cdot (2^{k+m+s-2} + 21 \cdot 2^{3(m+s-1)-4} - 11 \cdot 2^{2(m+s-1)-3}) \\
&\quad - 8 \cdot (21 \cdot 2^{3(m+s-1)-4} - 3 \cdot 2^{2(m+s-1)-3}) + 32 \cdot (21 \cdot 2^{3(m+s-2)-4} - 3 \cdot 2^{2(m+s-2)-3}) = -2^{k+m+s-1}
\end{aligned}$$

$$\underline{j = m+s+1}$$

$$\begin{aligned}
A_{2s+m+1}^{[s,m,l,k]} &= \Gamma_{m+s+1}^{\left[\begin{smallmatrix} m+s \\ m+s+l \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_{m+s}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l \end{smallmatrix} \right] \times k} - 8 \cdot \Gamma_{m+s}^{\left[\begin{smallmatrix} m+s \\ m+s+l-1 \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_{m+s-1}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l-1 \end{smallmatrix} \right] \times k} \\
&= (3 \cdot 2^{k+m+s-1} + 21 \cdot (2^{3m+3s-1} - 2^{2m+2s-1})) - (3 \cdot 2^{k+m+s} + 21 \cdot (2^{3m+3s-2} - 2^{2m+2s-1})) \\
&\quad - (2^{k+m+s+2} + 21 \cdot 2^{3m+3s-1} - 11 \cdot 2^{2m+2s}) + (2^{k+m+s+3} + 21 \cdot 2^{3m+3s-2} - 11 \cdot 2^{2m+2s}) = 5 \cdot 2^{k+m+s-1}
\end{aligned}$$

$$\underline{j = m+s+2}$$

$$\begin{aligned}
A_{2s+m+2}^{[s,m,l,k]} &= \Gamma_{m+s+2}^{\left[\begin{smallmatrix} m+s \\ m+s+l \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_{m+s+1}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l \end{smallmatrix} \right] \times k} - 8 \cdot \Gamma_{m+s+1}^{\left[\begin{smallmatrix} m+s \\ m+s+l-1 \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_{m+s}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l-1 \end{smallmatrix} \right] \times k} \\
&= (3 \cdot 2^{k+m+s+1} + 21 \cdot (2^{3m+3s+2} - 2^{2m+2s+2})) - (3 \cdot 2^{k+m+s+2} + 21 \cdot (2^{3m+3s+1} - 2^{2m+2s+2})) \\
&\quad - (3 \cdot 2^{k+m+s+2} + 21 \cdot (2^{3m+3s+2} - 2^{2m+2s+1})) + (3 \cdot 2^{k+m+s+3} + 21 \cdot (2^{3m+3s+1} - 2^{2m+2s+1})) = 3 \cdot 2^{k+m+s+1}
\end{aligned}$$

$$\underline{m+s+2 \leq j \leq m+s+l-1}$$

$$\begin{aligned}
A_{s+j}^{[s,m,l,k]} &= \Gamma_j^{\left[\begin{smallmatrix} m+s \\ m+s+l \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_{j-1}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l \end{smallmatrix} \right] \times k} - 8 \cdot \Gamma_{j-1}^{\left[\begin{smallmatrix} m+s \\ m+s+l-1 \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_{j-2}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l-1 \end{smallmatrix} \right] \times k} \\
&= (3 \cdot 2^{k-m-s+2j-3} + 21 \cdot (2^{3j-4} - 2^{-m-s+3j-4})) - (3 \cdot 2^{k-m-s+2j-2} + 21 \cdot (2^{3j-5} - 2^{-m-s+3j-4})) \\
&\quad - (3 \cdot 2^{k-m-s+2j-2} + 21 \cdot (2^{3j-4} - 2^{-m-s+3j-4})) + (3 \cdot 2^{k-m-s+2j-1} + 21 \cdot (2^{3j-5} - 2^{-m-s+3j-4})) \\
&= 3 \cdot 2^{k-m-s+2j-3}
\end{aligned}$$

$$\underline{j = m+s+l}$$

$$\begin{aligned}
A_{m+2s+l}^{[s,m,l,k]} &= \Gamma_{m+s+l}^{\left[\begin{smallmatrix} m+s \\ m+s+l \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_{m+s+l-1}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l \end{smallmatrix} \right] \times k} - 8 \cdot \Gamma_{m+s+l-1}^{\left[\begin{smallmatrix} m+s \\ m+s+l-1 \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_{m+s+l-2}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l-1 \end{smallmatrix} \right] \times k} \\
&= (11 \cdot 2^{k+m+s+2l-3} + 21 \cdot 2^{3m+3s+3l-4} - 53 \cdot 2^{2m+2s+3l-4}) - (3 \cdot 2^{k+m+s+2l-2} + 21 \cdot (2^{3m+3s+3l-5} - 2^{2m+2s+3l-4}))
\end{aligned}$$

$$\begin{aligned}
& - (11 \cdot 2^{k+m+s+2l-2} + 21 \cdot 2^{3m+3s+3l-4} - 53 \cdot 2^{2m+2s+3l-4}) + (3 \cdot 2^{k+m+s+2l-1} + 21 \cdot (2^{3m+3s+3l-5} - 2^{2m+2s+3l-4})) \\
& = -5 \cdot 2^{k+m+s+2l-3}
\end{aligned}$$

$$\underline{j = m + s + l + 1}$$

$$\begin{aligned}
A_{m+2s+l+1}^{[s,m,l,k]} &= \Gamma_{m+s+l+1}^{\left[\begin{smallmatrix} m+s \\ m+s+l \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_{m+s+l}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l \end{smallmatrix} \right] \times k} - 8 \cdot \Gamma_{m+s+l}^{\left[\begin{smallmatrix} m+s \\ m+s+l-1 \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_{m+s+l-1}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l-1 \end{smallmatrix} \right] \times k} \\
&= (21 \cdot (2^{k+m+s+2l-1} + 2^{3m+3s+3l-1} - 5 \cdot 2^{2m+2s+3l-1})) - (11 \cdot 2^{k+m+s+2l} + 21 \cdot 2^{3m+3s+3l-2} - 53 \cdot 2^{2m+2s+3l-1}) \\
&- (21 \cdot (2^{k+m+s+2l} + 2^{3m+3s+3l-1} - 5 \cdot 2^{2m+2s+3l-1})) + (11 \cdot 2^{k+m+s+2l+1} + 21 \cdot 2^{3m+3s+3l-2} - 53 \cdot 2^{2m+2s+3l-1}) \\
&= 2^{k+m+s+2l-1}
\end{aligned}$$

$$\underline{m + s + l + 2 \leq j \leq 2m + 2s + l - 2}$$

$$\begin{aligned}
A_{s+j}^{[s,m,l,k]} &= \Gamma_j^{\left[\begin{smallmatrix} m+s \\ m+s+l \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_{j-1}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l \end{smallmatrix} \right] \times k} - 8 \cdot \Gamma_{j-1}^{\left[\begin{smallmatrix} m+s \\ m+s+l-1 \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_{j-2}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l-1 \end{smallmatrix} \right] \times k} \\
&= (21 \cdot (2^{k-2m-2s-l+3j-4} + 2^{3j-4} - 5 \cdot 2^{4j-2s-2m-l-5})) - (21 \cdot (2^{k-2m-2s-l+3j-4} + 2^{3j-5} - 5 \cdot 2^{4j-2s-2m-l-6})) \\
&- (21 \cdot (2^{k-2m-2s-l+3j-3} + 2^{3j-4} - 5 \cdot 2^{4j-2s-2m-l-5})) + (21 \cdot (2^{k-2m-2s-l+3j-3} + 2^{3j-5} - 5 \cdot 2^{4j-2s-2m-l-6})) \\
&= 0
\end{aligned}$$

$$\underline{j = 2m + 2s + l}$$

$$\begin{aligned}
A_{2m+3s+l}^{[s,m,l,k]} &= \Gamma_{2m+2s+l}^{\left[\begin{smallmatrix} m+s \\ m+s+l \end{smallmatrix} \right] \times k} - 4 \cdot \Gamma_{2m+2s+l-1}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l \end{smallmatrix} \right] \times k} - 8 \cdot \Gamma_{2m+2s+l-1}^{\left[\begin{smallmatrix} m+s \\ m+s+l-1 \end{smallmatrix} \right] \times k} + 32 \cdot \Gamma_{2m+2s+l-2}^{\left[\begin{smallmatrix} m+s-1 \\ m+s+l-1 \end{smallmatrix} \right] \times k} \\
&= (2^{2k+2s+2m+l-2} - 3 \cdot 2^{k+4s+4m+2l-4} + 2^{6s+6m+3l-5}) - (2^{2k+2s+2m+l-1} - 3 \cdot 2^{k+4s+4m+2l-4} + 2^{6s+6m+3l-6}) \\
&- (2^{2k+2s+2m+l} - 3 \cdot 2^{k+4s+4m+2l-3} + 2^{6s+6m+3l-5}) + (2^{2k+2s+2m+l+1} - 3 \cdot 2^{k+4s+4m+2l-3} + 2^{6s+6m+3l-6}) \\
&= 3 \cdot 2^{2k+2s+2m+l-2}
\end{aligned}$$

□

7.5. Computation of $\omega_i(s, m, l, k+1) - \omega_i(s, m, l, k)$.

Lemma 7.4. *We have*

The case $s = 2$

The case $l = 0, m = 0, s = 2, k > i$

$$(7.28) \quad \omega_i(2, 0, 0, k+1) - \omega_i(2, 0, 0, k) = \begin{cases} 2^{k+1} & \text{if } i = 2 \\ -3 \cdot 2^{k+1} & \text{if } i = 3 \\ -9 \cdot 2^{2k+2} + 2^{k+2} & \text{if } i = 4 \\ 9 \cdot 2^{2k+2} & \text{if } i = 5 \\ 21 \cdot 2^{3k+3} & \text{if } i = 6 \end{cases}$$

The case $l = 0, m = 1, s = 2, k > i$

$$(7.29) \quad \omega_i(2, 1, 0, k+1) - \omega_i(2, 1, 0, k) = \begin{cases} 2^{k+1} & \text{if } i = 2 \\ -3 \cdot 2^{k+1} & \text{if } i = 3 \\ 2^{k+2} & \text{if } i = 4 \\ -9 \cdot 2^{2k+3} & \text{if } i = 5 \\ 9 \cdot 2^{2k+3} & \text{if } i = 6 \\ 0 & \text{if } i = 7 \\ 21 \cdot 2^{3k+5} & \text{if } i = 8 \end{cases}$$

The case $l = 0, m \geq 2, s = 2, k > i$

$$(7.30) \quad \omega_i(2, m, 0, k+1) - \omega_i(2, m, 0, k) = \begin{cases} 2^{k+1} & \text{if } i = 2 \\ -3 \cdot 2^{k+1} & \text{if } i = 3 \\ 2^{k+2} & \text{if } i = 4 \\ 0 & \text{if } 5 \leq i \leq m+3 \\ -9 \cdot 2^{2k+m+2} & \text{if } i = m+4 \\ 9 \cdot 2^{2k+m+2} & \text{if } i = m+5 \\ 0 & \text{if } m+7 \leq i \leq 2m+5 \\ 21 \cdot 2^{3k+2m+3} & \text{if } i = 2m+6 \end{cases}$$

The case $l = 1, m = 0, s = 2, k > i$

$$(7.31) \quad \omega_i(2, 0, 1, k+1) - \omega_i(2, 0, 1, k) = \begin{cases} 2^{k+1} & \text{if } i = 2 \\ -3 \cdot 2^{k+1} & \text{if } i = 3 \\ -3 \cdot 2^{2k+2} + 2^{k+2} & \text{if } i = 4 \\ -9 \cdot 2^{2k+2} & \text{if } i = 5 \\ 3 \cdot 2^{2k+4} & \text{if } i = 6 \\ 21 \cdot 2^{3k+4} & \text{if } i = 7 \end{cases}$$

The case $l = 1, m = 1, s = 2, k > i$

$$(7.32) \quad \omega_i(2, 1, 1, k+1) - \omega_i(2, 1, 1, k) = \begin{cases} 2^{k+1} & \text{if } i = 2 \\ -3 \cdot 2^{k+1} & \text{if } i = 3 \\ 2^{k+2} & \text{if } i = 4 \\ -3 \cdot 2^{2k+3} & \text{if } i = 5 \\ -9 \cdot 2^{2k+3} & \text{if } i = 6 \\ 3 \cdot 2^{2k+5} & \text{if } i = 7 \\ 0 & \text{if } i = 8 \\ 21 \cdot 2^{3k+6} & \text{if } i = 9 \end{cases}$$

The case $l = 1, m \geq 2, s = 2, k > i$

$$(7.33) \quad \omega_i(2, m, 1, k+1) - \omega_i(2, m, 1, k) = \begin{cases} 2^{k+1} & \text{if } i = 2 \\ -3 \cdot 2^{k+1} & \text{if } i = 3 \\ 2^{k+2} & \text{if } i = 4 \\ 0 & \text{if } 5 \leq i \leq m+3 \\ 3 \cdot 2^{2k+m+2} & \text{if } i = m+4 \\ -9 \cdot 2^{2k+m+2} & \text{if } i = m+5 \\ 3 \cdot 2^{2k+m+4} & \text{if } i = m+6 \\ 0 & \text{if } m+7 \leq i \leq 2m+6 \\ 21 \cdot 2^{3k+2m+4} & \text{if } i = 2m+7 \end{cases}$$

The case $l = 2, m = 0, s = 2, k > i$

$$(7.34) \quad \omega_i(2, 0, 2, k+1) - \omega_i(2, 0, 2, k) = \begin{cases} 2^{k+1} & \text{if } i = 2 \\ -3 \cdot 2^{k+1} & \text{if } i = 3 \\ -3 \cdot 2^{2k+2} + 2^{k+2} & \text{if } i = 4 \\ 15 \cdot 2^{2k+2} & \text{if } i = 5 \\ -15 \cdot 2^{2k+4} & \text{if } i = 6 \\ 3 \cdot 2^{2k+6} & \text{if } i = 7 \\ 21 \cdot 2^{3k+5} & \text{if } i = 8 \end{cases}$$

The case $l = 2, m = 1, s = 2, k > i$

$$(7.35) \quad \omega_i(2, 1, 2, k+1) - \omega_i(2, 1, 2, k) = \begin{cases} 2^{k+1} & \text{if } i = 2 \\ -3 \cdot 2^{k+1} & \text{if } i = 3 \\ 2^{k+2} & \text{if } i = 4 \\ -3 \cdot 2^{2k+3} & \text{if } i = 5 \\ 15 \cdot 2^{2k+3} & \text{if } i = 6 \\ -15 \cdot 2^{2k+5} & \text{if } i = 7 \\ 3 \cdot 2^{2k+7} & \text{if } i = 8 \\ 0 & \text{if } i = 9 \\ 21 \cdot 2^{3k+7} & \text{if } i = 10 \end{cases}$$

The case $l = 2, m \geq 2, s = 2, k > i$

$$(7.36) \quad \omega_i(2, m, 2, k+1) - \omega_i(2, m, 2, k) = \begin{cases} 2^{k+1} & \text{if } i = 2 \\ -3 \cdot 2^{k+1} & \text{if } i = 3 \\ 2^{k+2} & \text{if } i = 4 \\ 0 & \text{if } 5 \leq i \leq m+3 \\ -3 \cdot 2^{2k+m+2} & \text{if } i = m+4 \\ 15 \cdot 2^{2k+m+2} & \text{if } i = m+5 \\ -15 \cdot 2^{2k+m+4} & \text{if } i = m+6 \\ 3 \cdot 2^{2k+m+6} & \text{if } i = m+7 \\ 0 & \text{if } m+8 \leq i \leq 2m+7 \\ 21 \cdot 2^{3k+2m+5} & \text{if } i = 2m+8 \end{cases}$$

The case $l \geq 3, m = 0, s = 2, k > i$

$$(7.37) \quad \omega_i(2, 0, l, k+1) - \omega_i(2, 0, l, k) = \begin{cases} 2^{k+1} & \text{if } i = 2 \\ -3 \cdot 2^{k+1} & \text{if } i = 3 \\ -3 \cdot 2^{2k+2} + 2^{k+2} & \text{if } i = 4 \\ 15 \cdot 2^{2k+2} & \text{if } i = 5 \\ 9 \cdot 2^{2k+2i-8} & \text{if } 6 \leq i \leq l+3 \\ -15 \cdot 2^{2k+2l} & \text{if } i = l+4 \\ 3 \cdot 2^{2k+2l+2} & \text{if } i = l+5 \\ 21 \cdot 2^{3k+l+3} & \text{if } i = l+6 \end{cases}$$

The case $l \geq 3, m = 1, s = 2, k > i$

$$(7.38) \quad \omega_i(2, 1, l, k+1) - \omega_i(2, 1, l, k) = \begin{cases} 2^{k+1} & \text{if } i = 2 \\ -3 \cdot 2^{k+1} & \text{if } i = 3 \\ 2^{k+2} & \text{if } i = 4 \\ -3 \cdot 2^{2k+3} & \text{if } i = 5 \\ 15 \cdot 2^{2k+3} & \text{if } i = 6 \\ 9 \cdot 2^{2k+2i-9} & \text{if } 7 \leq i \leq l+4 \\ -15 \cdot 2^{2k+2l+1} & \text{if } i = l+5 \\ 3 \cdot 2^{2k+2l+3} & \text{if } i = l+6 \\ 0 & \text{if } i = l+7 \\ 21 \cdot 2^{3k+l+5} & \text{if } i = l+8 \end{cases}$$

The case $l \geq 3, m \geq 2, s = 2, k > i$

(7.39)

$$\omega_i(2, m, l, k+1) - \omega_i(2, m, l, k) = \begin{cases} 2^{k+1} & \text{if } i = 2 \\ -3 \cdot 2^{k+1} & \text{if } i = 3 \\ 2^{k+2} & \text{if } i = 4 \\ 0 & \text{if } 5 \leq i \leq m+3 \\ -3 \cdot 2^{2k+m+2} & \text{if } i = m+4 \\ 15 \cdot 2^{2k+m+2} & \text{if } i = m+5 \\ 9 \cdot 2^{2k-m+2i-8} & \text{if } m+6 \leq i \leq m+l+3 \\ -15 \cdot 2^{2k+m+2l} & \text{if } i = m+l+4 \\ 3 \cdot 2^{2k+m+2l+2} & \text{if } i = m+l+5 \\ 0 & \text{if } m+l+6 \leq i \leq 2m+l+5 \\ 21 \cdot 2^{3k+2m+l+3} & \text{if } i = 2m+l+6 \end{cases}$$

The case $s = 3$

The case $l = 0, m = 0, s = 3, k > i$

$$(7.40) \quad \omega_i(3, 0, 0, k+1) - \omega_i(3, 0, 0, k) = \begin{cases} 2^{k+2} & \text{if } i = 3 \\ -3 \cdot 2^{k+2} & \text{if } i = 4 \\ 2^{k+3} & \text{if } i = 5 \\ -9 \cdot 2^{2k+4} & \text{if } i = 6 \\ 9 \cdot 2^{2k+4} & \text{if } i = 7 \\ 0 & \text{if } i = 8 \\ 21 \cdot 2^{3k+6} & \text{if } i = 9 \end{cases}$$

The case $l = 0, m \geq 1, s = 3, k > i$

$$(7.41) \quad \omega_i(3, m, 0, k+1) - \omega_i(3, m, 0, k) = \begin{cases} 2^{k+2} & \text{if } i = 3 \\ -3 \cdot 2^{k+2} & \text{if } i = 4 \\ 2^{k+3} & \text{if } i = 5 \\ 0 & \text{if } 6 \leq i \leq m+5 \\ -9 \cdot 2^{2k+m+4} & \text{if } i = m+6 \\ 9 \cdot 2^{2k+m+4} & \text{if } i = m+7 \\ 0 & \text{if } m+8 \leq i \leq 2m+8 \\ 21 \cdot 2^{3k+2m+6} & \text{if } i = 2m+9 \end{cases}$$

The case $l = 1, m = 0, s = 3, k > i$

$$(7.42) \quad \omega_i(3, 0, 1, k+1) - \omega_i(3, 0, 1, k) = \begin{cases} 2^{k+2} & \text{if } i = 3 \\ -3 \cdot 2^{k+2} & \text{if } i = 4 \\ 2^{k+3} & \text{if } i = 5 \\ -3 \cdot 2^{2k+4} & \text{if } i = 6 \\ -9 \cdot 2^{2k+4} & \text{if } i = 7 \\ 3 \cdot 2^{2k+6} & \text{if } i = 8 \\ 0 & \text{if } i = 9 \\ 21 \cdot 2^{3k+7} & \text{if } i = 10 \end{cases}$$

The case $l = 1, m \geq 1, s = 3, k > i$

$$(7.43) \quad \omega_i(3, m, 1, k+1) - \omega_i(3, m, 1, k) = \begin{cases} 2^{k+2} & \text{if } i = 3 \\ -3 \cdot 2^{k+2} & \text{if } i = 4 \\ 2^{k+3} & \text{if } i = 5 \\ 0 & \text{if } 6 \leq i \leq m+5 \\ -3 \cdot 2^{2k+m+4} & \text{if } i = m+6 \\ -9 \cdot 2^{2k+m+4} & \text{if } i = m+7 \\ 3 \cdot 2^{2k+m+6} & \text{if } i = m+8 \\ 0 & \text{if } m+9 \leq i \leq 2m+9 \\ 21 \cdot 2^{3k+2m+7} & \text{if } i = 2m+10 \end{cases}$$

The case $l = 2, m = 0, s = 3, k > i$

$$(7.44) \quad \omega_i(3, 0, 2, k+1) - \omega_i(3, 0, 2, k) = \begin{cases} 2^{k+2} & \text{if } i = 3 \\ -3 \cdot 2^{k+2} & \text{if } i = 4 \\ 2^{k+3} & \text{if } i = 5 \\ -3 \cdot 2^{2k+4} & \text{if } i = 6 \\ 15 \cdot 2^{2k+4} & \text{if } i = 7 \\ -15 \cdot 2^{2k+6} & \text{if } i = 8 \\ 3 \cdot 2^{2k+8} & \text{if } i = 9 \\ 0 & \text{if } i = 10 \\ 21 \cdot 2^{3k+8} & \text{if } i = 11 \end{cases}$$

The case $l = 2, m \geq 1, s = 3, k > i$

$$(7.45) \quad \omega_i(3, m, 2, k+1) - \omega_i(3, m, 2, k) = \begin{cases} 2^{k+2} & \text{if } i = 3 \\ -3 \cdot 2^{k+2} & \text{if } i = 4 \\ 2^{k+3} & \text{if } i = 5 \\ 0 & \text{if } 6 \leq i \leq m+5 \\ -3 \cdot 2^{2k+m+4} & \text{if } i = m+6 \\ 15 \cdot 2^{2k+m+4} & \text{if } i = m+7 \\ -15 \cdot 2^{2k+m+6} & \text{if } i = m+8 \\ 3 \cdot 2^{2k+m+8} & \text{if } i = m+9 \\ 0 & \text{if } m+10 \leq i \leq 2m+10 \\ 21 \cdot 2^{3k+2m+8} & \text{if } i = 2m+11 \end{cases}$$

The case $l \geq 3, m = 0, s = 3, k > i$

$$(7.46) \quad \omega_i(3, 0, l, k+1) - \omega_i(3, 0, l, k) = \begin{cases} 2^{k+2} & \text{if } i = 3 \\ -3 \cdot 2^{k+2} & \text{if } i = 4 \\ 2^{k+3} & \text{if } i = 5 \\ -3 \cdot 2^{2k+4} & \text{if } i = 6 \\ 15 \cdot 2^{2k+4} & \text{if } i = 7 \\ 9 \cdot 2^{2k+2i-10} & \text{if } 8 \leq i \leq l+5 \\ -15 \cdot 2^{2k+2l+2} & \text{if } i = l+6 \\ 3 \cdot 2^{2k+2l+4} & \text{if } i = l+7 \\ 0 & \text{if } i = l+8 \\ 21 \cdot 2^{3k+l+6} & \text{if } i = l+9 \end{cases}$$

The case $l \geq 3, m \geq 1, s = 3, k > i$

$$(7.47) \quad \omega_i(3, m, l, k+1) - \omega_i(3, m, l, k) = \begin{cases} 2^{k+2} & \text{if } i = 3 \\ -3 \cdot 2^{k+2} & \text{if } i = 4 \\ 2^{k+3} & \text{if } i = 5 \\ 0 & \text{if } 6 \leq i \leq m+5 \\ -3 \cdot 2^{2k+m+4} & \text{if } i = m+6 \\ 15 \cdot 2^{2k+m+4} & \text{if } i = m+7 \\ 9 \cdot 2^{2k-m+2i-10} & \text{if } m+8 \leq i \leq m+l+5 \\ -15 \cdot 2^{2k+m+2l+2} & \text{if } i = m+l+6 \\ 3 \cdot 2^{2k+m+2l+4} & \text{if } i = m+l+7 \\ 0 & \text{if } m+l+8 \leq i \leq 2m+l+8 \\ 21 \cdot 2^{3k+2m+l+6} & \text{if } i = 2m+l+9 \end{cases}$$

The case $s \geq 4$

The case $l = 0, m \geq 0, s \geq 4, k > i$

(7.48)

$$\omega_i(s, m, 0, k+1) - \omega_i(s, m, 0, k) = \begin{cases} 2^{k+s-1} & \text{if } i = s \\ -3 \cdot 2^{k+s-1} & \text{if } i = s+1 \\ 2^{k+s} & \text{if } i = s+2 \\ 0 & \text{if } s+3 \leq i \leq m+2s-1 \\ -9 \cdot 2^{2k+2s+m-2} & \text{if } i = m+2s \\ 9 \cdot 2^{2k+2s+m-2} & \text{if } i = m+2s+1 \\ 0 & \text{if } m+2s+2 \leq i \leq 2m+3s-1 \\ 21 \cdot 2^{3k+3s+2m-3} & \text{if } i = 2m+3s \end{cases}$$

The case $l = 1, m \geq 0, s \geq 4, k > i$

(7.49)

$$\omega_i(s, m, 1, k+1) - \omega_i(s, m, 1, k) = \begin{cases} 2^{k+s-1} & \text{if } i = s \\ -3 \cdot 2^{k+s-1} & \text{if } i = s+1 \\ 2^{k+s} & \text{if } i = s+2 \\ 0 & \text{if } s+3 \leq i \leq m+2s-1 \\ -3 \cdot 2^{2k+2s+m-2} & \text{if } i = m+2s \\ -9 \cdot 2^{2k+2s+m-2} & \text{if } i = m+2s+1 \\ 3 \cdot 2^{2k+2s+m} & \text{if } i = m+2s+2 \\ 0 & \text{if } m+2s+3 \leq i \leq 2m+3s \\ 21 \cdot 2^{3k+3s+2m-2} & \text{if } i = 2m+3s+1 \end{cases}$$

The case $l = 2, m \geq 0, s \geq 4, k > i$

(7.50)

$$\omega_i(s, m, 2, k+1) - \omega_i(s, m, 2, k) = \begin{cases} 2^{k+s-1} & \text{if } i = s \\ -3 \cdot 2^{k+s-1} & \text{if } i = s+1 \\ 2^{k+s} & \text{if } i = s+2 \\ 0 & \text{if } s+3 \leq i \leq m+2s-1 \\ -3 \cdot 2^{2k+2s+m-2} & \text{if } i = m+2s \\ 15 \cdot 2^{2k+2s+m-2} & \text{if } i = m+2s+1 \\ -15 \cdot 2^{2k+2s+m} & \text{if } i = m+2s+2 \\ 3 \cdot 2^{2k+2s+m+2} & \text{if } i = m+2s+3 \\ 0 & \text{if } m+2s+4 \leq i \leq 2m+3s+1 \\ 21 \cdot 2^{3k+3s+2m-1} & \text{if } i = 2m+3s+2 \end{cases}$$

The case $l \geq 3, m \geq 0, s \geq 4, k > i$

(7.51)

$$\omega_i(s, m, l, k+1) - \omega_i(s, m, l, k) = \begin{cases} 2^{k+s-1} & \text{if } i = s \\ -3 \cdot 2^{k+s-1} & \text{if } i = s+1 \\ 2^{k+s} & \text{if } i = s+2 \\ 0 & \text{if } s+3 \leq i \leq m+2s-1 \\ -3 \cdot 2^{2k+2s+m-2} & \text{if } i = m+2s \\ 15 \cdot 2^{2k+2s+m-2} & \text{if } i = m+2s+1 \\ 9 \cdot 2^{2k-2s-m+2i-4} & \text{if } m+2s+2 \leq i \leq m+2s+l-1 \\ -15 \cdot 2^{2k+2s+m+2l-4} & \text{if } i = m+2s+l \\ 3 \cdot 2^{2k+2s+m+2l-2} & \text{if } i = m+2s+l+1 \\ 0 & \text{if } m+2s+l+2 \leq i \leq 2m+3s+l-1 \\ 21 \cdot 2^{3k+3s+2m+l-3} & \text{if } i = 2m+3s+l \end{cases}$$

Proof. By way of example we prove (7.51)

From (7.2) we have

$$\omega_i(s, m, l, k+1) - \omega_i(s, m, l, k) = 2^{k+s} A_i^{[s, m, l, k+1]} - 2^{k+s-1} A_i^{[s, m, l, k]}$$

We now deduce from (7.26)

$i = s$

$$\omega_s(s, m, l, k+1) - \omega_s(s, m, l, k) = 2^{k+s} \cdot 1 - 2^{k+s-1} \cdot 1 = 2^{k+s-1}$$

$i = s+1$

$$\omega_{s+1}(s, m, l, k+1) - \omega_{s+1}(s, m, l, k) = 2^{k+s} \cdot (-3) - 2^{k+s-1} \cdot (-3) = -3 \cdot 2^{k+s-1}$$

$i = s+2$

$$\omega_{s+2}(s, m, l, k+1) - \omega_{s+2}(s, m, l, k) = 2^{k+s} \cdot 2 - 2^{k+s-1} \cdot 2 = 2^{k+s}$$

$s+3 \leq i \leq m+2s-1$

$$\omega_i(s, m, l, k+1) - \omega_i(s, m, l, k) = 2^{k+s} \cdot 0 - 2^{k+s-1} \cdot 0 = 0$$

$i = m+2s$

$$\omega_{m+2s}(s, m, l, k+1) - \omega_{m+2s}(s, m, l, k) = 2^{k+s} \cdot (-2^{k+m+s}) - 2^{k+s-1} \cdot (-2^{k+m+s-1}) = -3 \cdot 2^{2k+m+2s-2}$$

$i = m+2s+1$

$$\omega_{m+2s+1}(s, m, l, k+1) - \omega_{m+2s+1}(s, m, l, k) = 2^{k+s} \cdot (5 \cdot 2^{k+m+s}) - 2^{k+s-1} \cdot (5 \cdot 2^{k+m+s-1}) = 15 \cdot 2^{2k+m+2s-2}$$

$m+2s+2 \leq i \leq m+2s+l-1$

$$\omega_i(s, m, l, k+1) - \omega_i(s, m, l, k) = 2^{k+s} \cdot (3 \cdot 2^{k-m-3s+2i-2}) - 2^{k+s-1} \cdot (3 \cdot 2^{k-m-3s+2i-3}) = 9 \cdot 2^{2k-2s-m+2i-4}$$

$i = m+2s+l$

$$\omega_{m+2s+l}(s, m, l, k+1) - \omega_{m+2s+l}(s, m, l, k) = 2^{k+s} \cdot (-5 \cdot 2^{k+m+s+2l-2}) - 2^{k+s-1} \cdot (-5 \cdot 2^{k+m+s+2l-3}) = -15 \cdot 2^{2k+2s+m+2l-4}$$

$i = m+2s+l+1$

$$\omega_{m+2s+l+1}(s, m, l, k+1) - \omega_{m+2s+l+1}(s, m, l, k) = 2^{k+s} \cdot 2^{k+m+s+2l} - 2^{k+s-1} \cdot 2^{k+m+s+2l-1} = 3 \cdot 2^{2k+2s+m+2l-2}$$

$m+2s+l+2 \leq i \leq 2m+3s+l-1$

$$\omega_i(s, m, l, k+1) - \omega_i(s, m, l, k) = 2^{k+s} \cdot 0 - 2^{k+s-1} \cdot 0 = 0$$

$i = 2m+3s+l$

$$\begin{aligned} \omega_{2m+3s+l}(s, m, l, k+1) - \omega_{2m+3s+l}(s, m, l, k) &= 2^{k+s} \cdot (3 \cdot 2^{2k+2s+2m+l}) - 2^{k+s-1} \cdot (3 \cdot 2^{2k+2s+2m+l-2}) \\ &= 21 \cdot 2^{3k+3s+2m+l-3} \end{aligned}$$

□

8. Computation of $\Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}}$ for $0 \leq j \leq 2s$

8.1. Notation.

Definition 8.1. We define

$$\bullet \quad \Delta_k \Gamma_i^{\begin{bmatrix} s \\ s+m \\ s+m+l \end{bmatrix}} = \Gamma_i^{\begin{bmatrix} s \\ s+m \\ s+m+l \end{bmatrix}} \times (k+1) - \Gamma_i^{\begin{bmatrix} s \\ s+m \\ s+m+l \end{bmatrix}} \times k$$

$$\bullet \quad \Delta_k \omega_i(s, m, l) = \omega_i(s, m, l, k+1) - \omega_i(s, m, l, k)$$

8.2. Introduction. We adapt the method used in Section 12 of [2] to compute explicitly

$$\Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} \quad \text{for } 0 \leq j \leq 2s$$

8.3. A reduction formula for $\Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}}$ for $0 \leq j \leq 2s$.

Lemma 8.2. We have the following reduction formulas :

$$(8.1) \quad \Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} = 4^{2s-3} \cdot \left[\Delta_k \Gamma_{j-s+3}^{\begin{bmatrix} \frac{1}{2} \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{j-s+2}^{\begin{bmatrix} \frac{1}{2} \\ s-1 \\ s \end{bmatrix}} \right] \\ + \Delta_k \omega_{s+j}(s, 0, 0) + \sum_{q=1}^{s-2} 4^{2q-1} \cdot \Delta_k \omega_{s-q+(j-q+1)}(s-q, 1, q-1) + \sum_{q=1}^{s-2} 4^{2q} \cdot \Delta_k \omega_{s-q+(j-q)}(s-q, 0, q) \quad \text{for } 0 \leq j \leq s-1$$

$$(8.2) \quad \Delta_k \Gamma_{2s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} = 4^{2s-3} \cdot \left[\Delta_k \Gamma_{j+3}^{\begin{bmatrix} \frac{1}{2} \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{j+2}^{\begin{bmatrix} \frac{1}{2} \\ s-1 \\ s \end{bmatrix}} \right] \\ + \Delta_k \omega_{2s+j}(s, 0, 0) + \sum_{q=1}^{s-2} 4^{2q-1} \cdot \Delta_k \omega_{2(s-q)+(j+1)}(s-q, 1, q-1) + \sum_{q=1}^{s-2} 4^{2q} \cdot \Delta_k \omega_{2(s-q)+j}(s-q, 0, q) \quad \text{for } 0 \leq j \leq s$$

Proof. From Definition 5.1 and Definition 8.1 we obtain the following formula

$$(8.3) \quad \Delta_k \omega_i(s, m, l) = \Delta_k \Gamma_i^{\begin{bmatrix} s \\ s+m \\ s+m+l \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{i-1}^{\begin{bmatrix} s \\ s+m-1 \\ s+m+l \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{i-1}^{\begin{bmatrix} s \\ s+m \\ s+m+l-1 \end{bmatrix}} + 32 \cdot \Delta_k \Gamma_{i-2}^{\begin{bmatrix} s \\ s+m-1 \\ s+m+l-1 \end{bmatrix}}$$

The case $i = s + j$, $0 \leq j \leq s - 1$

From (8.3) we obtain successively the following equations

$$\begin{aligned}
& \Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} = 8 \cdot \left[\Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} \right] + \Delta_k \omega_{s+j}(s, 0, 0) \\
& 4 \cdot \left[\Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-2}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} \right] = 4 \cdot \left[8 \cdot \left[\Delta_k \Gamma_{s+j-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} \right] + \Delta_k \omega_{s+j-1}(s-1, 1, 0) \right] \\
& 4^2 \cdot \left[\Delta_k \Gamma_{s+j-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-3}^{\begin{bmatrix} s-2 \\ s-1 \\ s \end{bmatrix}} \right] = 4^2 \cdot \left[8 \cdot \left[\Delta_k \Gamma_{s+j-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-4}^{\begin{bmatrix} s-2 \\ s-1 \\ s-1 \end{bmatrix}} \right] + \Delta_k \omega_{s+j-2}(s-1, 0, 1) \right] \\
& 4^3 \cdot \left[\Delta_k \Gamma_{s+j-3}^{\begin{bmatrix} s-2 \\ s-1 \\ s \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-4}^{\begin{bmatrix} s-2 \\ s \\ s \end{bmatrix}} \right] = 4^3 \cdot \left[8 \cdot \left[\Delta_k \Gamma_{s+j-4}^{\begin{bmatrix} s-2 \\ s-1 \\ s-1 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-5}^{\begin{bmatrix} s-2 \\ s-1 \\ s-1 \end{bmatrix}} \right] + \Delta_k \omega_{s+j-3}(s-2, 1, 1) \right] \\
& 4^4 \cdot \left[\Delta_k \Gamma_{s+j-4}^{\begin{bmatrix} s-2 \\ s \\ s \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-5}^{\begin{bmatrix} s-3 \\ s-2 \\ s \end{bmatrix}} \right] = 4^4 \cdot \left[8 \cdot \left[\Delta_k \Gamma_{s+j-5}^{\begin{bmatrix} s-2 \\ s-1 \\ s-1 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-6}^{\begin{bmatrix} s-3 \\ s-2 \\ s-1 \end{bmatrix}} \right] + \Delta_k \omega_{s+j-4}(s-2, 0, 2) \right] \\
& \vdots \\
& 4^{2p-1} \cdot \left[\Delta_k \Gamma_{s+j-(2p-1)}^{\begin{bmatrix} s-p \\ s-(p-1) \\ s \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-2p}^{\begin{bmatrix} s-p \\ s \\ s \end{bmatrix}} \right] = 4^{2p-1} \cdot \left[8 \cdot \left[\Delta_k \Gamma_{s+j-2p}^{\begin{bmatrix} s-p \\ s-1 \\ s-1 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-2p-1}^{\begin{bmatrix} s-p \\ s-p \\ s-1 \end{bmatrix}} \right] + \Delta_k \omega_{s+j-(2p-1)}(s-p, 1, p-1) \right] \\
& 4^{2p} \cdot \left[\Delta_k \Gamma_{s+j-2p}^{\begin{bmatrix} s-p \\ s-p \\ s \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-(2p+1)}^{\begin{bmatrix} s-p-1 \\ s-p \\ s \end{bmatrix}} \right] = 4^{2p} \cdot \left[8 \cdot \left[\Delta_k \Gamma_{s+j-(2p+1)}^{\begin{bmatrix} s-p \\ s-1 \\ s-1 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-(2p+2)}^{\begin{bmatrix} s-p-1 \\ s-p \\ s-1 \end{bmatrix}} \right] + \Delta_k \omega_{s+j-2p}(s-p, 0, p) \right]
\end{aligned}$$

By summing the left- hand side of the above equations we obtain:

$$\begin{aligned}
& \Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} + \sum_{q=1}^p 4^{2q-1} \cdot \left[\Delta_k \Gamma_{s+j-(2q-1)}^{\begin{bmatrix} s-q \\ s-(q-1) \\ s \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-2q}^{\begin{bmatrix} s-q \\ s \\ s \end{bmatrix}} \right] + \sum_{q=1}^p 4^{2q} \cdot \left[\Delta_k \Gamma_{s+j-2q}^{\begin{bmatrix} s-q \\ s-q \\ s \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-(2q+1)}^{\begin{bmatrix} s-q-1 \\ s-q \\ s \end{bmatrix}} \right] \\
& = \Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} + \sum_{q=1}^p 4^{2q-1} \cdot \Delta_k \Gamma_{s+j-(2q-1)}^{\begin{bmatrix} s-q \\ s-(q-1) \\ s \end{bmatrix}} + \sum_{q=1}^p 4^{2q} \cdot \Delta_k \Gamma_{s+j-2q}^{\begin{bmatrix} s-q \\ s-q \\ s \end{bmatrix}} - \sum_{q=1}^p 4^{2q} \cdot \Delta_k \Gamma_{s+j-2q}^{\begin{bmatrix} s-q \\ s \\ s \end{bmatrix}} - \sum_{q=1}^p 4^{2q+1} \cdot \Delta_k \Gamma_{s+j-(2q+1)}^{\begin{bmatrix} s-q-1 \\ s-q \\ s \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} \\
& = \Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} + \sum_{q=1}^p 4^{2q-1} \cdot \Delta_k \Gamma_{s+j-(2q-1)}^{\begin{bmatrix} s-q \\ s-(q-1) \\ s \end{bmatrix}} + \sum_{q=1}^p 4^{2q} \cdot \Delta_k \Gamma_{s+j-2q}^{\begin{bmatrix} s-q \\ s-q \\ s \end{bmatrix}} - \sum_{q=1}^p 4^{2q} \cdot \Delta_k \Gamma_{s+j-2q}^{\begin{bmatrix} s-q \\ s \\ s \end{bmatrix}} - \sum_{q=0}^p 4^{2q+1} \cdot \Delta_k \Gamma_{s+j-(2q+1)}^{\begin{bmatrix} s-q-1 \\ s-q \\ s \end{bmatrix}} \\
& = \Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} + \sum_{q=1}^p 4^{2q-1} \cdot \Delta_k \Gamma_{s+j-(2q-1)}^{\begin{bmatrix} s-q \\ s-(q-1) \\ s \end{bmatrix}} + \sum_{q=1}^p 4^{2q} \cdot \Delta_k \Gamma_{s+j-2q}^{\begin{bmatrix} s-q \\ s-q \\ s \end{bmatrix}} - \sum_{q=1}^p 4^{2q} \cdot \Delta_k \Gamma_{s+j-2q}^{\begin{bmatrix} s-q \\ s \\ s \end{bmatrix}} - \sum_{q=1}^{p+1} 4^{2q-1} \cdot \Delta_k \Gamma_{s+j-(2q-1)}^{\begin{bmatrix} s-q \\ s-(q-1) \\ s \end{bmatrix}} \\
& = \Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 4^{2p+1} \cdot \Delta_k \Gamma_{s+j-2p-1}^{\begin{bmatrix} s-p-1 \\ s-p \\ s \end{bmatrix}}
\end{aligned}$$

By summing the right-hand side of the above equations we obtain:

$$\begin{aligned}
& 8 \cdot \left[\Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} \right] + \Delta_k \omega_{s+j}(s, 0, 0) \\
& + \sum_{q=1}^p 4^{2q-1} \cdot \left[8 \cdot \left[\Delta_k \Gamma_{s+j-2q}^{\begin{bmatrix} s-q \\ s-(q-1) \\ s-1 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-2q-1}^{\begin{bmatrix} s-q \\ s-q \\ s-1 \end{bmatrix}} \right] + \Delta_k \omega_{s+j-(2q-1)}(s-q, 1, q-1) \right] \\
& + \sum_{q=1}^p 4^{2q} \cdot \left[8 \cdot \left[\Delta_k \Gamma_{s+j-(2q+1)}^{\begin{bmatrix} s-q \\ s-q \\ s-1 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-(2q+2)}^{\begin{bmatrix} s-q-1 \\ s-q \\ s-1 \end{bmatrix}} \right] + \Delta_k \omega_{s+j-2q}(s-q, 0, q) \right] \\
& = 8 \cdot \left[\Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} \right] + \Delta_k \omega_{s+j}(s, 0, 0) + \sum_{q=1}^p 2 \cdot 4^{2q} \cdot \Delta_k \Gamma_{s+j-2q}^{\begin{bmatrix} s-q \\ s-(q-1) \\ s-1 \end{bmatrix}} + \sum_{q=1}^p 2 \cdot 4^{2q+1} \cdot \Delta_k \Gamma_{s+j-(2q+1)}^{\begin{bmatrix} s-q \\ s-q \\ s-1 \end{bmatrix}} \\
& - \sum_{q=1}^p 2 \cdot 4^{2q+1} \cdot \Delta_k \Gamma_{s+j-(2q+1)}^{\begin{bmatrix} s-q \\ s-q \\ s-1 \end{bmatrix}} - \sum_{q=1}^p 2 \cdot 4^{2q+2} \cdot \Delta_k \Gamma_{s+j-(2q+2)}^{\begin{bmatrix} s-q-1 \\ s-q \\ s-1 \end{bmatrix}} \\
& + \sum_{q=1}^p 4^{2q-1} \cdot \Delta_k \omega_{s+j-(2q-1)}(s-q, 1, q-1) + \sum_{q=1}^p 4^{2q} \cdot \Delta_k \omega_{s+j-2q}(s-q, 0, q) \\
& = 8 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} + \Delta_k \omega_{s+j}(s, 0, 0) + \sum_{q=1}^p 2 \cdot 4^{2q} \cdot \Delta_k \Gamma_{s+j-2q}^{\begin{bmatrix} s-q \\ s-(q-1) \\ s-1 \end{bmatrix}} - \sum_{q=0}^p 2 \cdot 4^{2q+2} \cdot \Delta_k \Gamma_{s+j-(2q+2)}^{\begin{bmatrix} s-q-1 \\ s-q \\ s-1 \end{bmatrix}} \\
& + \sum_{q=1}^p 4^{2q-1} \cdot \Delta_k \omega_{s+j-(2q-1)}(s-q, 1, q-1) + \sum_{q=1}^p 4^{2q} \cdot \Delta_k \omega_{s+j-2q}(s-q, 0, q) \\
& = 8 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} + \Delta_k \omega_{s+j}(s, 0, 0) + \sum_{q=1}^p 2 \cdot 4^{2q} \cdot \Delta_k \Gamma_{s+j-2q}^{\begin{bmatrix} s-q \\ s-(q-1) \\ s-1 \end{bmatrix}} - \sum_{q=1}^{p+1} 2 \cdot 4^{2q} \cdot \Delta_k \Gamma_{s+j-2q}^{\begin{bmatrix} s-q \\ s-(q-1) \\ s-1 \end{bmatrix}} \\
& + \sum_{q=1}^p 4^{2q-1} \cdot \Delta_k \omega_{s+j-(2q-1)}(s-q, 1, q-1) + \sum_{q=1}^p 4^{2q} \cdot \Delta_k \omega_{s+j-2q}(s-q, 0, q) \\
& = 8 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} + \Delta_k \omega_{s+j}(s, 0, 0) - 2 \cdot 4^{2(p+1)} \cdot \Delta_k \Gamma_{s+j-2(p+1)}^{\begin{bmatrix} s-(p+1) \\ s-p \\ s-1 \end{bmatrix}} \\
& + \sum_{q=1}^p 4^{2q-1} \cdot \Delta_k \omega_{s+j-(2q-1)}(s-q, 1, q-1) + \sum_{q=1}^p 4^{2q} \cdot \Delta_k \omega_{s+j-2q}(s-q, 0, q)
\end{aligned}$$

Therefore

(8.4)

$$\begin{aligned}
& \Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 4^{2p+1} \cdot \Delta_k \Gamma_{s+j-2p-1}^{\begin{bmatrix} s-p-1 \\ s-p \\ s \end{bmatrix}} \\
& = 8 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} + \Delta_k \omega_{s+j}(s, 0, 0) - 2 \cdot 4^{2(p+1)} \cdot \Delta_k \Gamma_{s+j-2(p+1)}^{\begin{bmatrix} s-(p+1) \\ s-p \\ s-1 \end{bmatrix}} \\
& + \sum_{q=1}^p 4^{2q-1} \cdot \Delta_k \omega_{s+j-(2q-1)}(s-q, 1, q-1) + \sum_{q=1}^p 4^{2q} \cdot \Delta_k \omega_{s+j-2q}(s-q, 0, q) \\
& \iff
\end{aligned}$$

(8.5)

$$\begin{aligned}
& \Delta_k \Gamma_{s+j} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{s+j-1} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} \\
&= 4^{2p+1} \cdot \Delta_k \Gamma_{s+j-2p-1} \begin{bmatrix} s-p-1 \\ s-p \\ s-1 \end{bmatrix} - 2 \cdot 4^{2(p+1)} \cdot \Delta_k \Gamma_{s+j-2(p+1)} \begin{bmatrix} s-(p+1) \\ s-p \\ s-1 \end{bmatrix} \\
&+ \Delta_k \omega_{s+j}(s, 0, 0) + \sum_{q=1}^p 4^{2q-1} \cdot \Delta_k \omega_{s-q+(j-q+1)}(s-q, 1, q-1) + \sum_{q=1}^p 4^{2q} \cdot \Delta_k \omega_{s-q+(j-q)}(s-q, 0, q)
\end{aligned}$$

Setting $p = s-2$ in (8.5) we obtain the formula (8.1), that is

$$\begin{aligned}
& \Delta_k \Gamma_{s+j} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{s+j-1} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} \\
&= 4^{2s-3} \cdot \left[\Delta_k \Gamma_{j-s+3} \begin{bmatrix} \frac{1}{2} \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{j-s+2} \begin{bmatrix} \frac{1}{2} \\ s-1 \\ s \end{bmatrix} \right] \\
&+ \Delta_k \omega_{s+j}(s, 0, 0) + \sum_{q=1}^{s-2} 4^{2q-1} \cdot \Delta_k \omega_{s-q+(j-q+1)}(s-q, 1, q-1) + \sum_{q=1}^{s-2} 4^{2q} \cdot \Delta_k \omega_{s-q+(j-q)}(s-q, 0, q)
\end{aligned}$$

The case $i = 2s+j$, $0 \leq j \leq s$

Proceeding as in the first case we obtain

(8.6)

$$\begin{aligned}
& \Delta_k \Gamma_{2s+j} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{2s+j-1} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} \\
&= 4^{2p+1} \cdot \Delta_k \Gamma_{2s+j-2p-1} \begin{bmatrix} s-p-1 \\ s-p \\ s-1 \end{bmatrix} - 2 \cdot 4^{2(p+1)} \cdot \Delta_k \Gamma_{2s+j-2(p+1)} \begin{bmatrix} s-(p+1) \\ s-p \\ s-1 \end{bmatrix} \\
&+ \Delta_k \omega_{2s+j}(s, 0, 0) + \sum_{q=1}^p 4^{2q-1} \cdot \Delta_k \omega_{2(s-q)+(j+1)}(s-q, 1, q-1) + \sum_{q=1}^p 4^{2q} \cdot \Delta_k \omega_{2(s-q)+j}(s-q, 0, q)
\end{aligned}$$

Setting $p = s-2$ in (8.6) we get the formula (8.2), that is :

$$\begin{aligned}
& \Delta_k \Gamma_{2s+j} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{2s+j-1} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} \\
&= 4^{2s-3} \cdot \left[\Delta_k \Gamma_{j+3} \begin{bmatrix} \frac{1}{2} \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{j+2} \begin{bmatrix} \frac{1}{2} \\ s-1 \\ s \end{bmatrix} \right] \\
&+ \Delta_k \omega_{2s+j}(s, 0, 0) + \sum_{q=1}^{s-2} 4^{2q-1} \cdot \Delta_k \omega_{2(s-q)+(j+1)}(s-q, 1, q-1) + \sum_{q=1}^{s-2} 4^{2q} \cdot \Delta_k \omega_{2(s-q)+j}(s-q, 0, q)
\end{aligned}$$

□

8.4. An explicit formula for the difference

$$\Delta_k \Gamma_{s+j} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{s+j-1} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} \quad \text{for } 0 \leq j \leq 2s.$$

Lemma 8.3. *We have for $s \geq 5$*

$$(8.7) \quad \Delta_k \Gamma_{j-s+3} \begin{bmatrix} \frac{1}{2} \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{j-s+2} \begin{bmatrix} \frac{1}{2} \\ s-1 \\ s \end{bmatrix} = \begin{cases} 0 & \text{if } 0 \leq j \leq s-3 \\ 2^k & \text{if } j = s-2 \\ 9 \cdot 2^k & \text{if } j = s-1 \end{cases}$$

$$(8.8) \quad \Delta_k \Gamma_{j+3}^{\begin{bmatrix} 1 \\ 2 \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{j+2}^{\begin{bmatrix} \frac{1}{2} \\ s-1 \end{bmatrix}} = \begin{cases} 3 \cdot 2^{2k+1} - 17 \cdot 2^{k+1} & \text{if } j = 0 \\ -15 \cdot 2^{2k+1} + 3 \cdot 2^{k+3} & \text{if } j = 1 \\ -9 \cdot 2^{2k+2j-1} & \text{if } 2 \leq j \leq s-3 \\ -33 \cdot 2^{2k+2s-5} & \text{if } j = s-2 \\ -159 \cdot 2^{2k+2s-3} & \text{if } j = s-1 \\ -21 \cdot 2^{3k+s} + 21 \cdot 2^{2k+2s} & \text{if } j = s \end{cases}$$

Proof. From Lemma 1.22 we deduce easily Lemma 8.3. $\square$

Lemma 8.4. *We have :*

$$(8.9) \quad \Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} = \begin{cases} 3 \cdot 2^{k+s-1} & \text{if } j = 0 \\ 15 \cdot 2^{k+s-1} & \text{if } j = 1 \\ 63 \cdot 2^{k+s+3j-6} & \text{if } 2 \leq j \leq s-1 \end{cases}$$

$$(8.10) \quad \Delta_k \Gamma_{2s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} = \begin{cases} -15 \cdot 2^{2k+2s-2} - 33 \cdot 2^{k+4s-6} & \text{if } j = 0 \\ -321 \cdot 2^{2k+2s-2} + 3 \cdot 2^{k+4s-3} & \text{if } j = 1 \\ -315 \cdot 2^{2k+2s+4j-6} & \text{if } 2 \leq j \leq s-1 \\ -21 \cdot 2^{3k+3s-3} + 21 \cdot 2^{2k+6s-6} & \text{if } j = s \end{cases}$$

Proof. proof of (8.9)

From the reduction formula (8.1) using Lemma 7.4 and the formula (8.7) we deduce

The case $j = 0$

$$\begin{aligned} & \Delta_k \Gamma_s^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} \\ &= 2^{4s-6} \cdot \left[\Delta_k \Gamma_{-s+3}^{\begin{bmatrix} 1 \\ 2 \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{-s+2}^{\begin{bmatrix} \frac{1}{2} \\ s-1 \end{bmatrix}} \right] \\ &+ \Delta_k \omega_s(s, 0, 0) + \sum_{q=1}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{s-q+(-q+1)}(s-q, 1, q-1) + \sum_{q=1}^{s-2} 2^{4q} \cdot \Delta_k \omega_{s-q+(-q)}(s-q, 0, q) \\ &= \Delta_k \omega_s(s, 0, 0) + 2^2 \cdot \Delta_k \omega_{s-1}(s-1, 1, 0) = 2^{k+s-1} + 2^2 \cdot 2^{k+s-2} = 3 \cdot 2^{k+s-1} \end{aligned}$$

The case $j = 1$

$$\begin{aligned} & \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_s^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} \\ &= 2^{4s-6} \cdot \left[\Delta_k \Gamma_{-s+4}^{\begin{bmatrix} 1 \\ 2 \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{-s+3}^{\begin{bmatrix} \frac{1}{2} \\ s-1 \end{bmatrix}} \right] \\ &+ \Delta_k \omega_{s+1}(s, 0, 0) + \sum_{q=1}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{s-q+(-q+2)}(s-q, 1, q-1) + \sum_{q=1}^{s-2} 2^{4q} \cdot \Delta_k \omega_{s-q+(-q+1)}(s-q, 0, q) \\ &= \Delta_k \omega_{s+1}(s, 0, 0) + \sum_{q=1}^2 2^{4q-2} \cdot \Delta_k \omega_{s-q+(-q+2)}(s-q, 1, q-1) + \sum_{q=1}^1 2^{4q} \cdot \Delta_k \omega_{s-q+(-q+1)}(s-q, 0, q) \\ &= \Delta_k \omega_{s+1}(s, 0, 0) + 2^2 \cdot \Delta_k \omega_{s-1+1}(s-1, 1, 0) + 2^6 \cdot \Delta_k \omega_{s-2}(s-2, 1, 1) + 2^4 \cdot \Delta_k \omega_{s-1}(s-1, 0, 1) \\ &= -3 \cdot 2^{k+s-1} + 2^2 \cdot (-3 \cdot 2^{k+s-2}) + 2^6 \cdot 2^{k+s-3} + 2^4 \cdot 2^{k+s-2} = 15 \cdot 2^{k+s-1} \end{aligned}$$

The case $j = 2$

$$\begin{aligned}
& \Delta_k \Gamma_{s+2} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{s+1} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} \\
&= 2^{4s-6} \cdot \left[\Delta_k \Gamma_{-s+5} \begin{bmatrix} \frac{1}{2} \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{-s+4} \begin{bmatrix} \frac{1}{2} \\ s-1 \end{bmatrix} \right] \\
&+ \Delta_k \omega_{s+2}(s, 0, 0) + \sum_{q=1}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{s-q+(-q+3)}(s-q, 1, q-1) + \sum_{q=1}^{s-2} 2^{4q} \cdot \Delta_k \omega_{s-q+(-q+2)}(s-q, 0, q) \\
&= \Delta_k \omega_{s+2}(s, 0, 0) + \sum_{q=1}^3 2^{4q-2} \cdot \Delta_k \omega_{s-q+(-q+3)}(s-q, 1, q-1) + \sum_{q=1}^2 2^{4q} \cdot \Delta_k \omega_{s-q+(-q+2)}(s-q, 0, q) \\
&= \Delta_k \omega_{s+2}(s, 0, 0) + 2^2 \cdot \Delta_k \omega_{s-1+2}(s-1, 1, 0) + 2^6 \cdot \Delta_k \omega_{s-2+1}(s-2, 1, 1) + 2^{10} \cdot \Delta_k \omega_{s-3}(s-3, 1, 2) \\
&+ 2^4 \cdot \Delta_k \omega_{s-1+1}(s-1, 0, 1) + 2^8 \cdot \Delta_k \omega_{s-2}(s-2, 0, 2) \\
&= 2^{k+s} + 2^2 \cdot 2^{k+s-1} + 2^6 \cdot (-3 \cdot 2^{k+s-3}) + 2^{10} \cdot 2^{k+s-4} + 2^4 \cdot (-3 \cdot 2^{k+s-2}) + 2^8 \cdot 2^{k+s-3} = 63 \cdot 2^{k+s}
\end{aligned}$$

The case $3 \leq j \leq s-3$

$$\begin{aligned}
& \Delta_k \omega_{s-q+(j-q+1)}(s-q, 1, q-1) = 0 \\
& \iff s-q+3 \leq s-q+(j-q+1) \leq 1+2 \cdot (s-q) - 1 \iff q \leq j-2 \\
& \Delta_k \omega_{s-q+(j-q)}(s-q, 0, q) = 0 \\
& \iff s-q+3 \leq s-q+(j-q) \leq 2 \cdot (s-q) - 1 \iff q \leq j-3
\end{aligned}$$

$$\begin{aligned}
& \Delta_k \Gamma_{s+j} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{s+j-1} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} \\
&= 4^{2s-3} \cdot \left[\Delta_k \Gamma_{j-s+3} \begin{bmatrix} \frac{1}{2} \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{j-s+2} \begin{bmatrix} \frac{1}{2} \\ s-1 \end{bmatrix} \right] \\
&+ \Delta_k \omega_{s+j}(s, 0, 0) + \sum_{q=1}^{s-2} 4^{2q-1} \cdot \Delta_k \omega_{s-q+(j-q+1)}(s-q, 1, q-1) + \sum_{q=1}^{s-2} 4^{2q} \cdot \Delta_k \omega_{s-q+(j-q)}(s-q, 0, q) \\
&= \Delta_k \omega_{s+j}(s, 0, 0) + \sum_{q=1}^{j+1} 4^{2q-1} \cdot \Delta_k \omega_{s-q+(j-q+1)}(s-q, 1, q-1) + \sum_{q=1}^j 4^{2q} \cdot \Delta_k \omega_{s-q+(j-q)}(s-q, 0, q) \\
&= \Delta_k \omega_{s+j}(s, 0, 0) + \sum_{q=j-1}^{j+1} 4^{2q-1} \cdot \Delta_k \omega_{s-q+(j-q+1)}(s-q, 1, q-1) + \sum_{q=j-2}^j 4^{2q} \cdot \Delta_k \omega_{s-q+(j-q)}(s-q, 0, q) \\
&= [2^{4(j-1)-2} \cdot 2^{k+s-(j-1)} + 2^{4j-2} \cdot (-3 \cdot 2^{k+(s-j)-1}) + 2^{4j+2} \cdot 2^{k+s-(j+1)-1}] \\
&+ [2^{4(j-2)} \cdot 2^{k+s-(j-2)} + 2^{4(j-1)} \cdot (-3 \cdot 2^{k+s-(j-1)-1}) + 2^{4j} \cdot 2^{k+(s-j)-1}] = 63 \cdot 2^{k+s+3j-6}
\end{aligned}$$

The case $j = s-2$

$$\begin{aligned}
& \Delta_k \omega_{s-q+(s-q-1)}(s-q, 1, q-1) = 0 \\
& \iff s-q+3 \leq s-q+(s-q-1) \leq 1+2 \cdot (s-q) - 1 \iff q \leq s-4 \\
& \Delta_k \omega_{s-q+(s-q-2)}(s-q, 0, q) = 0 \\
& \iff s-q+3 \leq s-q+(s-q-2) \leq 2 \cdot (s-q) - 1 \iff q \leq s-5
\end{aligned}$$

$$\begin{aligned}
& \Delta_k \Gamma_{2s-2} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{2s-3} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} \\
&= 2^{4s-6} \cdot \left[\Delta_k \Gamma_1 \begin{bmatrix} \frac{1}{2} \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_0 \begin{bmatrix} \frac{1}{2} \\ s-1 \end{bmatrix} \right] \\
&+ \Delta_k \omega_{2s-2}(s, 0, 0) + \sum_{q=1}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{s-q+(s-q-1)}(s-q, 1, q-1) + \sum_{q=1}^{s-2} 2^{4q} \cdot \Delta_k \omega_{s-q+(s-q-2)}(s-q, 0, q) \\
&= 2^{4s-6} \cdot 2^k + \Delta_k \omega_{2s-2}(s, 0, 0) + \sum_{q=s-3}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{s-q+(s-q-1)}(s-q, 1, q-1) + \sum_{q=s-4}^{s-2} 2^{4q} \cdot \Delta_k \omega_{s-q+(s-q-2)}(s-q, 0, q) \\
&= 2^{k+4s-6} + [2^{4(s-3)-2} \cdot 2^{k+s-(s-3)} + 2^{4(s-2)-2} \cdot (-3 \cdot 2^{k+s-(s-2)-1})] \\
&+ [2^{4(s-4)} \cdot 2^{k+s-(s-4)} + 2^{4(s-3)} \cdot (-3 \cdot 2^{k+s-(s-3)-1}) + 2^{4(s-2)} \cdot 2^{k+s-(s-2)-1}] \\
&= 63 \cdot 2^{k+4s-12}
\end{aligned}$$

The case $j = s-1$

$$\begin{aligned}
& \Delta_k \omega_{s-q+(s-q)}(s-q, 1, q-1) = 0 \\
& \iff s-q+3 \leq s-q+(s-q) \leq 1+2 \cdot (s-q)-1 \iff q \leq s-3 \\
& \Delta_k \omega_{s-q+(s-q-1)}(s-q, 0, q) = 0 \\
& \iff s-q+3 \leq s-q+(s-q-1) \leq 2 \cdot (s-q)-1 \iff q \leq s-4
\end{aligned}$$

$$\begin{aligned}
& \Delta_k \Gamma_{2s-1} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{2s-2} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} \\
&= 2^{4s-6} \cdot \left[\Delta_k \Gamma_2 \begin{bmatrix} \frac{1}{2} \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_1 \begin{bmatrix} \frac{1}{2} \\ s-1 \end{bmatrix} \right] \\
&+ \Delta_k \omega_{2s-1}(s, 0, 0) + \sum_{q=1}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{s-q+(s-q)}(s-q, 1, q-1) + \sum_{q=1}^{s-2} 2^{4q} \cdot \Delta_k \omega_{s-q+(s-q-1)}(s-q, 0, q) \\
&= 2^{4s-6} \cdot \left[\Delta_k \Gamma_2 \begin{bmatrix} \frac{1}{2} \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_1 \begin{bmatrix} \frac{1}{2} \\ s-1 \end{bmatrix} \right] + \Delta_k \omega_{2s-1}(s, 0, 0) + \sum_{q=s-2}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{s-q+(s-q)}(s-q, 1, q-1) \\
&+ \sum_{q=s-3}^{s-2} 2^{4q} \cdot \Delta_k \omega_{s-q+(s-q-1)}(s-q, 0, q) \\
&= 2^{k+4s-6} \cdot 9 \cdot 2^k + 2^{4(s-2)-2} \cdot 2^{k+s-(s-2)} + 2^{4(s-3)} \cdot 2^{k+3} + 2^{4(s-2)} \cdot (-3 \cdot 2^{k+1}) = 63 \cdot 2^{k+4s-9}
\end{aligned}$$

proof of (8.10)

From the reduction formula (8.2) using Lemma 7.4 and the formula (8.8) we deduce

The case $j = 0$

$$\begin{aligned}
& \Delta_k \Gamma_{2s} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{2s-1} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} \\
&= 4^{2s-3} \cdot \left[\Delta_k \Gamma_3 \begin{bmatrix} \frac{1}{2} \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_2 \begin{bmatrix} \frac{1}{2} \\ s-1 \end{bmatrix} \right] \\
&+ \Delta_k \omega_{2s}(s, 0, 0) + \sum_{q=1}^{s-2} 4^{2q-1} \cdot \Delta_k \omega_{2(s-q)+1}(s-q, 1, q-1) + \sum_{q=1}^{s-2} 4^{2q} \cdot \Delta_k \omega_{2(s-q)}(s-q, 0, q)
\end{aligned}$$

$$\begin{aligned}
&= 2^{4s-6} \cdot [3 \cdot 2^{2k+1} - 17 \cdot 2^{k+1}] - 9 \cdot 2^{2k+2s-2} + 2^2 \cdot [-9 \cdot 2^{2k+2s-3}] \\
&+ \sum_{q=2}^{s-2} 2^{4q-2} \cdot [-3 \cdot 2^{2k+2(s-q)+1-2}] + \sum_{q=1}^{s-3} 2^{4q} \cdot [-3 \cdot 2^{2k+2(s-q)-2}] + 2^{4s-8} \cdot [-3 \cdot 2^{2k+2} + 2^{k+2}] \\
&= -15 \cdot 2^{2k+2s-2} - 33 \cdot 2^{k+4s-6}
\end{aligned}$$

The case $j = 1$

We have

$$\Delta_k \omega_{2s+1}(s, 0, 0) = 9 \cdot 2^{2k+2s-2}$$

$$\Delta_k \omega_{2(s-1)+2}(s-1, 1, 0) = 9 \cdot 2^{2k+2s-3}$$

$$\Delta_k \omega_{2(s-2)+2}(s-2, 1, 1) = -9 \cdot 2^{2k+2s-5}$$

$$\Delta_k \omega_{2(s-q)+2}(s-q, 1, q-1) = 15 \cdot 2^{2k+2(s-q)-1} \quad \text{if } 3 \leq q \leq s-2$$

$$\Delta_k \omega_{2(s-1)+1}(s-1, 0, 1) = -9 \cdot 2^{2k+2s-4}$$

$$\Delta_k \omega_{2(s-q)+1}(s-q, 0, q) = 15 \cdot 2^{2k+2(s-q)-2} \quad \text{if } 2 \leq q \leq s-2$$

$$\begin{aligned}
&\Delta_k \Gamma_{2s+1} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{2s} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} = 2^{4s-6} \cdot \left[\Delta_k \Gamma_4 \begin{bmatrix} \frac{1}{2} \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_3 \begin{bmatrix} \frac{1}{2} \\ s-1 \end{bmatrix} \right] \\
&+ \Delta_k \omega_{2s+1}(s, 0, 0) + \sum_{q=1}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{2(s-q)+2}(s-q, 1, q-1) + \sum_{q=1}^{s-2} 2^{4q} \cdot \Delta_k \omega_{2(s-q)+1}(s-q, 0, q) \\
&= 2^{4s-6} \cdot (3 \cdot 2^{2k+1} - 17 \cdot 2^{k+1}) + 9 \cdot 2^{2k+2s-2} + 2^2 \cdot 9 \cdot 2^{2k+2s-3} + 2^6 \cdot (-9 \cdot 2^{2k+2s-5}) \\
&+ \sum_{q=3}^{s-2} 2^{4q-2} \cdot 15 \cdot 2^{2k+2(s-q)-1} + 2^4 \cdot (-9 \cdot 2^{2k+2s-4}) + \sum_{q=2}^{s-2} 2^{4q} \cdot 15 \cdot 2^{2k+2(s-q)-2} \\
&= -15 \cdot 2^{2k+4s-5} + 3 \cdot 2^{k+4s-3} - 81 \cdot 2^{2k+2s-2} + 15 \cdot 2^{2k+4s-5} - 5 \cdot 2^{2k+2s+3} - 5 \cdot 2^{2k+2s+2} \\
&= -321 \cdot 2^{2k+2s-2} + 3 \cdot 2^{k+4s-3}
\end{aligned}$$

The case $2 \leq j \leq s-4$

$$\begin{aligned}
&\Delta_k \Gamma_{2s+j} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{2s+j-1} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} \\
&= 2^{4s-6} \cdot \left[\Delta_k \Gamma_{j+3} \begin{bmatrix} \frac{1}{2} \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{j+2} \begin{bmatrix} \frac{1}{2} \\ s-1 \end{bmatrix} \right] \\
&+ \Delta_k \omega_{2s+j}(s, 0, 0) + \sum_{q=1}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{2(s-q)+(j+1)}(s-q, 1, q-1) + \sum_{q=1}^{s-2} 2^{4q} \cdot \Delta_k \omega_{2(s-q)+j}(s-q, 0, q)
\end{aligned}$$

We have

$$\Delta_k \omega_{2s+j}(s, 0, 0) = 0 \quad \text{if } 2 \leq j \leq s-4,$$

$$\Delta_k \omega_{2(s-q)+(j+1)}(s-q, 1, q-1) = \begin{cases} 0 & \text{if } 1 \leq q \leq j-1 \\ 3 \cdot 2^{2k+2s-3} & \text{if } q = j \\ -15 \cdot 2^{2k+2s-5} & \text{if } q = j+1 \\ 9 \cdot 2^{2k+2s-2q+2j-3} & \text{if } q \geq j+2, \end{cases}$$

$$\Delta_k \omega_{2(s-q)+j}(s-q, 0, q) = \begin{cases} 0 & \text{if } 1 \leq q \leq j-2 \\ 3 \cdot 2^{2k+2s-2} & \text{if } q = j-1 \\ -15 \cdot 2^{2k+2s-4} & \text{if } q = j \\ 9 \cdot 2^{2k+2s-2q+2j-4} & \text{if } q \geq j+1. \end{cases}$$

$$\begin{aligned} \sum_{q=1}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{2(s-q)+(j+1)}(s-q, 1, q-1) &= \sum_{q=j}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{2(s-q)+(j+1)}(s-q, 1, q-1) \\ &= 2^{4j-2} \cdot (3 \cdot 2^{2k+2s-3}) + 2^{4(j+1)-2} \cdot (-15 \cdot 2^{2k+2s-5}) + \sum_{q=j+2}^{s-2} 2^{4q-2} \cdot 9 \cdot 2^{2k+2s-2q+2j-3} \\ &= 3 \cdot 2^{2k+2s+4j-5} - 15 \cdot 2^{2k+2s+4j-3} + 9 \cdot 2^{2k+2s+2j-5} \sum_{q=j+2}^{s-2} 2^{2q} = 3 \cdot 2^{2k+4s+2j-7} - 105 \cdot 2^{2k+2s+4j-5} \end{aligned}$$

and

$$\begin{aligned} \sum_{q=1}^{s-2} 2^{4q} \cdot \Delta_k \omega_{2(s-q)+j}(s-q, 0, q) &= \sum_{q=j-1}^{s-2} 2^{4q} \cdot \Delta_k \omega_{2(s-q)+j}(s-q, 0, q) \\ &= 2^{4(j-1)} \cdot (3 \cdot 2^{2k+2s-2}) + 2^{4j} \cdot (-15 \cdot 2^{2k+2s-4}) + \sum_{q=j+1}^{s-2} 2^{4q} \cdot 9 \cdot 2^{2k+2s-2q+2j-4} \\ &= 3 \cdot 2^{2k+2s+4j-6} - 15 \cdot 2^{2k+2s+4j-4} + 9 \cdot 2^{2k+2s+2j-4} \sum_{q=j+1}^{s-2} 2^{2q} = 3 \cdot 2^{2k+4s+2j-6} - 105 \cdot 2^{2k+2s+4j-6} \end{aligned}$$

Combining the above results we obtain for $2 \leq j \leq s-4$

$$\begin{aligned} &\Delta_k \Gamma_{2s+j} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{2s+j-1} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} \\ &= 2^{4s-6} \cdot \left[\Delta_k \Gamma_{j+3} \begin{bmatrix} \frac{1}{2} \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{j+2} \begin{bmatrix} \frac{1}{2} \\ s-1 \end{bmatrix} \right] \\ &+ \Delta_k \omega_{2s+j}(s, 0, 0) + \sum_{q=1}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{2(s-q)+(j+1)}(s-q, 1, q-1) + \sum_{q=1}^{s-2} 2^{4q} \cdot \Delta_k \omega_{2(s-q)+j}(s-q, 0, q) \\ &= -9 \cdot 2^{2k+4s+2j-7} + (3 \cdot 2^{2k+4s+2j-7} - 105 \cdot 2^{2k+2s+4j-5}) + (3 \cdot 2^{2k+4s+2j-6} - 105 \cdot 2^{2k+2s+4j-6}) \\ &= -315 \cdot 2^{2k+2s+4j-6} \end{aligned}$$

The case $j = s-3$

$$\begin{aligned} &\Delta_k \Gamma_{2s+(s-3)} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{2s+(s-4)} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} \\ &= 2^{4s-6} \cdot \left[\Delta_k \Gamma_s \begin{bmatrix} \frac{1}{2} \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{s-1} \begin{bmatrix} \frac{1}{2} \\ s-1 \end{bmatrix} \right] \end{aligned}$$

$$+ \Delta_k \omega_{2s+(s-3)}(s, 0, 0) + \sum_{q=1}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{2(s-q)+(s-2)}(s-q, 1, q-1) + \sum_{q=1}^{s-2} 2^{4q} \cdot \Delta_k \omega_{2(s-q)+(s-3)}(s-q, 0, q)$$

We have

$$\Delta_k \omega_{2s+(s-3)}(s, 0, 0) = 0 \quad \text{if } j = s-3,$$

$$\Delta_k \omega_{2(s-q)+(s-2)}(s-q, 1, q-1) = \begin{cases} 0 & \text{if } 1 \leq q \leq s-4 \\ 3 \cdot 2^{2k+2s-3} & \text{if } q = s-3 \\ -15 \cdot 2^{2k+2s-5} & \text{if } q = s-2, \end{cases}$$

$$\Delta_k \omega_{2(s-q)+(s-3)}(s-q, 0, q) = \begin{cases} 0 & \text{if } 1 \leq q \leq s-5 \\ 3 \cdot 2^{2k+2s-2} & \text{if } q = s-4 \\ -15 \cdot 2^{2k+2s-4} & \text{if } q = s-3 \\ 9 \cdot 2^{2k+2s-6} & \text{if } q = s-2, \end{cases}$$

$$\begin{aligned} \sum_{q=1}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{2(s-q)+(s-2)}(s-q, 1, q-1) &= \sum_{q=s-3}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{2(s-q)+(s-2)}(s-q, 1, q-1) \\ &= 2^{4(s-3)-2} \cdot (3 \cdot 2^{2k+2s-3}) + 2^{4(s-2)-2} \cdot (-15 \cdot 2^{2k+2s-5}) = -57 \cdot 2^{2k+6s-17}, \end{aligned}$$

$$\begin{aligned} \sum_{q=1}^{s-2} 2^{4q} \cdot \Delta_k \omega_{2(s-q)+(s-3)}(s-q, 0, q) &= \sum_{q=s-4}^{s-2} 2^{4q} \cdot \Delta_k \omega_{2(s-q)+(s-3)}(s-q, 0, q) \\ &= 2^{4(s-4)} \cdot (3 \cdot 2^{2k+2s-2}) + 2^{4(s-3)} \cdot (-15 \cdot 2^{2k+2s-4}) + 2^{4(s-2)} \cdot (9 \cdot 2^{2k+2s-6}) = 87 \cdot 2^{2k+6s-18}. \end{aligned}$$

As before we deduce from the above results

$$\begin{aligned} \Delta_k \Gamma_{2s+(s-3)} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{2s+(s-4)} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} &= 2^{4s-6} \cdot \left[\Delta_k \Gamma_s \begin{bmatrix} \frac{1}{2} \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{s-1} \begin{bmatrix} \frac{1}{2} \\ \frac{1}{2} \end{bmatrix} \right] \\ &+ \Delta_k \omega_{2s+(s-3)}(s, 0, 0) + \sum_{q=1}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{2(s-q)+(s-2)}(s-q, 1, q-1) + \sum_{q=1}^{s-2} 2^{4q} \cdot \Delta_k \omega_{2(s-q)+(s-3)}(s-q, 0, q) \\ &= -9 \cdot 2^{2k+6s-13} - 57 \cdot 2^{2k+6s-17} + 87 \cdot 2^{2k+6s-18} = -315 \cdot 2^{2k+6s-18} \end{aligned}$$

The case $j = s-2$

$$\begin{aligned} \Delta_k \Gamma_{2s+(s-2)} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{2s+(s-3)} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} \\ &= 2^{4s-6} \cdot \left[\Delta_k \Gamma_{s+1} \begin{bmatrix} \frac{1}{2} \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_s \begin{bmatrix} \frac{1}{2} \\ s-1 \end{bmatrix} \right] \\ &+ \Delta_k \omega_{2s+(s-2)}(s, 0, 0) + \sum_{q=1}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{2(s-q)+(s-1)}(s-q, 1, q-1) + \sum_{q=1}^{s-2} 2^{4q} \cdot \Delta_k \omega_{2(s-q)+(s-2)}(s-q, 0, q) \end{aligned}$$

We have

$$\Delta_k \omega_{2s+(s-2)}(s, 0, 0) = 0 \quad \text{if } j = s-2,$$

$$\Delta_k \omega_{2(s-q)+(s-1)}(s-q, 1, q-1) = \begin{cases} 0 & \text{if } 1 \leq q \leq s-3 \\ 3 \cdot 2^{2k+2s-3} & \text{if } q = s-2, \end{cases}$$

$$\Delta_k \omega_{2(s-q)+(s-2)}(s-q, 0, q) = \begin{cases} 0 & \text{if } 1 \leq q \leq s-4 \\ 3 \cdot 2^{2k+2s-2} & \text{if } q = s-3 \\ -15 \cdot 2^{2k+2s-4} & \text{if } q = s-2, \end{cases}$$

$$\begin{aligned} \sum_{q=1}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{2(s-q)+(s-1)}(s-q, 1, q-1) &= \sum_{q=s-2}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{2(s-q)+(s-1)}(s-q, 1, q-1) \\ &= 2^{4(s-2)-2} \cdot (3 \cdot 2^{2k+2s-3}) = 3 \cdot 2^{2k+6s-13}, \end{aligned}$$

$$\begin{aligned} \sum_{q=1}^{s-2} 2^{4q} \cdot \Delta_k \omega_{2(s-q)+(s-2)}(s-q, 0, q) &= \sum_{q=s-3}^{s-2} 2^{4q} \cdot \Delta_k \omega_{2(s-q)+(s-2)}(s-q, 0, q) \\ &= 2^{4(s-3)} \cdot (3 \cdot 2^{2k+2s-2}) + 2^{4(s-2)} \cdot (-15 \cdot 2^{2k+2s-4}) = 3 \cdot 2^{2k+6s-14} - 15 \cdot 2^{2k+6s-12} = -57 \cdot 2^{2k+6s-14}. \end{aligned}$$

As before we deduce from the above results

$$\begin{aligned} &\Delta_k \Gamma_{2s+(s-2)} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{2s+(s-3)} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} \\ &= 2^{4s-6} \cdot \left[\Delta_k \Gamma_{s+1} \begin{bmatrix} \frac{1}{2} \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_s \begin{bmatrix} \frac{1}{2} \\ s-1 \end{bmatrix} \right] \\ &+ \Delta_k \omega_{2s+(s-2)}(s, 0, 0) + \sum_{q=1}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{2(s-q)+(s-1)}(s-q, 1, q-1) + \sum_{q=1}^{s-2} 2^{4q} \cdot \Delta_k \omega_{2(s-q)+(s-2)}(s-q, 0, q) \\ &= -33 \cdot 2^{2k+6s-11} + 3 \cdot 2^{2k+6s-13} - 57 \cdot 2^{2k+6s-14} \\ &= -315 \cdot 2^{2k+6s-14} \end{aligned}$$

The case $j = s$

$$\begin{aligned} &\Delta_k \Gamma_{2s+s} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{2s+(s-1)} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} \\ &= 2^{4s-6} \cdot \left[\Delta_k \Gamma_{s+3} \begin{bmatrix} \frac{1}{2} \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{s+2} \begin{bmatrix} \frac{1}{2} \\ s-1 \end{bmatrix} \right] \\ &+ \Delta_k \omega_{3s}(s, 0, 0) + \sum_{q=1}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{2(s-q)+s+1}(s-q, 1, q-1) + \sum_{q=1}^{s-2} 2^{4q} \cdot \Delta_k \omega_{2(s-q)+s}(s-q, 0, q) \end{aligned}$$

We have

$$\begin{aligned} \Delta_k \omega_{3s}(s, 0, 0) &= 21 \cdot 2^{3k+3s-3}, \\ \Delta_k \omega_{2(s-q)+s+1}(s-q, 1, q-1) &= 21 \cdot 2^{3k+3(s-q)+q-2} & \text{if } 1 \leq q \leq s-2, \\ \Delta_k \omega_{2(s-q)+s}(s-q, 0, q) &= 21 \cdot 2^{3k+3(s-q)+q-3} & \text{if } 1 \leq q \leq s-2, \end{aligned}$$

$$\sum_{q=1}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{2(s-q)+s+1}(s-q, 1, q-1) = \sum_{q=1}^{s-2} 2^{4q-2} \cdot 21 \cdot 2^{3k+3(s-q)+q-2}$$

$$\begin{aligned}
&= 21 \cdot 2^{3k+3s-4} \cdot \sum_{q=1}^{s-2} 2^{2q} = 21 \cdot 2^{3k+3s-2} \cdot \sum_{q=0}^{s-3} 2^{2q} = 7 \cdot 2^{3k+5s-6} - 7 \cdot 2^{3k+3s-2}, \\
&\sum_{q=1}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{2(s-q)+s}(s-q, 0, q) = \sum_{q=1}^{s-2} 2^{4q} \cdot 21 \cdot 2^{3k+3(s-q)+q-3} \\
&= 21 \cdot 2^{3k+3s-3} \cdot \sum_{q=1}^{s-2} 2^{2q} = 21 \cdot 2^{3k+3s-1} \cdot \sum_{q=0}^{s-3} 2^{2q} = 7 \cdot 2^{3k+5s-5} - 7 \cdot 2^{3k+3s-1}.
\end{aligned}$$

As before we deduce from the above results

$$\begin{aligned}
&\Delta_k \Gamma_{2s+s} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{2s+(s-1)} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} \\
&= 2^{4s-6} \cdot \left[\Delta_k \Gamma_{s+3} \begin{bmatrix} \frac{1}{2} \\ s \\ s \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{s+2} \begin{bmatrix} \frac{1}{2} \\ s-1 \\ s \end{bmatrix} \right] \\
&+ \Delta_k \omega_{3s}(s, 0, 0) + \sum_{q=1}^{s-2} 2^{4q-2} \cdot \Delta_k \omega_{2(s-q)+s+1}(s-q, 1, q-1) + \sum_{q=1}^{s-2} 2^{4q} \cdot \Delta_k \omega_{2(s-q)+s}(s-q, 0, q) \\
&= -21 \cdot 2^{3k+5s-6} + 21 \cdot 2^{2k+6s-6} + 21 \cdot 2^{3k+3s-3} + 7 \cdot 2^{3k+5s-6} - 7 \cdot 2^{3k+3s-2} + 7 \cdot 2^{3k+5s-5} - 7 \cdot 2^{3k+3s-1} \\
&= 21 \cdot 2^{2k+6s-6} - 21 \cdot 2^{3k+3s-3}
\end{aligned}$$

□

9. An explicit formula for $\Delta_k \Gamma_i \begin{bmatrix} s \\ s \\ s \end{bmatrix}$, for $0 \leq i \leq 3s$

9.1. Notation.

Definition 9.1. We recall the formula obtained in Section 8 (see (8.3)):

$$\Delta_k \omega_i(s, m, l) = \Delta_k \Gamma_i \begin{bmatrix} s+m \\ s+m+l \end{bmatrix} - 4 \cdot \Delta_k \Gamma_{i-1} \begin{bmatrix} s+m-1 \\ s+m+l \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{i-1} \begin{bmatrix} s+m \\ s+m+l-1 \end{bmatrix} + 32 \cdot \Delta_k \Gamma_{i-2} \begin{bmatrix} s+m-1 \\ s+m+l-1 \end{bmatrix}$$

9.2. Introduction. Using the results concerning $\Delta_k \omega_i(s, m, l)$ obtained in Lemma 7.4 we adapt the method of Section 12 in [2] to compute $\Delta_k \Gamma_i \begin{bmatrix} s \\ s \\ s \end{bmatrix}$ for $0 \leq i \leq 3s$ applying successively the formula 8.3.

$$\text{9.3. Computation of } \Delta_k \Gamma_{s+j} \begin{bmatrix} s+m \\ s+m+l \end{bmatrix} = \Gamma_{s+j} \begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times (k+1) - \Gamma_{s+j} \begin{bmatrix} s+m \\ s+m+l \end{bmatrix} \times k \quad \text{for } j=0,1,2.$$

Lemma 9.2. We have

$$(9.1) \quad \Delta_k \Gamma_s \begin{bmatrix} s+m \\ s+m+l \end{bmatrix} = \begin{cases} 2^{k+s-1} & \text{if } m \geq 1, l \geq 0 \\ 3 \cdot 2^{k+s-1} & \text{if } m = 0, l \geq 1 \\ 7 \cdot 2^{k+s-1} & \text{if } m = 0, l = 0, \end{cases}$$

$$(9.2) \quad \Delta_k \Gamma_{s+1} \begin{bmatrix} s+m \\ s+m+l \end{bmatrix} = \begin{cases} 9 \cdot 2^{k+s-1} & \text{if } m \geq 2, l \geq 0 \\ 17 \cdot 2^{k+s-1} & \text{if } m = 1, l \geq 1 \\ 33 \cdot 2^{k+s-1} & \text{if } m = 1, l = 0 \\ 147 \cdot 2^{k+s-1} & \text{if } m = 0, l = 0 \\ 71 \cdot 2^{k+s-1} & \text{if } m = 0, l = 1 \\ 39 \cdot 2^{k+s-1} & \text{if } m = 0, l \geq 2, \end{cases}$$

$$(9.3) \quad \Delta_k \Gamma_{s+2}^{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix}} = \begin{cases} 78 \cdot 2^{k+s-1} & \text{if } m \geq 3, l \geq 0 \\ 110 \cdot 2^{k+s-1} & \text{if } m = 2, l \geq 1 \\ 174 \cdot 2^{k+s-1} & \text{if } m = 2, l = 0 \\ 198 \cdot 2^{k+s-1} & \text{if } m = 1, l \geq 2 \\ 326 \cdot 2^{k+s-1} & \text{if } m = 1, l = 1 \\ 630 \cdot 2^{k+s-1} & \text{if } m = 1, l = 0 \\ 438 \cdot 2^{k+s-1} & \text{if } m = 0, l \geq 3 \\ 694 \cdot 2^{k+s-1} & \text{if } m = 0, l = 2 \\ 1302 \cdot 2^{k+s-1} & \text{if } m = 0, l = 1 \\ 2646 \cdot 2^{k+s-1} & \text{if } m = 0, l = 0. \end{cases}$$

Proof. The proof is based on Lemma 7.4 and (6.33).

We remark that to obtain the result in one case we are using some of the results in the precedent cases.

proof of (9.1)

The case $m \geq 1, l \geq 0$

$$\Delta_k \Gamma_s^{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s+m-1 \\ s+m+l \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s+m \\ s+m+l-1 \end{bmatrix}} + 32 \cdot \Delta_k \Gamma_{s-2}^{\begin{bmatrix} s+m-1 \\ s+m+l-1 \end{bmatrix}} = \Delta_k \omega_s(s, m, l) = 2^{k+s-1}$$

$$\text{since } \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s+m-1 \\ s+m+l \end{bmatrix}} = \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s+m \\ s+m+l-1 \end{bmatrix}} = \Delta_k \Gamma_{s-2}^{\begin{bmatrix} s+m-1 \\ s+m+l-1 \end{bmatrix}} = 0 \quad \text{we get } \Delta_k \Gamma_s^{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix}} = 2^{k+s-1}$$

The case $m = 0, l \geq 1$

$$\Delta_k \Gamma_s^{\begin{bmatrix} s \\ s+l \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s-1 \\ s+l \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s \\ s+l-1 \end{bmatrix}} + 32 \cdot \Delta_k \Gamma_{s-2}^{\begin{bmatrix} s-1 \\ s+l-1 \end{bmatrix}} = \Delta_k \omega_s(s, 0, l) = 2^{k+s-1}$$

$$\text{since } \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s \\ s+l-1 \end{bmatrix}} = \Delta_k \Gamma_{s-2}^{\begin{bmatrix} s-1 \\ s+l-1 \end{bmatrix}} = 0 \text{ and } \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s-1 \\ s+l \end{bmatrix}} = 2^{k+s-2} \quad \text{we get } \Delta_k \Gamma_s^{\begin{bmatrix} s \\ s+l \end{bmatrix}} = 3 \cdot 2^{k+s-1}$$

The case $m = 0, l = 0$

$$\Delta_k \Gamma_s^{\begin{bmatrix} s \\ s \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s-1 \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s \\ s-1 \end{bmatrix}} + 32 \cdot \Delta_k \Gamma_{s-2}^{\begin{bmatrix} s-1 \\ s-1 \end{bmatrix}} = \Delta_k \omega_s(s, 0, 0) = 2^{k+s-1}$$

$$\text{since } \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s-1 \\ s \end{bmatrix}} = 2^{k+s-2}, \quad \Delta_k \Gamma_{s-2}^{\begin{bmatrix} s-1 \\ s-1 \end{bmatrix}} = 0 \quad \text{we get } \Delta_k \Gamma_s^{\begin{bmatrix} s \\ s \end{bmatrix}} = 2^{k+s-1} + 12 \cdot 2^{k+s-2} = 7 \cdot 2^{k+s-1}$$

proof of (9.2)

The case $m \geq 2, l \geq 0$

$$\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_s^{\begin{bmatrix} s+m-1 \\ s+m+l \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_s^{\begin{bmatrix} s+m \\ s+m+l-1 \end{bmatrix}} + 32 \cdot \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s+m-1 \\ s+m+l-1 \end{bmatrix}} = \Delta_k \omega_{s+1}(s, m, l) = -3 \cdot 2^{k+s-1}$$

$$\text{since } \Delta_k \Gamma_s^{\begin{bmatrix} s+m-1 \\ s+m+l \end{bmatrix}} = \Delta_k \Gamma_s^{\begin{bmatrix} s+m \\ s+m+l-1 \end{bmatrix}} = 2^{k+s-1} \text{ and } \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s+m-1 \\ s+m+l-1 \end{bmatrix}} = 0$$

$$\text{we get } \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s+m \\ s+m+l \end{bmatrix}} = 12 \cdot 2^{k+s-1} - 3 \cdot 2^{k+s-1} = 9 \cdot 2^{k+s-1}$$

The case $m = 1, l \geq 1$

$$\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s+1 \\ s+1+l \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_s^{\begin{bmatrix} s \\ s+1+l \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_s^{\begin{bmatrix} s+1 \\ s+l \end{bmatrix}} + 32 \cdot \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s \\ s+l \end{bmatrix}} = \Delta_k \omega_{s+1}(s, 1, l) = -3 \cdot 2^{k+s-1}$$

since $\Delta_k \Gamma_s \left[\begin{smallmatrix} s \\ s+1+l \end{smallmatrix} \right] = 3 \cdot 2^{k+s-1}$, $\Delta_k \Gamma_s \left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix} \right] = 2^{k+s-1}$ and $\Delta_k \Gamma_{s-1} \left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right] = 0$

we get $\Delta_k \Gamma_{s+1} \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] = 12 \cdot 2^{k+s-1} + 8 \cdot 2^{k+s-1} - 3 \cdot 2^{k+s-1} = 17 \cdot 2^{k+s-1}$

The case $m = 1, l = 0$

$$\Delta_k \Gamma_{s+1} \left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix} \right] - 4 \cdot \Delta_k \Gamma_s \left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_s \left[\begin{smallmatrix} s \\ s \end{smallmatrix} \right] + 32 \cdot \Delta_k \Gamma_{s-1} \left[\begin{smallmatrix} s \\ s \end{smallmatrix} \right] = \Delta_k \omega_{s+1}(s, 1, 0) = -3 \cdot 2^{k+s-1}$$

since $\Delta_k \Gamma_s \left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix} \right] = 3 \cdot 2^{k+s-1}$ and $\Delta_k \Gamma_{s-1} \left[\begin{smallmatrix} s \\ s \end{smallmatrix} \right] = 0$ we get $\Delta_k \Gamma_{s+1} \left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix} \right] = 36 \cdot 2^{k+s-1} - 3 \cdot 2^{k+s-1} = 33 \cdot 2^{k+s-1}$

The case $m = 0, l = 0$

$$\Delta_k \Gamma_{s+1} \left[\begin{smallmatrix} s \\ s \end{smallmatrix} \right] - 4 \cdot \Delta_k \Gamma_s \left[\begin{smallmatrix} s-1 \\ s \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_s \left[\begin{smallmatrix} s-1 \\ s \end{smallmatrix} \right] + 32 \cdot \Delta_k \Gamma_{s-1} \left[\begin{smallmatrix} s-1 \\ s-1 \end{smallmatrix} \right] = \Delta_k \omega_{s+1}(s, 0, 0) = -3 \cdot 2^{k+s-1}$$

since $\Delta_k \Gamma_s \left[\begin{smallmatrix} s-1 \\ s \end{smallmatrix} \right] = 33 \cdot 2^{k+s-2}$ and $\Delta_k \Gamma_{s-1} \left[\begin{smallmatrix} s-1 \\ s-1 \end{smallmatrix} \right] = 3 \cdot 2^{k+s-2}$ we get $\Delta_k \Gamma_{s+1} \left[\begin{smallmatrix} s \\ s \end{smallmatrix} \right]$
 $= 12 \cdot 33 \cdot 2^{k+s-2} - 32 \cdot 3 \cdot 2^{k+s-2} - 3 \cdot 2^{k+s-1} = 147 \cdot 2^{k+s-1}$

The case $m = 0, l = 1$

$$\Delta_k \Gamma_{s+1} \left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix} \right] - 4 \cdot \Delta_k \Gamma_s \left[\begin{smallmatrix} s-1 \\ s+1 \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_s \left[\begin{smallmatrix} s \\ s \end{smallmatrix} \right] + 32 \cdot \Delta_k \Gamma_{s-1} \left[\begin{smallmatrix} s-1 \\ s \end{smallmatrix} \right] = \Delta_k \omega_{s+1}(s, 0, 1) = -3 \cdot 2^{k+s-1}$$

since $\Delta_k \Gamma_s \left[\begin{smallmatrix} s-1 \\ s+1 \end{smallmatrix} \right] = 17 \cdot 2^{k+s-2}$, $\Delta_k \Gamma_s \left[\begin{smallmatrix} s \\ s \end{smallmatrix} \right] = 7 \cdot 2^{k+s-1}$ and $\Delta_k \Gamma_{s-1} \left[\begin{smallmatrix} s-1 \\ s \end{smallmatrix} \right] = 2^{k+s-2}$ we get

$$\Delta_k \Gamma_{s+1} \left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix} \right] = 4 \cdot 17 \cdot 2^{k+s-2} + 8 \cdot 7 \cdot 2^{k+s-1} - 32 \cdot 2^{k+s-2} - 3 \cdot 2^{k+s-1} = 71 \cdot 2^{k+s-1}$$

The case $m = 0, l \geq 2$

$$\Delta_k \Gamma_{s+1} \left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right] - 4 \cdot \Delta_k \Gamma_s \left[\begin{smallmatrix} s-1 \\ s+l \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_s \left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right] + 32 \cdot \Delta_k \Gamma_{s-1} \left[\begin{smallmatrix} s-1 \\ s+l-1 \end{smallmatrix} \right] = \Delta_k \omega_{s+1}(s, 0, l) = -3 \cdot 2^{k+s-1}$$

since $\Delta_k \Gamma_s \left[\begin{smallmatrix} s-1 \\ s+l \end{smallmatrix} \right] = 17 \cdot 2^{k+s-2}$, $\Delta_k \Gamma_s \left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right] = 3 \cdot 2^{k+s-1}$ and $\Delta_k \Gamma_{s-1} \left[\begin{smallmatrix} s-1 \\ s+l-1 \end{smallmatrix} \right] = 2^{k+s-2}$ we get

$$\Delta_k \Gamma_{s+1} \left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right] = 4 \cdot 17 \cdot 2^{k+s-2} + 8 \cdot 3 \cdot 2^{k+s-1} - 32 \cdot 2^{k+s-2} - 3 \cdot 2^{k+s-1} = 39 \cdot 2^{k+s-1}$$

proof of (9.3)

The case $m \geq 3, l \geq 0$

$$\Delta_k \Gamma_{s+2} \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] - 4 \cdot \Delta_k \Gamma_{s+1} \left[\begin{smallmatrix} s+m-1 \\ s+m+l \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_{s+1} \left[\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix} \right] + 32 \cdot \Delta_k \Gamma_s \left[\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] = \Delta_k \omega_{s+2}(s, m, l) = 2^{k+s}$$

since $\Delta_k \Gamma_{s+1} \left[\begin{smallmatrix} s+m-1 \\ s+m+l \end{smallmatrix} \right] = \Delta_k \Gamma_{s+1} \left[\begin{smallmatrix} s+m \\ s+m+l-1 \end{smallmatrix} \right] = 9 \cdot 2^{k+s-1}$ and $\Delta_k \Gamma_s \left[\begin{smallmatrix} s+m-1 \\ s+m+l-1 \end{smallmatrix} \right] = 2^{k+s-1}$

we get $\Delta_k \Gamma_{s+2} \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] = 12 \cdot 9 \cdot 2^{k+s-1} - 32 \cdot 2^{k+s-1} + 2^{k+s} = 78 \cdot 2^{k+s-1}$

The case $m = 2, l \geq 1$

$$\Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s+2+l \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+2+l \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+2+l-1 \end{bmatrix}} + 32 \cdot \Delta_k \Gamma_s^{\begin{bmatrix} s \\ s+l+1 \end{bmatrix}} = \Delta_k \omega_{s+2}(s, 2, l) = 2^{k+s}$$

since $\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+2+l \end{bmatrix}} = 17 \cdot 2^{k+s-1}$, $\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+l+1 \end{bmatrix}} = 9 \cdot 2^{k+s-1}$ and $\Delta_k \Gamma_s^{\begin{bmatrix} s \\ s+l+1 \end{bmatrix}} = 2^{k+s-1}$

we get $\Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s+2+l \end{bmatrix}} = 4 \cdot 17 \cdot 2^{k+s-1} + 8 \cdot 9 \cdot 2^{k+s-1} - 32 \cdot 2^{k+s-1} + 2^{k+s} = 110 \cdot 2^{k+s-1}$

The case $m = 2, l = 0$

$$\Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s+2 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+2 \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+2 \end{bmatrix}} + 32 \cdot \Delta_k \Gamma_s^{\begin{bmatrix} s \\ s+1 \end{bmatrix}} = \Delta_k \omega_{s+2}(s, 2, 0) = 2^{k+s}$$

since $\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+2 \end{bmatrix}} = 17 \cdot 2^{k+s-1}$ and $\Delta_k \Gamma_s^{\begin{bmatrix} s \\ s+1 \end{bmatrix}} = 2^{k+s-1}$

we get $\Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s+2 \end{bmatrix}} = 12 \cdot 17 \cdot 2^{k+s-1} - 32 \cdot 2^{k+s-1} + 2^{k+s} = 174 \cdot 2^{k+s-1}$

The case $m = 1, l \geq 2$

$$\Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s+1+l \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+1+l \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+l \end{bmatrix}} + 32 \cdot \Delta_k \Gamma_s^{\begin{bmatrix} s \\ s+l \end{bmatrix}} = \Delta_k \omega_{s+2}(s, 1, l) = 2^{k+s}$$

since $\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+1+l \end{bmatrix}} = 39 \cdot 2^{k+s-1}$, $\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+l \end{bmatrix}} = 17 \cdot 2^{k+s-1}$ and $\Delta_k \Gamma_s^{\begin{bmatrix} s \\ s+l \end{bmatrix}} = 3 \cdot 2^{k+s-1}$

we get $\Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s+1+l \end{bmatrix}} = 4 \cdot 39 \cdot 2^{k+s-1} + 8 \cdot 17 \cdot 2^{k+s-1} - 32 \cdot 3 \cdot 2^{k+s-1} + 2^{k+s} = 198 \cdot 2^{k+s-1}$

The case $m = 1, l = 1$

$$\Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}} + 32 \cdot \Delta_k \Gamma_s^{\begin{bmatrix} s \\ s+1 \end{bmatrix}} = \Delta_k \omega_{s+2}(s, 1, 1) = 2^{k+s}$$

since $\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}} = 39 \cdot 2^{k+s-1}$, $\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}} = 33 \cdot 2^{k+s-1}$ and $\Delta_k \Gamma_s^{\begin{bmatrix} s \\ s+1 \end{bmatrix}} = 3 \cdot 2^{k+s-1}$

we get $\Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}} = 4 \cdot 39 \cdot 2^{k+s-1} + 8 \cdot 33 \cdot 2^{k+s-1} - 32 \cdot 3 \cdot 2^{k+s-1} + 2^{k+s} = 326 \cdot 2^{k+s-1}$

The case $m = 1, l = 0$

$$\Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}} + 32 \cdot \Delta_k \Gamma_s^{\begin{bmatrix} s \\ s \end{bmatrix}} = \Delta_k \omega_{s+2}(s, 1, 0) = 2^{k+s}$$

since $\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}} = 71 \cdot 2^{k+s-1}$, and $\Delta_k \Gamma_s^{\begin{bmatrix} s \\ s \end{bmatrix}} = 7 \cdot 2^{k+s-1}$

we get $\Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}} = 12 \cdot 71 \cdot 2^{k+s-1} - 32 \cdot 7 \cdot 2^{k+s-1} + 2^{k+s} = 630 \cdot 2^{k+s-1}$

The case $m = 0, l \geq 3$

$$\Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s-l \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+l \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+l-1 \end{bmatrix}} + 32 \cdot \Delta_k \Gamma_s^{\begin{bmatrix} s-1 \\ s+l-1 \end{bmatrix}} = \Delta_k \omega_{s+2}(s, 0, l) = 2^{k+s}$$

since $\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s-1 \\ s \\ s+l \end{bmatrix}} = 198 \cdot 2^{k+s-2}$, $\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s \\ s+l-1 \end{bmatrix}} = 39 \cdot 2^{k+s-1}$ and $\Delta_k \Gamma_s^{\begin{bmatrix} s-1 \\ s \\ s+l-1 \end{bmatrix}} = 17 \cdot 2^{k+s-2}$

we get $\Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = 4 \cdot 198 \cdot 2^{k+s-2} + 8 \cdot 39 \cdot 2^{k+s-1} - 32 \cdot 17 \cdot 2^{k+s-2} + 2^{k+s} = 438 \cdot 2^{k+s-1}$

The case $m = 0, l = 2$

$\Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s-1 \\ s \\ s+2 \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s \\ s+1 \end{bmatrix}} + 32 \cdot \Delta_k \Gamma_s^{\begin{bmatrix} s-1 \\ s \\ s+1 \end{bmatrix}} = \Delta_k \omega_{s+2}(s, 0, 2) = 2^{k+s}$

since $\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s-1 \\ s \\ s+2 \end{bmatrix}} = 198 \cdot 2^{k+s-2}$, $\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s \\ s+1 \end{bmatrix}} = 71 \cdot 2^{k+s-1}$ and $\Delta_k \Gamma_s^{\begin{bmatrix} s-1 \\ s \\ s+1 \end{bmatrix}} = 17 \cdot 2^{k+s-2}$

we get $\Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = 4 \cdot 198 \cdot 2^{k+s-2} + 8 \cdot 71 \cdot 2^{k+s-1} - 32 \cdot 17 \cdot 2^{k+s-2} + 2^{k+s} = 694 \cdot 2^{k+s-1}$

The case $m = 0, l = 1$

$\Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s-1 \\ s \\ s+1 \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} + 32 \cdot \Delta_k \Gamma_s^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} = \Delta_k \omega_{s+2}(s, 0, 1) = 2^{k+s}$

since $\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s-1 \\ s \\ s+1 \end{bmatrix}} = 326 \cdot 2^{k+s-2}$, $\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = 147 \cdot 2^{k+s-1}$ and $\Delta_k \Gamma_s^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} = 33 \cdot 2^{k+s-2}$

we get $\Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = 4 \cdot 326 \cdot 2^{k+s-2} + 8 \cdot 147 \cdot 2^{k+s-1} - 32 \cdot 33 \cdot 2^{k+s-2} + 2^{k+s} = 1302 \cdot 2^{k+s-1}$

The case $m = 0, l = 0$

$\Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s-1 \\ s \\ s+1 \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} + 32 \cdot \Delta_k \Gamma_s^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} = \Delta_k \omega_{s+2}(s, 0, 0) = 2^{k+s}$

since $\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s-1 \\ s \\ s+1 \end{bmatrix}} = 630 \cdot 2^{k+s-2}$, and $\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = 71 \cdot 2^{k+s-1}$

we get $\Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = 12 \cdot 630 \cdot 2^{k+s-2} - 32 \cdot 71 \cdot 2^{k+s-1} + 2^{k+s} = 2646 \cdot 2^{k+s-1}$

□

9.4. **Computation of** $\Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{s+j-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}}$, for $0 \leq j \leq s-1$.

Lemma 9.3. *We have*

$$(9.4) \quad \Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{s+j-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} = \begin{cases} 7 \cdot 2^{k+s-1} & \text{if } j = 0 \\ 147 \cdot 2^{k+s-1} & \text{if } j = 1 \\ 427 \cdot 2^{k+s} & \text{if } j = 2 \\ 441 \cdot 2^{k+s+3j-6} & \text{if } 3 \leq j \leq s-1 \end{cases}$$

Proof. We use Lemma 7.4, Lemma 9.2, (6.33) and (8.9)

The case $j = 0$

$$\Delta_k \Gamma_s^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{s-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} = \Delta_k \Gamma_s^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = 7 \cdot 2^{k+s-1}$$

The case $j = 1$

$$\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{s-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} = \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = 147 \cdot 2^{k+s-1}$$

The case $j = 2$

$$\begin{aligned} \Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} &= 63 \cdot 2^{k+s} \iff 4 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 32 \cdot \Delta_k \Gamma_s^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} + \Delta_k \omega_{s+2}(s, 0, 0) = 63 \cdot 2^{k+s} \\ \iff \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_s^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} &= 2^{-2} \cdot [63 \cdot 2^{k+s} - \Delta_k \omega_{s+2}(s, 0, 0)] = 2^{-2} \cdot [63 \cdot 2^{k+s} - 2^{k+s}] = 31 \cdot 2^{k+s-1} \\ \iff 4 \cdot \Delta_k \Gamma_s^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} - 32 \cdot \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} &+ \Delta_k \omega_{s+1}(s-1, 1, 0) = 31 \cdot 2^{k+s-1} \\ \iff \Delta_k \Gamma_s^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} &= 2^{-2} \cdot [31 \cdot 2^{k+s-1} - \Delta_k \omega_{s+1}(s-1, 1, 0)] = 2^{-2} \cdot [31 \cdot 2^{k+s-1} - 2^{k+s-1}] = 15 \cdot 2^{k+s-2} \end{aligned}$$

We then obtain the following equalities,

$$\begin{aligned} \Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} &= 63 \cdot 2^{k+s} \\ 2^3 \cdot (\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_s^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}}) &= 2^3 \cdot 31 \cdot 2^{k+s-1} \\ 2^6 \cdot (\Delta_k \Gamma_s^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}}) &= 2^6 \cdot 15 \cdot 2^{k+s-2}. \end{aligned}$$

Summing the above equalities we get

$$\Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} = 2^{k+s} \cdot [63 + 31 \cdot 2^2 + 15 \cdot 2^4] = 427 \cdot 2^{k+s}$$

The case $3 \leq j \leq s-1$

$$\begin{aligned} \Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} &= 63 \cdot 2^{k+s+3j-6} \\ \Updownarrow & \\ 4 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 32 \cdot \Delta_k \Gamma_{s+j-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} &+ \Delta_k \omega_{s+j}(s, 0, 0) = 63 \cdot 2^{k+s+3j-6} \\ \Updownarrow & \\ \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+j-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} &= 2^{-2} \cdot [63 \cdot 2^{k+s+3j-6} - \Delta_k \omega_{s+j}(s, 0, 0)] = 2^{-2} \cdot [63 \cdot 2^{k+s+3j-6} - 0] = 63 \cdot 2^{k+s+3j-8} \\ \Updownarrow & \\ 4 \cdot \Delta_k \Gamma_{s+j-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} - 32 \cdot \Delta_k \Gamma_{s+j-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} &+ \Delta_k \omega_{s+j-1}(s-1, 1, 0) = 63 \cdot 2^{k+s+3j-8} \\ \Updownarrow & \\ \Delta_k \Gamma_{s+j-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} &= 2^{-2} \cdot [63 \cdot 2^{k+s+3j-8} - \Delta_k \omega_{s+j-1}(s-1, 1, 0)] = [63 \cdot 2^{k+s+3j-10} - 0] = 63 \cdot 2^{k+s+3j-10} \end{aligned}$$

We then obtain the following equalities

$$\begin{aligned}\Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} &= 63 \cdot 2^{k+s+3j-6} \\ 2^3 \cdot (\Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+j-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}}) &= 2^3 \cdot 63 \cdot 2^{k+s+3j-8} \\ 2^6 \cdot (\Delta_k \Gamma_{s+j-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+j-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}}) &= 2^6 \cdot 63 \cdot 2^{k+s+3j-10}\end{aligned}$$

Summing the above equalities we get

$$\Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} = 63 \cdot 2^{k+s+3j-6} \cdot [1 + 2 + 2^2] = 63 \cdot 7 \cdot 2^{k+s+3j-6} = 441 \cdot 2^{k+s+3j-6}$$

□

9.5. Computation of $\Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}}$ for $0 \leq j \leq s-1$.

Lemma 9.4. *We have*

$$(9.5) \quad \Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = \begin{cases} 7 \cdot 2^{k+s-1} & \text{if } j = 0 \\ 147 \cdot [5 \cdot 2^{k+s+4j-7} - 2^{k+s+3j-6}] = 147 \cdot (5 \cdot 2^{j-1} - 1) \cdot 2^{k+s+3j-6} & \text{if } 1 \leq j \leq s-1 \end{cases}$$

Proof. We obtain from (9.4)

The case $j = 0$

$$\Delta_k \Gamma_s^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = 7 \cdot 2^{k+s-1}$$

The case $j = 1$

$$\Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = 147 \cdot 2^{k+s-1}$$

The case $j = 2$

$$\Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = 427 \cdot 2^{k+s} + 2^9 \cdot \Delta_k \Gamma_{s-1}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} = 427 \cdot 2^{k+s} + 2^9 \cdot 7 \cdot 2^{k+s-2} = 1323 \cdot 2^{k+s}$$

The case $3 \leq j \leq s-1$, $j \equiv 0 \pmod{2}$

$$\begin{aligned}\Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{s+j-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} &= 441 \cdot 2^{k+s+3j-6} \\ 2^9 \cdot (\Delta_k \Gamma_{s+j-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{s+j-6}^{\begin{bmatrix} s-2 \\ s-2 \\ s-2 \end{bmatrix}}) &= 441 \cdot 2^{k+(s-1)+3(j-2)-6} \cdot 2^9 \\ (2^9)^2 \cdot (\Delta_k \Gamma_{s+j-6}^{\begin{bmatrix} s-2 \\ s-2 \\ s-2 \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{s+j-9}^{\begin{bmatrix} s-3 \\ s-3 \\ s-3 \end{bmatrix}}) &= 441 \cdot 2^{k+(s-2)+3(j-4)-6} \cdot (2^9)^2 \\ &\vdots\end{aligned}$$

$$\begin{aligned}
(2^9)^{\frac{j}{2}-2} \cdot \left(\Delta_k \Gamma_{s+j-3(\frac{j}{2}-2)}^{\begin{bmatrix} s-(\frac{j}{2}-2) \\ s-(\frac{j}{2}-2) \\ s-(\frac{j}{2}-2) \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{s+j-3(\frac{j}{2}-1)}^{\begin{bmatrix} s-(\frac{j}{2}-1) \\ s-(\frac{j}{2}-1) \\ s-(\frac{j}{2}-1) \end{bmatrix}} \right) &= 441 \cdot 2^{k+(s-(\frac{j}{2}-2)+3(j-(j-4)))-6} \cdot (2^9)^{\frac{j}{2}-2} \\
(2^9)^{\frac{j}{2}-1} \cdot \left(\Delta_k \Gamma_{s+j-3(\frac{j}{2}-1)}^{\begin{bmatrix} s-(\frac{j}{2}-1) \\ s-(\frac{j}{2}-1) \\ s-(\frac{j}{2}-1) \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{s+j-3(\frac{j}{2})}^{\begin{bmatrix} s-(\frac{j}{2}) \\ s-(\frac{j}{2}) \\ s-(\frac{j}{2}) \end{bmatrix}} \right) &= 427 \cdot 2^{k+s-\frac{j}{2}+1} \cdot (2^9)^{\frac{j}{2}-1} \\
(2^9)^{\frac{j}{2}} \cdot \left(\Delta_k \Gamma_{s+j-3(\frac{j}{2})}^{\begin{bmatrix} s-(\frac{j}{2}) \\ s-(\frac{j}{2}) \\ s-(\frac{j}{2}) \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{s+j-3(\frac{j}{2}+1)}^{\begin{bmatrix} s-(\frac{j}{2}+1) \\ s-(\frac{j}{2}+1) \\ s-(\frac{j}{2}+1) \end{bmatrix}} \right) &= 7 \cdot 2^{k+s-\frac{j}{2}-1} \cdot (2^9)^{\frac{j}{2}}
\end{aligned}$$

We obtain by summing the above equations

$$\begin{aligned}
\Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} &= \sum_{d=0}^{\frac{j}{2}-2} 441 \cdot 2^{k+(s-d)+3(j-2d)-6} \cdot 2^{qd} + 427 \cdot 2^{k+s-\frac{j}{2}+1} \cdot (2^9)^{\frac{j}{2}-1} + 7 \cdot 2^{k+s-\frac{j}{2}-1} \cdot (2^9)^{\frac{j}{2}} \\
&= 441 \cdot 2^{k+s+3j-6} \sum_{d=0}^{\frac{j}{2}-2} 2^{2d} + 427 \cdot 2^{k+s+4j-8} + 7 \cdot 2^{k+s+4j-1} \\
&= 441 \cdot 2^{k+s+3j-6} \cdot \frac{2^{j-2}-1}{2^2-1} + 427 \cdot 2^{k+s+4j-8} + 7 \cdot 2^{k+s+4j-1} \\
&= 147 \cdot 2^{k+s+4j-8} - 147 \cdot 2^{k+s+3j-6} + 427 \cdot 2^{k+s+4j-8} + 7 \cdot 2^{k+s+4j-1} \\
&= -147 \cdot 2^{k+s+3j-6} + 735 \cdot 2^{k+s+4j-7} \\
&= 147 \cdot [5 \cdot 2^{k+s+4j-7} - 2^{k+s+3j-6}]
\end{aligned}$$

The case $3 \leq j \leq s-1$, $j \equiv 1 \pmod{2}$

Similarly to the proof in the case $j \equiv 0 \pmod{2}$

□

9.6. Computation of $\Delta_k \Gamma_{2s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{2s+j-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}}$, for $0 \leq j \leq s$.

Lemma 9.5. *We have*

$$(9.6) \quad \Delta_k \Gamma_{2s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{2s+j-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} = \begin{cases} 21 \cdot 2^{2k+2s-2} - 231 \cdot 2^{k+4s-6} & \text{if } j = 0 \\ -2373 \cdot 2^{2k+2s-2} + 21 \cdot 2^{k+4s-3} & \text{if } j = 1 \\ -7 \cdot 315 \cdot 2^{2k+2s+4j-6} & \text{if } 2 \leq j \leq s-1 \\ -441 \cdot 2^{3k+3s-3} + 147 \cdot 2^{2k+6s-6} & \text{if } j = s \end{cases}$$

Proof. We use Lemma 7.4, (8.10) and (8.3)

The case $j = 0$

$$\begin{aligned}
\Delta_k \Gamma_{2s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s-1}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} &= -15 \cdot 2^{2k+2s-2} - 33 \cdot 2^{k+4s-6} \\
\Downarrow & \\
4 \cdot \Delta_k \Gamma_{2s-1}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} - 32 \cdot \Delta_k \Gamma_{2s-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} + \Delta_k \omega_{2s}(s, 0, 0) &= -15 \cdot 2^{2k+2s-2} - 33 \cdot 2^{k+4s-6} \\
\Downarrow & \\
\Delta_k \Gamma_{2s-1}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} &= 2^{-2} \cdot [-15 \cdot 2^{2k+2s-2} - 33 \cdot 2^{k+4s-6} - \Delta_k \omega_{2s}(s, 0, 0)] \\
&= 2^{-2} \cdot [-15 \cdot 2^{2k+2s-2} - 33 \cdot 2^{k+4s-6} + 9 \cdot 2^{2k+2s-2}] = -3 \cdot 2^{2k+2s-3} - 33 \cdot 2^{k+4s-8} \\
\Downarrow &
\end{aligned}$$

$$\begin{aligned}
& 4 \cdot \Delta_k \Gamma_{2s-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} - 32 \cdot \Delta_k \Gamma_{2s-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} + \Delta_k \omega_{2s-1}(s-1, 1, 0) = -3 \cdot 2^{2k+2s-3} - 33 \cdot 2^{k+4s-8} \\
& \Updownarrow \\
& \Delta_k \Gamma_{2s-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} = 2^{-2} \cdot [-3 \cdot 2^{2k+2s-3} - 33 \cdot 2^{k+4s-8} - \Delta_k \omega_{2s-1}(s-1, 1, 0)] \\
& = 2^{-2} \cdot [-3 \cdot 2^{2k+2s-3} - 33 \cdot 2^{k+4s-8} + 9 \cdot 2^{2k+2s-3}] = -3 \cdot 2^{2k+2s-4} - 33 \cdot 2^{k+4s-10}
\end{aligned}$$

We then obtain the following equalities

$$\begin{aligned}
& \Delta_k \Gamma_{2s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s-1}^{\begin{bmatrix} s-1 \\ s \\ s-1 \end{bmatrix}} = -15 \cdot 2^{2k+2s-2} - 33 \cdot 2^{k+4s-6} \\
& 2^3 \cdot (\Delta_k \Gamma_{2s-1}^{\begin{bmatrix} s-1 \\ s \\ s-1 \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}}) = 2^3 \cdot [-3 \cdot 2^{2k+2s-3} - 33 \cdot 2^{k+4s-8}] \\
& 2^6 \cdot (\Delta_k \Gamma_{2s-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}}) = 2^6 \cdot [3 \cdot 2^{2k+2s-4} - 33 \cdot 2^{k+4s-10}]
\end{aligned}$$

Summing the above equalities we get

$$\Delta_k \Gamma_{2s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{2s-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} = 2^{2k+2s-2} \cdot [-15 - 12 + 48] + 2^{k+4s-6} \cdot [-33 - 33 \cdot 2 - 33 \cdot 2^2] = 21 \cdot 2^{2k+2s-2} - 231 \cdot 2^{k+4s-6}$$

The case $j = 1$

$$\begin{aligned}
& \Delta_k \Gamma_{2s+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s}^{\begin{bmatrix} s-1 \\ s \\ s-1 \end{bmatrix}} = -321 \cdot 2^{2k+2s-2} + 3 \cdot 2^{k+4s-3} \\
& \Updownarrow \\
& 4 \cdot \Delta_k \Gamma_{2s}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 32 \cdot \Delta_k \Gamma_{2s-1}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} + \Delta_k \omega_{2s+1}(s, 0, 0) = -321 \cdot 2^{2k+2s-2} + 3 \cdot 2^{k+4s-3} \\
& \Updownarrow \\
& \Delta_k \Gamma_{2s}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s-1}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} = 2^{-2} \cdot [-321 \cdot 2^{2k+2s-2} + 3 \cdot 2^{k+4s-3} - \Delta_k \omega_{2s+1}(s, 0, 0)] \\
& = 2^{-2} \cdot [-321 \cdot 2^{2k+2s-2} + 3 \cdot 2^{k+4s-3} - 9 \cdot 2^{2k+2s-2}] = -165 \cdot 2^{2k+2s-3} + 3 \cdot 2^{k+4s-5} \\
& \Updownarrow \\
& 4 \cdot \Delta_k \Gamma_{2s-1}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} - 32 \cdot \Delta_k \Gamma_{2s-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} + \Delta_k \omega_{2s}(s-1, 1, 0) = -165 \cdot 2^{2k+2s-3} + 3 \cdot 2^{k+4s-5} \\
& \Updownarrow \\
& \Delta_k \Gamma_{2s-1}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} = 2^{-2} \cdot [-165 \cdot 2^{2k+2s-3} + 3 \cdot 2^{k+4s-5} - \Delta_k \omega_{2s}(s-1, 1, 0)] \\
& = 2^{-2} \cdot [-165 \cdot 2^{2k+2s-3} + 3 \cdot 2^{k+4s-5} - 9 \cdot 2^{2k+2s-3}] = -87 \cdot 2^{2k+2s-4} + 3 \cdot 2^{k+4s-7}
\end{aligned}$$

We then obtain the following equalities

$$\begin{aligned}
& \Delta_k \Gamma_{2s+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s}^{\begin{bmatrix} s-1 \\ s \\ s-1 \end{bmatrix}} = -321 \cdot 2^{2k+2s-2} + 3 \cdot 2^{k+4s-3} \\
& 2^3 \cdot (\Delta_k \Gamma_{2s}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s-1}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}}) = 2^3 \cdot [-165 \cdot 2^{2k+2s-3} + 3 \cdot 2^{k+4s-5}] \\
& 2^6 \cdot (\Delta_k \Gamma_{2s-1}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}}) = 2^6 \cdot [-87 \cdot 2^{2k+2s-4} + 3 \cdot 2^{k+4s-7}]
\end{aligned}$$

Summing the above equalities we get

$$\begin{aligned} \Delta_k \Gamma_{2s+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{2s-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} &= 2^{2k+2s-2} \cdot [-321 + (-165) \cdot 2^2 + (-87) \cdot 2^4] + 2^{k+4s-3} \cdot [3 + 3 \cdot 2 + 3 \cdot 2^2] \\ &= -2373 \cdot 2^{2k+2s-2} + 21 \cdot 2^{k+4s-3} \end{aligned}$$

The case $2 \leq j \leq s-1$

$$\begin{aligned} \Delta_k \Gamma_{2s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} &= -315 \cdot 2^{2k+2s+4j-6} \\ \Downarrow \\ 4 \cdot \Delta_k \Gamma_{2s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 32 \cdot \Delta_k \Gamma_{2s+j-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} + \Delta_k \omega_{2s+j}(s, 0, 0) &= -315 \cdot 2^{2k+2s+4j-6} \\ \Downarrow \\ \Delta_k \Gamma_{2s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+j-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} &= 2^{-2} \cdot [-315 \cdot 2^{2k+2s+4j-6} - \Delta_k \omega_{2s+j}(s, 0, 0)] \\ &= 2^{-2} \cdot [-315 \cdot 22k + 2s + 4j - 6 - 0] = -315 \cdot 2^{2k+2s+4j-8} \\ \Downarrow \\ 4 \cdot \Delta_k \Gamma_{2s+j-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} - 32 \cdot \Delta_k \Gamma_{2s+j-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} + \Delta_k \omega_{2s+j-1}(s-1, 1, 0) &= -315 \cdot 2^{2k+2s+4j-8} \\ \Downarrow \\ \Delta_k \Gamma_{2s+j-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+j-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} &= 2^{-2} \cdot [-315 \cdot 2^{2k+2s+4j-8} - \Delta_k \omega_{2s+j-1}(s-1, 1, 0)] \\ &= 2^{-2} \cdot [-315 \cdot 2^{2k+2s+4j-8} - 0] = -315 \cdot 2^{2k+2s+4j-10} \end{aligned}$$

We then obtain the following equalities

$$\begin{aligned} \Delta_k \Gamma_{2s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} &= -315 \cdot 2^{2k+2s+4j-6} \\ 2^3 \cdot (\Delta_k \Gamma_{2s+j-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+j-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}}) &= 2^3 \cdot [-315 \cdot 2^{2k+2s+4j-8}] \\ 2^6 \cdot (\Delta_k \Gamma_{2s+j-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+j-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}}) &= 2^6 \cdot [-315 \cdot 2^{2k+2s+4j-10}] \end{aligned}$$

Summing the above equalities we get

$$\Delta_k \Gamma_{2s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{2s+j-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} = -315 \cdot 2^{2k+2s+4j-6} \cdot [1 + 2 + 2^2] = 7 \cdot (-315) \cdot 2^{2k+2s+4j-6}$$

The case $j = s$

$$\begin{aligned} \Delta_k \Gamma_{3s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{3s-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} &= -21 \cdot 2^{3k+3s-3} + 21 \cdot 2^{2k+6s-6} \\ \Downarrow \\ 4 \cdot \Delta_k \Gamma_{3s-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 32 \cdot \Delta_k \Gamma_{3s-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} + \Delta_k \omega_{3s}(s, 0, 0) &= -21 \cdot 2^{3k+3s-3} + 21 \cdot 2^{2k+6s-6} \\ \Downarrow \end{aligned}$$

$$\begin{aligned}
& \Delta_k \Gamma_{3s-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{3s-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} = 2^{-2} \cdot [-21 \cdot 2^{3k+3s-3} + 21 \cdot 2^{2k+6s-6} - \Delta_k \omega_{3s}(s, 0, 0)] \\
& = 2^{-2} \cdot [-21 \cdot 2^{3k+3s-3} + 21 \cdot 2^{2k+6s-6} - 21 \cdot 2^{3k+3s-3}] = -21 \cdot 2^{3k+3s-4} + 21 \cdot 2^{2k+6s-8} \\
& \Updownarrow \\
& 4 \cdot \Delta_k \Gamma_{3s-2}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 32 \cdot \Delta_k \Gamma_{3s-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} + \Delta_k \omega_{3s-1}(s-1, 1, 0) = -21 \cdot 2^{3k+3s-4} + 21 \cdot 2^{2k+6s-8} \\
& \Updownarrow \\
& \Delta_k \Gamma_{3s-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{3s-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} = 2^{-2} \cdot [-21 \cdot 2^{3k+3s-4} + 21 \cdot 2^{2k+6s-8} - \Delta_k \omega_{3s-1}(s-1, 1, 0)] \\
& = 2^{-2} \cdot [-21 \cdot 2^{3k+3s-4} + 21 \cdot 2^{2k+6s-8} - 21 \cdot 2^{3k+3s-4}] = -21 \cdot 2^{3k+3s-5} + 21 \cdot 2^{2k+6s-10}
\end{aligned}$$

We then obtain the following equalities

$$\begin{aligned}
& \Delta_k \Gamma_{3s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{3s-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} = -21 \cdot 2^{3k+3s-3} + 21 \cdot 2^{2k+6s-6} \\
& 2^3 \cdot (\Delta_k \Gamma_{3s-1}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{3s-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}}) = 2^3 \cdot [-21 \cdot 2^{3k+3s-4} + 21 \cdot 2^{2k+6s-8}] \\
& 2^6 \cdot (\Delta_k \Gamma_{3s-2}^{\begin{bmatrix} s-1 \\ s \\ s \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{3s-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}}) = 2^6 \cdot [-21 \cdot 2^{3k+3s-5} + 21 \cdot 2^{2k+6s-10}]
\end{aligned}$$

Summing the above equalities we get

$$\begin{aligned}
& \Delta_k \Gamma_{3s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{3(s-1)}^{\begin{bmatrix} s-1 \\ s-1 \\ s \end{bmatrix}} = -21 \cdot 2^{3k+3s-3} \cdot [1 + 2^2 + 2^4] + 21 \cdot 2^{2k+6s-6} \cdot [1 + 2 + 2^2] \\
& = -441 \cdot 2^{3k+3s-3} + 7 \cdot 21 \cdot 2^{2k+6s-6} = -(21)^2 \cdot 2^{3k+3s-3} + 7 \cdot 21 \cdot 2^{2k+6s-6}
\end{aligned}$$

□

9.7. Computation of $\Delta_k \Gamma_{2s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}}$ for $0 \leq j \leq s$.

Lemma 9.6. *We have*

$$(9.7) \quad \Delta_k \Gamma_{2s+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = \begin{cases} 21 \cdot [2^{2k+2s-2} + 35 \cdot 2^{k+5s-7} - 39 \cdot 2^{k+4s-6}] & \text{if } j = 0 \\ 105 \cdot [3 \cdot 2^{2k+2s+4j-6} + 7 \cdot 2^{k+5s+4j-7} - 31 \cdot 2^{k+4s+5j-8}] & \text{if } 1 \leq j \leq s-1 \\ 7 \cdot 2^{3k+3s-3} - 21 \cdot 2^{2k+6s-6} + 7 \cdot 2^{k+9s-8} & \text{if } j = s \end{cases}$$

Proof. The case $j = 0$

We have from (9.5) with $s \rightarrow s-1$, $j = (s-1) - 1$

$$\Delta_k \Gamma_{2s-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} = \Delta_k \Gamma_{(s-1)+(s-2)}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} = 147 \cdot (5 \cdot 2^{s-3} - 1) \cdot 2^{k+4s-13} = 147 \cdot 5 \cdot 2^{k+5s-16} - 147 \cdot 2^{k+4s-13}$$

We then deduce from (9.6) with $j = 0$

$$(9.8) \quad \Delta_k \Gamma_{2s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = 2^9 \cdot \Delta_k \Gamma_{2s-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} + 21 \cdot 2^{2k+2s-2} - 231 \cdot 2^{k+4s-6}$$

$$\begin{aligned}
&= 147 \cdot 5 \cdot 2^{k+5s-7} - 147 \cdot 2^{k+4s-4} + 21 \cdot 2^{2k+2s-2} - 231 \cdot 2^{k+4s-6} \\
&= 21 \cdot [2^{2k+2s-2} + 35 \cdot 2^{k+5s-7} - 39 \cdot 2^{k+4s-6}]
\end{aligned}$$

The case $j = 1$

From (9.8) with $s \rightarrow s-1$ we obtain

$$\Delta_k \Gamma_{2s-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} = \Delta_k \Gamma_{2(s-1)}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} = 21 \cdot [2^{2k+2s-4} + 35 \cdot 2^{k+5s-12} - 39 \cdot 2^{k+4s-10}]$$

We then deduce from (9.6) with $j = 1$

$$\begin{aligned}
(9.9) \quad \Delta_k \Gamma_{2s+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} &= 2^9 \cdot \Delta_k \Gamma_{2s-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} - 2373 \cdot 2^{2k+2s-2} + 21 \cdot 2^{k+4s-3} \\
&= 21 \cdot [2^{2k+2s+5} + 35 \cdot 2^{k+5s-3} - 39 \cdot 2^{k+4s-1}] - 21 \cdot 113 \cdot 2^{2k+2s-2} + 21 \cdot 2^{k+4s-3} \\
&= 105 \cdot [3 \cdot 2^{2k+2s-2} + 7 \cdot 2^{k+5s-3} - 31 \cdot 2^{k+4s-3}]
\end{aligned}$$

The case $j = 2$

From (9.9) with $s \rightarrow s-1$ we obtain

$$\begin{aligned}
\Delta_k \Gamma_{2(s-1)+1}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} &= 105 \cdot [3 \cdot 2^{2k+2(s-1)-2} + 7 \cdot 2^{k+5(s-1)-3} - 31 \cdot 2^{k+4(s-1)-3}] \\
&= 105 \cdot [3 \cdot 2^{2k+2s-4} + 7 \cdot 2^{k+5s-8} - 31 \cdot 2^{k+4s-7}]
\end{aligned}$$

We then deduce from (9.6) with $j = 2$

$$\begin{aligned}
(9.10) \quad \Delta_k \Gamma_{2s+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} &= 2^9 \cdot \Delta_k \Gamma_{2(s-1)+1}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} + 7 \cdot (-315) \cdot 2^{2k+2s+2} \\
&= 2^9 \cdot 105 \cdot [3 \cdot 2^{2k+2s-4} + 7 \cdot 2^{k+5s-8} - 31 \cdot 2^{k+4s-7}] + 105 \cdot (-21) \cdot 2^{2k+2s+2} \\
&= 105 \cdot [3 \cdot 2^{2k+2s+2} + 7 \cdot 2^{k+5s+1} - 31 \cdot 2^{k+4s+2}]
\end{aligned}$$

We observe that

$$\begin{aligned}
16^3 \cdot \Delta_{k-3} \Gamma_{2(s-1)+1}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} &= 2^{12} \cdot 105 \cdot [3 \cdot 2^{2(k-3)+2(s-1)-2} + 7 \cdot 2^{k-3+5(s-1)-3} - 31 \cdot 2^{k-3+4(s-1)-3}] \\
&= 105 \cdot [3 \cdot 2^{2k+2s+2} + 7 \cdot 2^{k+5s+1} - 31 \cdot 2^{k+4s+2}]
\end{aligned}$$

From (9.10) and the the above equation we deduce

$$(9.11) \quad \Delta_k \Gamma_{2s+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = 16^3 \cdot \Delta_{k-3} \Gamma_{2(s-1)+1}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}}$$

The case $2 \leq j \leq s-1$

We have for $1 \leq j \leq s-2$

$$(9.12) \quad (H_j) \quad \Delta_k \Gamma_{2s+1+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = 16^{3j} \cdot \Delta_{k-3j} \Gamma_{2(s-j)+1}^{\begin{bmatrix} s-j \\ s-j \\ s-j \end{bmatrix}}$$

We show (9.12) by induction on j .

By (9.11) (H_j) holds for $j = 1$.

We shall show that (H_j) holds implies that (H_{j+1}) holds.

We have

$$\begin{aligned}
\Delta_k \Gamma_{2s+1+(j+1)}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} &= 2^9 \cdot \Delta_k \Gamma_{2s+1+(j+1)-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} + 7 \cdot (-315) \cdot 2^{2k+2s+4(j+2)-6} \\
&= 2^9 \cdot \Delta_k \Gamma_{2(s-1)+1+j}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} + 7 \cdot (-315) \cdot 2^{2k+2s+4j+2} \\
&= 2^9 \cdot 2^{12j} \cdot \Delta_k \Gamma_{2(s-(j+1))+1}^{\begin{bmatrix} s-1-j \\ s-1-j \\ s-1-j \end{bmatrix}} + 7 \cdot (-315) \cdot 2^{2k+2s+4j+2} \\
&= 2^{12j+9} \cdot 105 \cdot [3 \cdot 2^{2(k-3j)+2(s-1-j)-2} + 7 \cdot 2^{k-3j+5(s-1-j)-3} - 31 \cdot 2^{k-3j+4(s-1-j)-3}] + 21 \cdot (-105) \cdot 2^{2k+2s+4j+2} \\
&= 105 \cdot [3 \cdot 2^{2k+2s+4j+2} + 7 \cdot 2^{k+5s+4j+1} - 31 \cdot 2^{k+4s+5j+2}]
\end{aligned}$$

On the other hand

$$\begin{aligned}
16^{3(j+1)} \cdot \Delta_k \Gamma_{2s-3(j+1)}^{\begin{bmatrix} s-(j+1) \\ s-(j+1) \\ s-(j+1) \end{bmatrix}} &= 2^{12j+12} \cdot 105 \cdot [3 \cdot 2^{2(k-3(j+1))+2(s-j-1)-2} + 7 \cdot 2^{k-3(j+1)+5(s-j-1)-3} - 31 \cdot 2^{k-3(j+1)+4(s-j-1)-3}] \\
&= 105 \cdot [3 \cdot 2^{2k+2s+4j+2} + 7 \cdot 2^{k+5s+4j+1} - 31 \cdot 2^{k+4s+5j+2}]
\end{aligned}$$

We have now proved that for $0 \leq j \leq s-2$ we have

$$\Delta_k \Gamma_{2s+1+j}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = 105 \cdot [3 \cdot 2^{2k+2s+4j-2} + 7 \cdot 2^{k+5s+4j-3} - 31 \cdot 2^{k+4s+5j-3}]$$

The case $j = s$

We have successively

$$\begin{aligned}
\Delta_k \Gamma_{3s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{3(s-1)}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} &= -441 \cdot 2^{3k+3s-3} + 147 \cdot 2^{2k+6s-6} \\
2^9 \cdot (\Delta_k \Gamma_{3(s-1)}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{3(s-2)}^{\begin{bmatrix} s-2 \\ s-2 \\ s-2 \end{bmatrix}}) &= (-441 \cdot 2^{3k+3(s-1)-3} + 147 \cdot 2^{2k+6(s-1)-6}) \cdot 2^9 \\
(2^9)^2 \cdot (\Delta_k \Gamma_{3(s-2)}^{\begin{bmatrix} s-2 \\ s-2 \\ s-2 \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{3(s-3)}^{\begin{bmatrix} s-3 \\ s-3 \\ s-3 \end{bmatrix}}) &= (-441 \cdot 2^{3k+3(s-2)-3} + 147 \cdot 2^{2k+6(s-2)-6}) \cdot (2^9)^2 \\
&\vdots \\
(2^9)^{s-2} \cdot (\Delta_k \Gamma_{3(s-(s-2))}^{\begin{bmatrix} s-(s-2) \\ s-(s-2) \\ s-(s-2) \end{bmatrix}} - 2^9 \cdot \Delta_k \Gamma_{3(s-(s-1))}^{\begin{bmatrix} s-(s-1) \\ s-(s-1) \\ s-(s-1) \end{bmatrix}}) &= (-441 \cdot 2^{3k+3(s-(s-2))-3} + 147 \cdot 2^{2k+6(s-(s-2))-6}) \cdot (2^9)^{s-2}
\end{aligned}$$

We obtain by summing the above equations

$$\Delta_k \Gamma_{3s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = \sum_{d=0}^{s-2} (-441 \cdot 2^{3k+3(s-d)-3} + 147 \cdot 2^{2k+6(s-d)-6}) \cdot (2^9)^d + (2^9)^{s-1} \cdot \Delta_k \Gamma_3^{\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}}$$

$$\begin{aligned}
&= (2^9)^{s-1} \cdot \Delta_k \Gamma_3 \begin{bmatrix} 1 \\ s \\ 1 \end{bmatrix} + \sum_{d=0}^{s-2} -441 \cdot 2^{3k+3s+6d-3} + 147 \cdot 2^{2k+3d+6s-6} \\
&= 2^{9s-9} \cdot (7 \cdot 2^{3k} - 21 \cdot 2^{2k} + 7 \cdot 2^{k+1}) - 441 \cdot 2^{3k+3s-3} \sum_{d=0}^{s-2} 2^{6d} + 147 \cdot 2^{2k+6s-6} \sum_{d=0}^{s-2} 2^{3d} \\
&= 7 \cdot 2^{3k+9s-9} - 21 \cdot 2^{2k+9s-9} + 7 \cdot 2^{k+9s-8} - 441 \cdot 2^{3k+3s-3} \cdot \frac{2^{6s-6} - 1}{2^6 - 1} + 147 \cdot 2^{2k+6s-6} \cdot \frac{2^{3s-3} - 1}{2^3 - 1} \\
&= 7 \cdot 2^{3k+9s-9} - 21 \cdot 2^{2k+9s-9} + 7 \cdot 2^{k+9s-8} - 7 \cdot 2^{3k+9s-9} + 7 \cdot 2^{3k+3s-3} + 21 \cdot 2^{2k+9s-9} - 21 \cdot 2^{2k+6s-9} \\
&= 7 \cdot 2^{3k+3s-3} - 21 \cdot 2^{2k+6s-6} + 7 \cdot 2^{k+9s-8}
\end{aligned}$$

□

Lemma 9.7. *We have*

$$(9.13) \quad \Delta_k \Gamma_{s+j-1} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} = \begin{cases} 2^{k+s-2} & \text{if } j = 0 \\ 33 \cdot 2^{k+s-2} & \text{if } j = 1 \\ 7 \cdot (105 \cdot 2^{j-2} - 15) \cdot 2^{k+s+3j-8} & \text{if } 2 \leq j \leq s-1 \\ 9 \cdot 2^{2k+2s-3} + 21 \cdot 35 \cdot 2^{k+5s-10} - 393 \cdot 2^{k+4s-8} & \text{if } j = s \\ 159 \cdot 2^{2k+2s-3} + 735 \cdot 2^{k+5s-6} - 1629 \cdot 2^{k+4s-5} & \text{if } j = s+1 \end{cases}$$

$$(9.14) \quad \Delta_k \Gamma_{2s+j-1} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} = \begin{cases} 105 \cdot (3 \cdot 2^{2k+2s+4j-8} + 7 \cdot 2^{k+5s+4j-10} - 31 \cdot 2^{k+4s+5j-11}) & \text{if } 2 \leq j \leq s-1 \\ 7 \cdot 2^{3k+3s-4} - 21 \cdot 2^{2k+6s-8} + 7 \cdot 2^{k+9s-11} & \text{if } j = s \end{cases}$$

$$(9.15) \quad \Delta_k \Gamma_{s+j-2} \begin{bmatrix} s-1 \\ s \\ s-1 \end{bmatrix} = \begin{cases} 3 \cdot 2^{k+s-2} & \text{if } j = 1 \\ 71 \cdot 2^{k+s-2} & \text{if } j = 2 \\ 21 \cdot (35 \cdot 2^{j-2} - 8) \cdot 2^{k+s+3j-11} & \text{if } 3 \leq j \leq s-1 \\ 3 \cdot 2^{2k+2s-4} + 735 \cdot 2^{k+5s-13} - 45 \cdot 2^{k+4s-8} & \text{if } j = s \\ 81 \cdot 2^{2k+2s-4} + 735 \cdot 2^{k+5s-9} - 51 \cdot 2^{k+4s-3} & \text{if } j = s+1 \end{cases}$$

$$(9.16) \quad \Delta_k \Gamma_{2s+j-2} \begin{bmatrix} s-1 \\ s \\ s-1 \end{bmatrix} = \begin{cases} 315 \cdot 2^{2k+2s+4j-10} + 735 \cdot 2^{k+5s+4j-13} - 3255 \cdot 2^{k+4s+5j-14} & \text{if } 2 \leq j \leq s-1 \\ 7 \cdot 2^{3k+3s-5} - 21 \cdot 2^{2k+6s-10} + 7 \cdot 2^{k+9s-14} & \text{if } j = s \end{cases}$$

Proof. **proof of** (9.13), (9.14)

Combining (8.9), (8.10) and (9.5), (9.7) give the desired formulas.

proof of (9.15), (9.16)

We obtain from (8.3) with $m = l = 0$

$$\begin{aligned}
&\Delta_k \Gamma_{s+j} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 4 \cdot \Delta_k \Gamma_{s+j-1} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} = 8 \cdot \left[\Delta_k \Gamma_{s+j-1} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} - 4 \cdot \Delta_k \Gamma_{s+j-2} \begin{bmatrix} s-1 \\ s \\ s-1 \end{bmatrix} \right] + \Delta_k \omega_{s+j}(s, 0, 0) \\
(9.17) \quad &\iff 32 \cdot \Delta_k \Gamma_{s+j-2} \begin{bmatrix} s-1 \\ s \\ s-1 \end{bmatrix} = 12 \cdot \Delta_k \Gamma_{s+j-1} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} - \Delta_k \Gamma_{s+j} \begin{bmatrix} s \\ s \\ s \end{bmatrix} + \Delta_k \omega_{s+j}(s, 0, 0) \\
&\Delta_k \Gamma_{2s+j} \begin{bmatrix} s \\ s \\ s \end{bmatrix} - 4 \cdot \Delta_k \Gamma_{2s+j-1} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} = 8 \cdot \left[\Delta_k \Gamma_{2s+j-1} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} - 4 \cdot \Delta_k \Gamma_{2s+j-2} \begin{bmatrix} s-1 \\ s \\ s-1 \end{bmatrix} \right] + \Delta_k \omega_{2s+j}(s, 0, 0) \\
(9.18) \quad &\iff 32 \cdot \Delta_k \Gamma_{2s+j-2} \begin{bmatrix} s-1 \\ s \\ s-1 \end{bmatrix} = 12 \cdot \Delta_k \Gamma_{2s+j-1} \begin{bmatrix} s-1 \\ s \\ s \end{bmatrix} - \Delta_k \Gamma_{2s+j} \begin{bmatrix} s \\ s \\ s \end{bmatrix} + \Delta_k \omega_{2s+j}(s, 0, 0)
\end{aligned}$$

(9.15), (9.16) follows now from (9.17), (9.18) and (7.48) $\square$

10. **An explicit formula for $\Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k}$, for $s \leq i \leq 3s$, $k \geq i$**

10.1. **Notation.**

Definition 10.1. We recall that $\Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k}$ denote the number of rank i matrices of the form

$$\left(\begin{array}{cccccc} \alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \dots & \alpha_{s+k-2} & \alpha_{s+k-1} \\ \hline \beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s-1} & \beta_s & \beta_{s+1} & \dots & \beta_{s+k-3} & \beta_{s+k-2} \\ \beta_s & \beta_{s+1} & \beta_{s+2} & \dots & \beta_{s+k-2} & \beta_{s+k-1} \\ \hline \gamma_1 & \gamma_2 & \gamma_3 & \dots & \gamma_{k-1} & \gamma_k \\ \gamma_2 & \gamma_3 & \gamma_4 & \dots & \gamma_k & \gamma_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{s-1} & \gamma_s & \gamma_{s+1} & \dots & \gamma_{s+k-3} & \gamma_{s+k-2} \\ \gamma_s & \gamma_{s+1} & \gamma_{s+2} & \dots & \gamma_{s+k-2} & \gamma_{s+k-1} \end{array} \right)$$

10.2. **Introduction.** We adapt the method of Sections 13, 14, 15, 16 and 17 in [2] to compute

$\Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k}$ for $s \leq i \leq \inf(3s, k)$ using the results obtained in Section 6 and Section 9.

10.3. **Computation of $\Gamma_{s+j}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k}$ for $j = 0, 1, 2$, $j \leq s-1$, $k \geq s+j$.**

We have respectively from (6.1) and (6.2) with $m = l = 0$

$$(10.1) \quad \sum_{i=0}^{\inf(3s, k)} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} = 2^{3k+3s-3}$$

and

$$(10.2) \quad \sum_{i=0}^{\inf(3s, k)} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} \cdot 2^{-i} = 2^{2k+3s-3} + 2^{3k-3} - 2^{2k-3}.$$

We get respectively from (6.39) and (6.40)

$$(10.3) \quad \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} = 7 \cdot 2^{4s-6} - 3 \cdot 2^{3s-5} \quad \text{if } k \geq s$$

and

$$(10.4) \quad \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} \cdot 2^{-i} = 15 \cdot 2^{3s-6} - 7 \cdot 2^{2s-5} \quad \text{if } k \geq s$$

Lemma 10.2. *We have*

$$(10.5) \quad \Gamma_s^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} = 7 \cdot 2^{k+s-1} - 7 \cdot 2^{2s} + 105 \cdot 2^{4s-6} - 21 \cdot 2^{3s-5} \quad \text{if } k \geq s+1, \quad s \geq 2$$

$$(10.6) \quad \Gamma_s^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times s} = 2^{6s-3} - 7 \cdot 2^{4s-6} + 3 \cdot 2^{3s-5} \quad \text{if } s \geq 2$$

$$(10.7) \quad \Gamma_{s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} = 147 \cdot 2^{k+s-1} + 105 \cdot 2^{4s-2} - 21 \cdot 2^{3s-2} - 315 \cdot 2^{2s} \quad \text{if } k \geq s+2, \quad s \geq 2$$

$$(10.8) \quad \Gamma_{s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+1)} = 2^{6s} - 7 \cdot 2^{4s-2} + 3 \cdot 2^{3s-2} \quad \text{if } s \geq 2$$

$$(10.9) \quad \Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} = 147 \cdot 9 \cdot 2^{k+s} + 21 \cdot [5 \cdot 2^{4s+2} - 2^{3s+1} - 275 \cdot 2^{2s+1}] \quad \text{if } k \geq s+3, \quad s \geq 3$$

$$(10.10) \quad \Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} = 2^{6s+3} - 7 \cdot 2^{4s+2} + 3 \cdot 2^{3s+1} + 7 \cdot 2^{2s+1} \quad \text{if } s \geq 2$$

$$(10.11) \quad \Gamma_{s+3}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+3)} = 2^{6s+6} - 7 \cdot 2^{4s+6} + 3 \cdot 2^{3s+4} + 21 \cdot 2^{2s+5} \quad \text{if } s \geq 3$$

Proof. Proof of (10.5)

By (6.42), we get

$$(10.12) \quad \Gamma_s^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+1)} = 105 \cdot 2^{4s-6} - 21 \cdot 2^{3s-5}$$

We have from (9.5) with $j=0$, $k > s$

$$(10.13) \quad \Delta_k \Gamma_s^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right]} = \Gamma_s^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (k+1)} - \Gamma_s^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} = 7 \cdot 2^{k+s-1}$$

We obtain from (10.12) and (10.13) for $k \geq s+1$

$$(10.14) \quad \begin{aligned} & \sum_{i=s+1}^k (\Gamma_s^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (i+1)} - \Gamma_s^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times i}) = \sum_{i=s+1}^k 7 \cdot 2^{k+s-1} \\ & \Downarrow \\ & \Gamma_s^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (k+1)} - \Gamma_s^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+1)} = 7 \cdot 2^{k+s} - 7 \cdot 2^{2s} \\ & \Downarrow \\ & \Gamma_s^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} = 7 \cdot 2^{k+s-1} - 7 \cdot 2^{2s} + 105 \cdot 2^{4s-6} - 21 \cdot 2^{3s-5} \quad \text{for } k \geq s+2 \end{aligned}$$

It follows from (10.12) that (10.14) is valid for $k \geq s+1$

proof of (10.6) By (6.41) with $m = l = 0, i = s$.

proof of (10.7) and (10.10) We get respectively from (10.1) and (10.2) with $k = s+2$

$$(10.15) \quad \sum_{i=0}^{s+2} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} = 2^{6s+3}$$

and

$$(10.16) \quad \sum_{i=0}^{s+2} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} \cdot 2^{-i} = 2^{5s+1} + 2^{3s+3} - 2^{2s+1}.$$

Applying (10.3) and (10.5) with $k = s+2$ we get from (10.15)

$$\begin{aligned} \sum_{i=0}^{s+2} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} &= \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} + \Gamma_s^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} + \Gamma_{s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} + \Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} \\ &= (7 \cdot 2^{4s-6} - 3 \cdot 2^{3s-5}) + (7 \cdot 2^{2s+1} - 7 \cdot 2^{2s} + 105 \cdot 2^{4s-6} - 21 \cdot 2^{3s-5}) + \Gamma_{s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} + \Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} = 2^{6s+3} \end{aligned}$$

Hence we obtain from the above equation

$$(10.17) \quad \Gamma_{s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} + \Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} = 2^{6s+3} - 7 \cdot 2^{4s-2} + 3 \cdot 2^{3s-2} - 7 \cdot 2^{2s}$$

Applying (10.4) and (10.5) with $k = s+2$ we get from (10.16)

$$\begin{aligned} \sum_{i=0}^{s+2} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} \cdot 2^{-i} &= \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} \cdot 2^{-i} + \Gamma_s^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} \cdot 2^{-s} + \Gamma_{s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} \cdot 2^{-(s+1)} + \Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} \cdot 2^{-(s+2)} \\ &= (15 \cdot 2^{3s-6} - 7 \cdot 2^{2s-5}) + (7 \cdot 2^s + 105 \cdot 2^{3s-6} - 21 \cdot 2^{2s-5}) \\ &\quad + \Gamma_{s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} \cdot 2^{-(s+1)} + \Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} \cdot 2^{-(s+2)} = 2^{5s+1} + 2^{3s+3} - 2^{2s+1} \end{aligned}$$

We then get from the above equation

$$\begin{aligned} &\Gamma_{s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} \cdot 2^{-(s+1)} + \Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} \cdot 2^{-(s+2)} = 2^{5s+1} + 49 \cdot 2^{3s-3} - 9 \cdot 2^{2s-3} - 7 \cdot 2^s \\ &\quad \updownarrow \\ (10.18) \quad &2 \cdot \Gamma_{s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} + \Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} = 2^{6s+3} + 49 \cdot 2^{4s-1} - 9 \cdot 2^{3s-1} - 7 \cdot 2^{2s+2} \end{aligned}$$

Now we deduce from (10.17) and (10.18)

$$(10.19) \quad \Gamma_{s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} = 105 \cdot 2^{4s-2} - 21 \cdot 2^{3s-2} - 21 \cdot 2^{2s}$$

$$(10.20) \quad \Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+2)} = 2^{6s+3} - 7 \cdot 2^{4s+2} + 3 \cdot 2^{3s+1} + 7 \cdot 2^{2s+1}$$

We have from (9.5) with $j = 1, k \geq s+2$

$$(10.21) \quad \Delta_k \Gamma_{s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right]} = \Gamma_{s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (k+1)} - \Gamma_{s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} = 147 \cdot 2^{k+s-3}$$

We obtain from (10.21) and (10.19) for $k \geq s+2$

$$\begin{aligned}
& \sum_{i=s+2}^k \left(\Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (i+1)} - \Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times i} \right) = \sum_{i=s+1}^k 147 \cdot 2^{k+s-3} \\
& \Updownarrow \\
& \Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (k+1)} - \Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+2)} = 147 \cdot 2^{k+s} - 147 \cdot 2^{2s+1} \\
& \Updownarrow \\
(10.22) \quad & \Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times k} = 147 \cdot 2^{k+s-1} - 147 \cdot 2^{2s+1} + [105 \cdot 2^{4s-2} - 21 \cdot 2^{3s-2} - 21 \cdot 2^{2s}] \quad \text{for } k \geq s+3
\end{aligned}$$

It follows from (10.19) that (10.22) is valid for $k \geq s+2$

proof of (10.6)
By (6.41) with $m = l = 0, i = s$.

Proof of (10.9) and (10.11) . We get respectively from (10.1) and (10.2) with $k = s+3$

$$(10.23) \quad \sum_{i=0}^{s+3} \Gamma_i^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+3)} = 2^{6s+6}$$

and

$$(10.24) \quad \sum_{i=0}^{s+3} \Gamma_i^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+3)} \cdot 2^{-i} = 2^{5s+3} + 2^{3s+6} - 2^{2s+3}.$$

Applying (10.3), (10.5) and (10.7) with $k = s+3$ we get from (10.23)

$$\begin{aligned}
\sum_{i=0}^{s+3} \Gamma_i^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+3)} &= \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+3)} + \Gamma_s^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+3)} + \Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+3)} + \Gamma_{s+2}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+3)} + \Gamma_{s+3}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+3)} \\
&= (7 \cdot 2^{4s-6} - 3 \cdot 2^{3s-5}) + (21 \cdot 2^{2s} + 105 \cdot 2^{4s-6} - 21 \cdot 2^{3s-5}) \\
&+ (147 \cdot 2^{2s+2} + 105 \cdot 2^{4s-2} - 21 \cdot 2^{3s-2} - 315 \cdot 2^{2s}) + \Gamma_{s+2}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+3)} + \Gamma_{s+3}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+3)} = 2^{6s+6}
\end{aligned}$$

Hence we obtain from the above equation

$$(10.25) \quad \Gamma_{s+2}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+3)} + \Gamma_{s+3}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+3)} = 2^{6s+6} - 7 \cdot 2^{4s+2} + 3 \cdot 2^{3s+1} - 147 \cdot 2^{2s+1}$$

Applying (10.4), (10.5) and (10.7) with $k = s+3$ we get from (10.24)

$$\begin{aligned}
\sum_{i=0}^{s+3} \Gamma_i^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+3)} \cdot 2^{-i} &= \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+3)} \cdot 2^{-i} + \Gamma_s^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+3)} \cdot 2^{-s} + \Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+3)} \cdot 2^{-(s+1)} \\
&+ \Gamma_{s+2}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+3)} \cdot 2^{-(s+2)} + \Gamma_{s+3}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+3)} \cdot 2^{-(s+3)} \\
&= (15 \cdot 2^{3s-6} - 7 \cdot 2^{2s-5}) + (21 \cdot 2^s + 105 \cdot 2^{3s-6} - 21 \cdot 2^{2s-5}) + (147 \cdot 2^{s+1} + 105 \cdot 2^{3s-3} - 21 \cdot 2^{2s-3} - 315 \cdot 2^{s-1}) \\
&+ \Gamma_{s+2}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+3)} \cdot 2^{-(s+2)} + \Gamma_{s+3}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (s+3)} \cdot 2^{-(s+3)} = 2^{5s+3} + 2^{3s+6} - 2^{2s+3}
\end{aligned}$$

We then get from the above equation

$$\begin{aligned}
 & \Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+3)} \cdot 2^{-(s+2)} + \Gamma_{s+3}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+3)} \cdot 2^{-(s+3)} = 2^{5s+3} + 49 \cdot 2^{3s} - 9 \cdot 2^{2s-1} - 315 \cdot 2^{s-1} \\
 & \Updownarrow \\
 (10.26) \quad & 2 \cdot \Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+3)} + \Gamma_{s+3}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+3)} = 2^{6s+6} + 49 \cdot 2^{4s+3} - 9 \cdot 2^{3s+2} - 315 \cdot 2^{2s+2}
 \end{aligned}$$

Now we deduce from (10.25) and (10.26)

$$(10.27) \quad \Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+3)} = 21 \cdot [5 \cdot 2^{4s+2} - 2^{3s+1} - 23 \cdot 2^{2s+1}]$$

$$(10.28) \quad \Gamma_{s+3}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+3)} = 2^{6s+6} - 7 \cdot 2^{4s+6} + 3 \cdot 2^{3s+4} + 21 \cdot 2^{2s+5}$$

We have from (9.5) with $j = 2$, $k \geq s + 3$

$$(10.29) \quad \Delta_k \Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (k+1)} = \Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (k+1)} - \Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} = 147 \cdot 9 \cdot 2^{k+s}$$

We obtain from (10.29) and (10.27) for $k \geq s + 3$

$$\begin{aligned}
 & \sum_{i=s+3}^k (\Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (i+1)} - \Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times i}) = \sum_{i=s+3}^k 147 \cdot 9 \cdot 2^{k+s} \\
 & \Updownarrow \\
 & \Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (k+1)} - \Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+3)} = 147 \cdot 9 \cdot 2^{k+s+1} - 147 \cdot 9 \cdot 2^{2s+3} \\
 & \Updownarrow \\
 (10.30) \quad & \Gamma_{s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} = 147 \cdot 9 \cdot 2^{k+s} + 21 \cdot [5 \cdot 2^{4s+2} - 2^{3s+1} - 275 \cdot 2^{2s+1}] \quad \text{for } k \geq s + 4
 \end{aligned}$$

It follows from (10.27) that (10.30) is valid for $k \geq s + 3$

□

10.4. An explicit formula for $\Gamma_{s+j}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k}$ for $1 \leq j \leq s - 1$, $k \geq s + j$.

Lemma 10.3. *We have*

$$\begin{aligned}
 (10.31) \quad & (H_j) \quad \Gamma_{s+j}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} = 147 \cdot (5 \cdot 2^{j-1} - 1) \cdot 2^{k+s+3j-6} \\
 & \quad + 21 \cdot [5 \cdot 2^{4s+4j-6} - 2^{3s+3j-5} - (155 \cdot 2^{j-1} - 35) \cdot 2^{2s+4j-7}] \quad \text{for } 1 \leq j \leq s - 1, k \geq s + j + 1. \\
 (10.32) \quad & \Gamma_{s+j}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j)} = 2^{6s+3j-3} + 7 \cdot 2^{2s+5j-8} - 7 \cdot 2^{2s+4j-7} - 7 \cdot 2^{4s+4j-6} + 3 \cdot 2^{3s+3j-5} \quad \text{for } 1 \leq j \leq s.
 \end{aligned}$$

Proof. The proof is by strong induction, that is:

If

- (H_1) is true, and
- for all $j \geq 1$, $(H_1) \wedge (H_2) \wedge \dots \wedge (H_j)$ implies (H_{j+1})

then (H_j) is true for all $j \geq 1$.

(H_1) is true

Indeed from (10.7) we have

$$\Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix}\right] \times k} = 147 \cdot 2^{k+s-1} + 105 \cdot 2^{4s-2} - 21 \cdot 2^{3s-2} - 315 \cdot 2^{2s} \quad \text{if } k \geq s+2, s \geq 2$$

and (H_j) with $j=1$ gives

$$147 \cdot (5-1) \cdot 2^{k+s-3} + 21 \cdot [5 \cdot 2^{4s-2} - 2^{3s-2} - (155-35) \cdot 2^{2s-3}] = \Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix}\right] \times k}$$

From (10.8) we see that $\Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix}\right] \times (s+1)}$ is equal to (10.32) with $j=1$.

(H_1) implies (H_2)

Recall the proof of (10.9).

$$(10.9) \quad \Gamma_{s+2}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix}\right] \times k} = 147 \cdot 9 \cdot 2^{k+s} + 21 \cdot [5 \cdot 2^{4s+2} - 2^{3s+1} - 275 \cdot 2^{2s+1}] \quad \text{if } k \geq s+3, s \geq 3.$$

(H_j) with $j=2$ gives

$$147 \cdot (5 \cdot 2 - 1) \cdot 2^{k+s} + 21 \cdot [5 \cdot 2^{4s+2} - 2^{3s+1} - (155 \cdot 2 - 35) \cdot 2^{2s+1}] = \Gamma_{s+2}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix}\right] \times k}$$

From (10.10) we see that $\Gamma_{s+2}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix}\right] \times (s+2)}$ is equal to (10.32) with $j=2$.

$(H_1) \wedge (H_2) \wedge \dots \wedge (H_j)$ implies (H_{j+1})

We get respectively from (10.1) and (10.2) with $k = s+j+2$

$$(10.33) \quad \sum_{i=0}^{s+j+2} \Gamma_i^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix}\right] \times (s+j+2)} = 2^{6s+3j+3}$$

and

$$(10.34) \quad \sum_{i=0}^{s+j+2} \Gamma_i^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix}\right] \times (s+j+2)} \cdot 2^{-i} = 2^{5s+2j+1} + 2^{3s+3j+3} - 2^{2s+2j+1}.$$

We have respectively from (10.3) and (10.4) with $k=s+j+2$

$$(10.35) \quad \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix}\right] \times (s+j+2)} = 7 \cdot 2^{4s-6} - 3 \cdot 2^{3s-5}$$

$$(10.36) \quad \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix}\right] \times (s+j+2)} \cdot 2^{-i} = 15 \cdot 2^{3s-6} - 7 \cdot 2^{2s-5}$$

By (10.5) with $k=s+j+2$

$$(10.37) \quad \Gamma_s^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix}\right] \times (s+j+2)} = 7 \cdot 2^{2s+j+1} - 7 \cdot 2^{2s} + 105 \cdot 2^{4s-6} - 21 \cdot 2^{3s-5}$$

$$(10.38) \quad \Gamma_s^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix}\right] \times (s+j+2)} \cdot 2^{-s} = 7 \cdot 2^{s+j+1} - 7 \cdot 2^s + 105 \cdot 2^{3s-6} - 21 \cdot 2^{2s-5}$$

We get from (H_q) with $k = s+j+2$ and $1 \leq q \leq j$

$$\Gamma_{s+q}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix}\right] \times (s+j+2)} = 147 \cdot (5 \cdot 2^{q-1} - 1) \cdot 2^{(s+j+2)+s+3q-6} + 21 \cdot [5 \cdot 2^{4s+4q-6} - 2^{3s+3q-5} - (155 \cdot 2^{q-1} - 35) \cdot 2^{2s+4q-7}]$$

$$\Updownarrow$$

(10.39)

$$\Gamma_{s+q}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} = 2^{5q} \cdot [-3255 \cdot 2^{2s-8}] + 2^{4q} \cdot [735 \cdot 2^{2s+j-5} + 105 \cdot 2^{4s-6} + 735 \cdot 2^{2s-7}] + 2^{3q} \cdot [-147 \cdot 2^{2s+j-4} - 21 \cdot 2^{3s-5}]$$

(10.40)

$$\Gamma_{s+q}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} \cdot 2^{-(s+q)} = 2^{4q} \cdot [-3255 \cdot 2^{s-8}] + 2^{3q} \cdot [735 \cdot 2^{s+j-5} + 105 \cdot 2^{3s-6} + 735 \cdot 2^{s-7}] + 2^{2q} \cdot [-147 \cdot 2^{s+j-4} - 21 \cdot 2^{2s-5}]$$

We obtain from (10.33), (10.35), (10.37) and (10.39).

$$\begin{aligned} \sum_{i=0}^{s+j+2} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} &= \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} + \Gamma_s^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} + \sum_{q=1}^j \Gamma_{s+q}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} + \Gamma_{s+j+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} + \Gamma_{s+j+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} \\ &= (7 \cdot 2^{4s-6} - 3 \cdot 2^{3s-5}) + (7 \cdot 2^{2s+j+1} - 7 \cdot 2^{2s} + 105 \cdot 2^{4s-6} - 21 \cdot 2^{3s-5}) \\ &\quad + [-3255 \cdot 2^{2s-8}] \cdot \sum_{q=1}^j 2^{5q} + [735 \cdot 2^{2s+j-5} + 105 \cdot 2^{4s-6} + 735 \cdot 2^{2s-7}] \cdot \sum_{q=1}^j 2^{4q} + [-147 \cdot 2^{2s+j-4} - 21 \cdot 2^{3s-5}] \cdot \sum_{q=1}^j 2^{3q} \\ &\quad + \Gamma_{s+j+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} + \Gamma_{s+j+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} = 2^{6s+3j+3} \end{aligned}$$

$$\Updownarrow$$

(10.41)

$$\Gamma_{s+j+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} + \Gamma_{s+j+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} = 2^{6s+3j+3} - 91 \cdot 2^{2s+5j-3} + 35 \cdot 2^{2s+4j-3} - 7 \cdot 2^{4s+4j-2} + 3 \cdot 2^{3s+3j-2}$$

We get in the same way from (10.34), (10.36), (10.38) and (10.40).

$$\begin{aligned} \sum_{i=0}^{s+j+2} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} \cdot 2^{-i} &= \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} \cdot 2^{-i} + \Gamma_s^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} \cdot 2^{-s} + \sum_{q=1}^j \Gamma_{s+q}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} \cdot 2^{-(s+q)} \\ &\quad + \Gamma_{s+j+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} \cdot 2^{-(s+j+1)} + \Gamma_{s+j+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} \cdot 2^{-(s+j+2)} \\ &= (15 \cdot 2^{3s-6} - 7 \cdot 2^{2s-5}) + (7 \cdot 2^{s+j+1} - 7 \cdot 2^s + 105 \cdot 2^{3s-6} - 21 \cdot 2^{2s-5}) \\ &\quad + [-3255 \cdot 2^{s-8}] \cdot \sum_{q=1}^j 2^{4q} + [735 \cdot 2^{s+j-5} + 105 \cdot 2^{3s-6} + 735 \cdot 2^{s-7}] \cdot \sum_{q=1}^j 2^{3q} + [-147 \cdot 2^{s+j-4} - 21 \cdot 2^{2s-5}] \cdot \sum_{q=1}^j 2^{2q} \\ &\quad + \Gamma_{s+j+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} \cdot 2^{-(s+j+1)} + \Gamma_{s+j+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} \cdot 2^{-(s+j+2)} = 2^{5s+2j+1} + 2^{3s+3j+3} - 2^{2s+2j+1} \end{aligned}$$

$$\Updownarrow$$

$$\begin{aligned} \Gamma_{s+j+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} \cdot 2^{-(s+j+1)} + \Gamma_{s+j+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} \cdot 2^{-(s+j+2)} \\ = 2^{5s+2j+1} + 49 \cdot 2^{3s+3j-3} - 203 \cdot 2^{s+4j-4} + 91 \cdot 2^{s+3j-4} - 9 \cdot 2^{2s+2j-3} \end{aligned}$$

$$\Updownarrow$$

(10.42)

$$2 \cdot \Gamma_{s+j+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} + \Gamma_{s+j+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} = 2^{6s+3j+3} + 49 \cdot 2^{4s+4j-1} - 203 \cdot 2^{2s+5j-2} + 91 \cdot 2^{2s+4j-2} - 9 \cdot 2^{3s+3j-1}$$

Hence by (10.41), (10.42) we deduce

$$(10.43) \quad \Gamma_{s+j+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (s+j+2)} = 105 \cdot 2^{4s+4j-2} - 315 \cdot 2^{2s+5j-3} + 147 \cdot 2^{2s+4j-3} - 21 \cdot 2^{3s+3j-2}$$

$$(10.44) \quad \Gamma_{s+j+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (s+j+2)} = 2^{6s+3j+3} + 7 \cdot 2^{2s+5j+2} - 7 \cdot 2^{2s+4j+1} - 7 \cdot 2^{4s+4j+2} + 3 \cdot 2^{3s+3j+1}$$

We have from (9.5) with $j \rightarrow j+1$, $j+1 \leq s-1$, $k \geq s+j+2$

$$(10.45) \quad \Delta_k \Gamma_{s+j+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = \Gamma_{s+j+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (k+1)} - \Gamma_{s+j+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times k} = 147 \cdot (5 \cdot 2^j - 1) \cdot 2^{k+s+3j-3}$$

We obtain from (10.45) and (10.43) for $k \geq s+j+2$

$$\begin{aligned} \sum_{i=s+j+2}^k (\Gamma_{s+j+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (i+1)} - \Gamma_{s+j+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times i}) &= \sum_{i=s+j+2}^k 147 \cdot (5 \cdot 2^j - 1) \cdot 2^{k+s+3j-3} \\ \Downarrow \\ \Gamma_{s+j+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (k+1)} - \Gamma_{s+j+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (s+j+2)} &= 147 \cdot (5 \cdot 2^j - 1) \cdot 2^{k+s+3j-2} - 147 \cdot (5 \cdot 2^j - 1) \cdot 2^{2s+4j-1} \\ \Downarrow \\ (10.46) \quad \Gamma_{s+j+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times k} &= 147 \cdot (5 \cdot 2^j - 1) \cdot 2^{k+s+3j-3} - 147 \cdot (5 \cdot 2^j - 1) \cdot 2^{2s+4j-1} \\ &\quad + 105 \cdot 2^{4s+4j-2} - 315 \cdot 2^{2s+5j-3} + 147 \cdot 2^{2s+4j-3} - 21 \cdot 2^{3s+3j-2} \quad \text{for } k \geq s+j+3 \end{aligned}$$

It follows from (10.43) that (10.46) is valid for $k \geq s+j+2$

By (10.31) we have with $j \rightarrow j+1$

$$\begin{aligned} (10.47) \quad (H_{j+1}) \quad \Gamma_{s+j+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times k} &= 147 \cdot (5 \cdot 2^j - 1) \cdot 2^{k+s+3j-3} \\ &\quad + 21 \cdot [5 \cdot 2^{4s+4j-2} - 2^{3s+3j-2} - (155 \cdot 2^j - 35) \cdot 2^{2s+4j-3}] \quad \text{for } 1 \leq j+1 \leq s-1, k \geq s+j+2. \end{aligned}$$

By comparing (10.46) with (10.47) we see that (H_{j+1}) holds. $\square$

10.5. Explicit formulas for $\Gamma_{2s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times k}$ where $k \geq 2s+1$ and for $\Gamma_{2s+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+1)}$.

Lemma 10.4. *We have*

$$(10.48) \quad \Gamma_{2s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times k} = 7 \cdot 2^{2k+2s-2} + 21 \cdot [35 \cdot 2^{k+5s-7} - 39 \cdot 2^{k+4s-6}] + 7 \cdot [15 \cdot 2^{8s-6} - 465 \cdot 2^{7s-8} + 349 \cdot 2^{6s-7}] \quad \text{for } k \geq 2s+1.$$

$$(10.49) \quad \Gamma_{2s+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+1)} = 2^{9s} - 7 \cdot 2^{8s-2} + 7 \cdot 2^{7s-3} - 2^{6s-3}$$

Proof. We get respectively from (10.43) and (10.44) with $j = s-1$

$$(10.50) \quad \Gamma_{2s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+1)} = 105 \cdot 2^{8s-6} - 315 \cdot 2^{7s-8} + 63 \cdot 2^{6s-7}$$

$$(10.51) \quad \Gamma_{2s+1}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+1)} = 2^{9s} + 7 \cdot 2^{7s-3} - 7 \cdot 2^{8s-2} - 2^{6s-3}$$

We have from (9.7) with $j = 0$ and $k \geq 2s+1$

$$(10.52) \quad \Delta_k \Gamma_{2s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = \Gamma_{2s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (k+1)} - \Gamma_{2s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times k} = 21 \cdot (2^{2k+2s-2} + 35 \cdot 2^{k+5s-7} - 39 \cdot 2^{k+4s-6})$$

We obtain from (10.52) for $k \geq 2s+1$ and (10.50)

$$\sum_{i=2s+1}^k \left(\Gamma_{2s}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (i+1)} - \Gamma_{2s}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times i} \right) = \sum_{i=2s+1}^k 21 \cdot (2^{2i+2s-2} + 35 \cdot 2^{i+5s-7} - 39 \cdot 2^{i+4s-6})$$

$\Downarrow$

$$\begin{aligned} \Gamma_{2s}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (k+1)} &= 21 \cdot 2^{2s-2} \cdot \sum_{i=2s+1}^k 2^{2i} + (735 \cdot 2^{5s-7} - 819 \cdot 2^{4s-6}) \cdot \sum_{i=2s+1}^k 2^i + 105 \cdot 2^{8s-6} - 315 \cdot 2^{7s-8} + 63 \cdot 2^{6s-7} \\ &= 7 \cdot 2^{2k+2s} + 735 \cdot 2^{k+5s-6} - 819 \cdot 2^{k+4s-5} + 105 \cdot 2^{8s-6} - 3255 \cdot 2^{7s-8} + 2443 \cdot 2^{6s-7} \end{aligned}$$

$\Downarrow$

(10.53)

$$\Gamma_{2s}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} = 7 \cdot 2^{2k+2s-2} + 21 \cdot [35 \cdot 2^{k+5s-7} - 39 \cdot 2^{k+4s-6}] + 7 \cdot [15 \cdot 2^{8s-6} - 465 \cdot 2^{7s-8} + 349 \cdot 2^{6s-7}] \quad \text{for } k \geq 2s+2$$

From (10.51) we see that (10.53) is valid for $k = 2s+1$.

□

10.6. Explicit formulas for $\Gamma_{2s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k}$ where $k \geq 2s+2$ and for $\Gamma_{2s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)}$.

Lemma 10.5. *We have*

$$\begin{aligned} \Gamma_{2s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} &= 105 \cdot (2^{2k+2s-2} + 7 \cdot 2^{k+5s-3} - 31 \cdot 2^{k+4s-3}) \\ &\quad + 105 \cdot (2^{8s-2} - 31 \cdot 2^{7s-3} + 93 \cdot 2^{6s-3}) \quad \text{for } k \geq 2s+2. \end{aligned} \quad (10.54)$$

$$\Gamma_{2s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} = 2^{9s+3} - 7 \cdot 2^{8s+2} + 7 \cdot 2^{7s+2} - 2^{6s+3} \quad (10.55)$$

Proof. We get respectively from (10.1) and (10.2) for $k = 2s+2$

$$\sum_{i=0}^{2s+2} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} = 2^{9s+3} \quad (10.56)$$

and

$$\sum_{i=0}^{2s+2} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} \cdot 2^{-i} = 2^{7s+1} + 2^{6s+3} - 2^{4s+1}. \quad (10.57)$$

We have from (10.3), (10.4) with $k = 2s+2$

$$\sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} = 7 \cdot 2^{4s-6} - 3 \cdot 2^{3s-5} \quad (10.58)$$

$$\sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} \cdot 2^{-i} = 15 \cdot 2^{3s-6} - 7 \cdot 2^{2s-5} \quad (10.59)$$

By (10.5) with $k=2s+2$

$$\Gamma_s^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} = 427 \cdot 2^{3s-5} - 7 \cdot 2^{2s} + 105 \cdot 2^{4s-6} \quad (10.60)$$

$$\Gamma_s^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} \cdot 2^{-s} = 427 \cdot 2^{2s-5} - 7 \cdot 2^s + 105 \cdot 2^{3s-6} \quad (10.61)$$

We obtain from (10.31) with $j \rightarrow q$ and $k \geq 2s$

$$\begin{aligned}
 (10.62) \quad & \sum_{q=1}^{s-1} \Gamma_{s+q}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times k} \\
 &= \sum_{q=1}^{s-1} 2^{4q} [735 \cdot 2^{k+s-7} + 105 \cdot 2^{4s-6} + 735 \cdot 2^{2s-7}] \\
 &+ \sum_{q=1}^{s-1} 2^{3q} [-147 \cdot 2^{k+s-6} - 21 \cdot 2^{3s-5}] + \sum_{q=1}^{s-1} 2^{5q} [-3255 \cdot 2^{2s-8}] \\
 &= 49 \cdot 2^{k+5s-7} - 7 \cdot 2^{k+s-1} - 21 \cdot 2^{k+4s-6} + 7 \cdot 2^{8s-6} + 37 \cdot 2^{6s-7} - 7 \cdot 2^{4s-2} - 105 \cdot 2^{7s-8} + 3 \cdot 2^{3s-2} + 7 \cdot 2^{2s}
 \end{aligned}$$

$$\begin{aligned}
 (10.63) \quad & \sum_{q=1}^{s-1} \Gamma_{s+q}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times k} \cdot 2^{-(s+q)} \\
 &= \sum_{q=1}^{s-1} 2^{3q} [735 \cdot 2^{k-7} + 105 \cdot 2^{3s-6} + 735 \cdot 2^{s-7}] \\
 &+ \sum_{q=1}^{s-1} 2^{2q} [-147 \cdot 2^{k-6} - 21 \cdot 2^{2s-5}] + \sum_{q=1}^{s-1} 2^{4q} [-3255 \cdot 2^{s-8}] \\
 &= 105 \cdot 2^{k+3s-7} - 49 \cdot 2^{k+2s-6} - 7 \cdot 2^{k-1} - 217 \cdot 2^{5s-8} + 15 \cdot 2^{6s-6} + 77 \cdot 2^{4s-7} - 15 \cdot 2^{3s-3} + 7 \cdot 2^{2s-3} + 7 \cdot 2^s
 \end{aligned}$$

From (10.62) and (10.63) with $k = 2k+2$ we obtain

$$\begin{aligned}
 (10.64) \quad & \sum_{q=1}^{s-1} \Gamma_{s+q}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+2)} \\
 &= 287 \cdot 2^{7s-8} - 53 \cdot 2^{3s-2} - 131 \cdot 2^{6s-7} + 7 \cdot 2^{8s-6} - 7 \cdot 2^{4s-2} + 7 \cdot 2^{2s}
 \end{aligned}$$

$$\begin{aligned}
 (10.65) \quad & \sum_{q=1}^{s-1} \Gamma_{s+q}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+2)} \cdot 2^{-(s+q)} \\
 &= 623 \cdot 2^{5s-8} + 15 \cdot 2^{6s-6} - 315 \cdot 2^{4s-7} - 105 \cdot 2^{2s-3} - 15 \cdot 2^{3s-3} + 7 \cdot 2^s
 \end{aligned}$$

We get from (10.48) with $k = 2s+2$

$$(10.66) \quad \Gamma_{2s}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+2)} = 105 \cdot 2^{8s-6} + 2625 \cdot 2^{7s-8} - 525 \cdot 2^{6s-7}$$

$$(10.67) \quad \Gamma_{2s}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+2)} \cdot 2^{-2s} = 105 \cdot 2^{6s-6} + 2625 \cdot 2^{5s-8} - 525 \cdot 2^{4s-7}$$

We obtain from (10.56), (10.58), (10.60), (10.64) and (10.66).

$$\begin{aligned}
 \sum_{i=0}^{2s+2} \Gamma_i^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+2)} &= \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+2)} + \Gamma_s^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+2)} + \sum_{q=1}^{s-1} \Gamma_{s+q}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+2)} + \Gamma_{2s}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+2)} + \Gamma_{2s+1}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+2)} + \Gamma_{2s+2}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+2)} \\
 &= (7 \cdot 2^{4s-6} - 3 \cdot 2^{3s-5}) + (427 \cdot 2^{3s-5} - 7 \cdot 2^{2s} + 105 \cdot 2^{4s-6}) \\
 &+ (287 \cdot 2^{7s-8} - 53 \cdot 2^{3s-2} - 131 \cdot 2^{6s-7} + 7 \cdot 2^{8s-6} - 7 \cdot 2^{4s-2} + 7 \cdot 2^{2s}) \\
 &+ (105 \cdot 2^{8s-6} + 2625 \cdot 2^{7s-8} - 525 \cdot 2^{6s-7}) + \Gamma_{2s+1}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+2)} + \Gamma_{2s+2}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+2)} = 2^{9s+3} \\
 &\updownarrow
 \end{aligned}$$

$$(10.68) \quad \Gamma_{2s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} + \Gamma_{2s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} = 2^{9s+3} - 7 \cdot 2^{8s-2} - 91 \cdot 2^{7s-3} + 41 \cdot 2^{6s-3}$$

We get in the same way from (10.57), (10.59), (10.61), (10.65) and (10.67)

$$\begin{aligned} \sum_{i=0}^{2s+2} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} \cdot 2^{-i} &= \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} \cdot 2^{-i} + \Gamma_s^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} \cdot 2^{-s} + \sum_{q=1}^{s-1} \Gamma_{s+q}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} \cdot 2^{-(s+q)} \\ &+ \Gamma_{2s}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} \cdot 2^{-2s} + \Gamma_{2s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} \cdot 2^{-(2s+1)} + \Gamma_{2s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} \cdot 2^{-(2s+2)} \\ &= (15 \cdot 2^{3s-6} - 7 \cdot 2^{2s-5}) + (427 \cdot 2^{2s-5} - 7 \cdot 2^s + 105 \cdot 2^{3s-6}) \\ &+ (623 \cdot 2^{5s-8} + 15 \cdot 2^{6s-6} - 315 \cdot 2^{4s-7} - 105 \cdot 2^{2s-3} - 15 \cdot 2^{3s-3} + 7 \cdot 2^s) \\ &+ (105 \cdot 2^{6s-6} + 2625 \cdot 2^{5s-8} - 525 \cdot 2^{4s-7}) + \Gamma_{2s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} \cdot 2^{-(2s+1)} + \Gamma_{2s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} \cdot 2^{-(2s+2)} = 2^{7s+1} + 2^{6s+3} - 2^{4s+1} \\ &\Updownarrow \end{aligned}$$

$$(10.69) \quad 2 \cdot \Gamma_{2s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} + \Gamma_{2s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} = 2^{9s+3} + 49 \cdot 2^{8s-1} - 203 \cdot 2^{7s-2} + 73 \cdot 2^{6s-2}$$

Hence by (10.68), (10.69)

$$(10.70) \quad \Gamma_{2s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} = 105 \cdot (2^{8s-2} - 3 \cdot 2^{7s-3} + 2^{6s-3})$$

$$(10.71) \quad \Gamma_{2s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} = 2^{9s+3} - 7 \cdot 2^{8s+2} + 7 \cdot 2^{7s+2} - 2^{6s+3}$$

We have from (9.7) with $j = 1$ and $k \geq 2s + 2$

$$(10.72) \quad \Delta_k \Gamma_{2s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right]} = \Gamma_{2s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (k+1)} - \Gamma_{2s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} = 105 \cdot (3 \cdot 2^{2k+2s-2} + 7 \cdot 2^{k+5s-3} - 31 \cdot 2^{k+4s-3})$$

We obtain from (10.72) for $k \geq 2s + 2$ and (10.70)

$$\begin{aligned} \sum_{i=2s+2}^k (\Gamma_{2s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (i+1)} - \Gamma_{2s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times i}) &= \sum_{i=2s+2}^k 105 \cdot (3 \cdot 2^{2i+2s-2} + 7 \cdot 2^{i+5s-3} - 31 \cdot 2^{i+4s-3}) \\ &\Updownarrow \\ \Gamma_{2s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (k+1)} - \Gamma_{2s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+2)} &= 315 \cdot 2^{2s-2} \cdot \sum_{i=2s+2}^k 2^{2i} + (735 \cdot 2^{5s-3} - 3255 \cdot 2^{4s-3}) \cdot \sum_{i=2s+2}^k 2^i \\ &= 105 \cdot 2^{2k+2s} + 735 \cdot 2^{k+5s-2} - 3255 \cdot 2^{k+4s-2} - 735 \cdot 2^{7s-1} + 2415 \cdot 2^{6s-1} \\ &\Updownarrow \\ \Gamma_{2s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} &= 105 \cdot 2^{2k+2s-2} + 735 \cdot 2^{k+5s-3} - 3255 \cdot 2^{k+4s-3} - 735 \cdot 2^{7s-1} + 2415 \cdot 2^{6s-1} + 105 \cdot (2^{8s-2} - 3 \cdot 2^{7s-3} + 2^{6s-3}) \\ &\Updownarrow \\ (10.73) \quad \Gamma_{2s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} &= 105 \cdot (2^{2k+2s-2} + 7 \cdot 2^{k+5s-3} - 31 \cdot 2^{k+4s-3}) + 105 \cdot (2^{8s-2} - 31 \cdot 2^{7s-3} + 93 \cdot 2^{6s-3}) \quad \text{for } k \geq 2s + 3 \end{aligned}$$

From (10.70) we see that (10.73) is valid for $k = 2s+2$.

□

10.7. **Explicit formulas for** $\Gamma_{2s+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times k}$ *where* $k \geq 2s+3$ *and for* $\Gamma_{2s+3}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+3)}$.

Lemma 10.6. *We have*

$$(10.74) \quad \Gamma_{2s+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times k} = 105 \cdot (2^{2k+2s+2} + 7 \cdot 2^{k+5s+1} - 31 \cdot 2^{k+4s+2}) \\ + 105 \cdot (2^{8s+2} - 31 \cdot 2^{7s+2} + 93 \cdot 2^{6s+3}) \quad \text{for } k \geq 2s+3.$$

$$(10.75) \quad \Gamma_{2s+3}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+3)} = 2^{9s+6} - 7 \cdot 2^{8s+6} + 7 \cdot 2^{7s+7} - 2^{6s+9}$$

$$(10.76) \quad \Gamma_{2s+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times k} = 16^3 \cdot \Gamma_{2(s-1)+1}^{\begin{bmatrix} s-1 \\ s-1 \\ s-1 \end{bmatrix} \times (k-3)} \quad \text{for } k \geq 2s+2.$$

Proof. We get respectively from (10.1) and (10.2) for $k = 2s+3$

$$(10.77) \quad \sum_{i=0}^{2s+3} \Gamma_i^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+3)} = 2^{9s+6}$$

and

$$(10.78) \quad \sum_{i=0}^{2s+3} \Gamma_i^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+3)} \cdot 2^{-i} = 2^{7s+3} + 2^{6s+6} - 2^{4s+3}.$$

We have from (10.3), (10.4) with $k = 2s+3$

$$(10.79) \quad \sum_{i=0}^{s-1} \Gamma_i^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+3)} = 7 \cdot 2^{4s-6} - 3 \cdot 2^{3s-5}$$

$$(10.80) \quad \sum_{i=0}^{s-1} \Gamma_i^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+3)} \cdot 2^{-i} = 15 \cdot 2^{3s-6} - 7 \cdot 2^{2s-5}$$

By (10.5) with $k=2s+3$

$$(10.81) \quad \Gamma_s^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+3)} = 875 \cdot 2^{3s-5} - 7 \cdot 2^{2s} + 105 \cdot 2^{4s-6}$$

$$(10.82) \quad \Gamma_s^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+3)} \cdot 2^{-s} = 875 \cdot 2^{2s-5} - 7 \cdot 2^s + 105 \cdot 2^{3s-6}$$

From (10.62) and (10.63) with $k = 2k+3$ we obtain

$$(10.83) \quad \sum_{q=1}^{s-1} \Gamma_{s+q}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+3)} \\ = 7 \cdot 2^{8s-6} + 679 \cdot 2^{7s-8} - 299 \cdot 2^{6s-7} - 7 \cdot 2^{4s-2} - 109 \cdot 2^{3s-2} + 7 \cdot 2^{2s}$$

$$(10.84) \quad \sum_{q=1}^{s-1} \Gamma_{s+q}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+3)} \cdot 2^{-(s+q)} \\ = 1463 \cdot 2^{5s-8} + 15 \cdot 2^{6s-6} - 707 \cdot 2^{4s-7} - 15 \cdot 2^{3s-3} - 217 \cdot 2^{2s-3} + 7 \cdot 2^s$$

We get from (10.48) with $k = 2s+3$

$$(10.85) \quad \Gamma_{2s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+3)} = 105 \cdot 2^{8s-6} + 8505 \cdot 2^{7s-8} + 3675 \cdot 2^{6s-7}$$

$$(10.86) \quad \Gamma_{2s}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} \cdot 2^{-2s} = 105 \cdot 2^{6s-6} + 8505 \cdot 2^{5s-8} + 3675 \cdot 2^{4s-7}$$

We get from (10.54) with $k = 2s+3$

$$(10.87) \quad \Gamma_{2s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} = 105 \cdot 2^{8s-2} + 2625 \cdot 2^{7s-3} - 2835 \cdot 2^{6s-3}$$

$$(10.88) \quad \Gamma_{2s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} \cdot 2^{-(2s+1)} = 105 \cdot 2^{6s-3} + 2625 \cdot 2^{5s-4} - 2835 \cdot 2^{4s-4}$$

We obtain from (10.77), (10.79), (10.81), (10.83), (10.85) and (10.87).

$$\begin{aligned} \sum_{i=0}^{2s+3} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} &= \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} + \Gamma_s^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} + \sum_{q=1}^{s-1} \Gamma_{s+q}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} \\ &+ \Gamma_{2s}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} + \Gamma_{2s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} + \Gamma_{2s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} + \Gamma_{2s+3}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} \\ &= (7 \cdot 2^{4s-6} - 3 \cdot 2^{3s-5}) + (875 \cdot 2^{3s-5} - 7 \cdot 2^{2s} + 105 \cdot 2^{4s-6}) \\ &+ (7 \cdot 2^{8s-6} + 679 \cdot 2^{7s-8} - 299 \cdot 2^{6s-7} - 7 \cdot 2^{4s-2} - 109 \cdot 2^{3s-2} + 7 \cdot 2^{2s}) \\ &+ (105 \cdot 2^{8s-6} + 8505 \cdot 2^{7s-8} + 3675 \cdot 2^{6s-7}) + (105 \cdot 2^{8s-2} + 2625 \cdot 2^{7s-3} - 2835 \cdot 2^{6s-3}) \\ &+ \Gamma_{2s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} + \Gamma_{2s+3}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} = 2^{9s+6} \\ &\Updownarrow \end{aligned}$$

$$(10.89) \quad \Gamma_{2s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} + \Gamma_{2s+3}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} = 2^{9s+6} - 7 \cdot 2^{8s+2} - 91 \cdot 2^{7s+2} + 41 \cdot 2^{6s+3}$$

We get in the same way from (10.78), (10.80), (10.82), (10.84), (10.86) and (10.88).

$$\begin{aligned} \sum_{i=0}^{2s+3} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} \cdot 2^{-i} &= \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} \cdot 2^{-i} + \Gamma_s^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} \cdot 2^{-s} + \sum_{q=1}^{s-1} \Gamma_{s+q}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} \cdot 2^{-(s+q)} \\ &+ \Gamma_{2s}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} \cdot 2^{-2s} + \Gamma_{2s+1}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} \cdot 2^{-(2s+1)} + \Gamma_{2s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} \cdot 2^{-(2s+2)} + \Gamma_{2s+3}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} \cdot 2^{-(2s+3)} \\ &= (15 \cdot 2^{3s-6} - 7 \cdot 2^{2s-5}) + (875 \cdot 2^{2s-5} - 7 \cdot 2^s + 105 \cdot 2^{3s-6}) \\ &+ (1463 \cdot 2^{5s-8} + 15 \cdot 2^{6s-6} - 707 \cdot 2^{4s-7} - 15 \cdot 2^{3s-3} - 217 \cdot 2^{2s-3} + 7 \cdot 2^s) \\ &+ (105 \cdot 2^{6s-6} + 8505 \cdot 2^{5s-8} + 3675 \cdot 2^{4s-7}) + (105 \cdot 2^{6s-3} + 2625 \cdot 2^{5s-4} - 2835 \cdot 2^{4s-4}) + \Gamma_{2s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} \cdot 2^{-(2s+2)} \\ &+ \Gamma_{2s+3}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} \cdot 2^{-(2s+3)} = 2^{7s+3} + 2^{6s+6} - 2^{4s+3} \\ &\Updownarrow \end{aligned}$$

$$(10.90) \quad 2 \cdot \Gamma_{2s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} + \Gamma_{2s+3}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} = 2^{9s+6} + 49 \cdot 2^{8s+3} - 203 \cdot 2^{7s+3} + 73 \cdot 2^{6s+4}$$

Hence by (10.89), (10.90)

$$(10.91) \quad \Gamma_{2s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} = 105 \cdot (2^{8s+2} - 3 \cdot 2^{7s+2} + 2^{6s+3})$$

$$(10.92) \quad \Gamma_{2s+3}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+3)} = 2^{9s+6} - 7 \cdot 2^{8s+6} + 7 \cdot 2^{7s+7} - 2^{6s+9}$$

We have from (9.7) with $j = 2$ and $k \geq 2s+3$

$$(10.93) \quad \Delta_k \Gamma_{2s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right]} = \Gamma_{2s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (k+1)} - \Gamma_{2s+2}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} = 105 \cdot (3 \cdot 2^{2k+2s+2} + 7 \cdot 2^{k+5s+1} - 31 \cdot 2^{k+4s+2})$$

We obtain from (10.93) for $k \geq 2s+3$ and (10.91)

$$\begin{aligned}
& \sum_{i=2s+3}^k \left(\Gamma_{2s+2}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (i+1)} - \Gamma_{2s+2}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times i} \right) = \sum_{i=2s+3}^k 105 \cdot (3 \cdot 2^{2i+2s+2} + 7 \cdot 2^{i+5s+1} - 31 \cdot 2^{i+4s+2}) \\
& \Downarrow \\
& \Gamma_{2s+2}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (k+1)} - \Gamma_{2s+2}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+3)} = 315 \cdot 2^{2s+2} \cdot \sum_{i=2s+3}^k 2^{2i} + (735 \cdot 2^{5s+1} - 3255 \cdot 2^{4s+2}) \cdot \sum_{i=2s+3}^k 2^i \\
& = 105 \cdot 2^{2k+2s+4} + 735 \cdot 2^{k+5s+2} - 3255 \cdot 2^{k+4s+3} - 735 \cdot 2^{7s+4} + 2415 \cdot 2^{6s+5} \\
& \Downarrow \\
& \Gamma_{2s+2}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (k+1)} = 105 \cdot 2^{2k+2s+4} + 735 \cdot 2^{k+5s+2} - 3255 \cdot 2^{k+4s+3} - 735 \cdot 2^{7s+4} + 2415 \cdot 2^{6s+5} + 105 \cdot (2^{8s+2} - 3 \cdot 2^{7s+2} + 2^{6s+3}) \\
& \Downarrow \\
& (10.94) \quad \Gamma_{2s+2}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times k} = 105 \cdot (2^{2k+2s+2} + 7 \cdot 2^{k+5s+1} - 31 \cdot 2^{k+4s+2}) + 105 \cdot (2^{8s+2} - 31 \cdot 2^{7s+2} + 93 \cdot 2^{6s+3}) \quad \text{for } k \geq 2s+4
\end{aligned}$$

From (10.91) we see that (10.94) is valid for $k = 2s+3$. It remains to prove (10.76). We have respectively from (10.49) with $s \rightarrow s-1$ and (10.55)

$$\begin{aligned}
16^3 \cdot \Gamma_{2(s-1)+1}^{\left[\begin{smallmatrix} s-1 \\ s-1 \\ s-1 \end{smallmatrix} \right] \times (2(s-1)+1)} &= 2^{12} \cdot (2^{9(s-1)} - 7 \cdot 2^{8(s-1)-2} + 7 \cdot 2^{7(s-1)-3} - 2^{6(s-1)-3}) \\
&= 2^{9s+3} - 7 \cdot 2^{8s+2} + 7 \cdot 2^{7s+2} - 2^{6s+3} = \Gamma_{2s+3}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+3)}
\end{aligned}$$

We get respectively from (10.54) with $s \rightarrow s-1$, $k \rightarrow k-3$, $k \geq 2s+3$ and (10.74)

$$\begin{aligned}
16^3 \cdot \Gamma_{2(s-1)+1}^{\left[\begin{smallmatrix} s-1 \\ s-1 \\ s-1 \end{smallmatrix} \right] \times (k-3)} &= 2^{12} \cdot 105 \cdot (2^{2(k-3)+2(s-1)-2} + 7 \cdot 2^{k-3+5(s-1)-3} - 31 \cdot 2^{k-3+4(s-1)-3}) \\
&+ 2^{12} \cdot 105 \cdot (2^{8(s-1)-2} - 31 \cdot 2^{7(s-1)-3} + 93 \cdot 2^{6(s-1)-3}) \\
&= 105 \cdot (2^{2k+2s+2} + 7 \cdot 2^{k+5s+1} - 31 \cdot 2^{k+4s+2}) + 105 \cdot (2^{8s+2} - 31 \cdot 2^{7s+2} + 93 \cdot 2^{6s+3}) = \Gamma_{2s+2}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times k}
\end{aligned}$$

□

10.8. Un explicit reduction formula for $\Gamma_{2s+1+j}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times k}$ for $0 \leq j \leq s-1$, $k \geq 2s+1+j$.

Lemma 10.7. *We have for $0 \leq j \leq s-2$, $k \geq 2s+2+j$*

(10.95)

$$\begin{aligned}
(H_j) \quad \Gamma_{2s+1+j}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times k} &= 16^{3j} \cdot \Gamma_{2(s-j)+1}^{\left[\begin{smallmatrix} s-j \\ s-j \\ s-j \end{smallmatrix} \right] \times (k-3j)} \\
&= 2^{12j} \cdot 105 \cdot (2^{2(k-3j)+2(s-j)-2} + 7 \cdot 2^{k-3j+5(s-j)-3} - 31 \cdot 2^{k-3j+4(s-j)-3}) \\
&+ 2^{12j} \cdot 105 \cdot (2^{8(s-j)-2} - 31 \cdot 2^{7(s-j)-3} + 93 \cdot 2^{6(s-j)-3}) \\
&= 105 \cdot (2^{2k+2s+4j-2} + 7 \cdot 2^{k+5s+4j-3} - 31 \cdot 2^{k+4s+5j-3}) \\
&+ 105 \cdot (2^{8s+4j-2} - 31 \cdot 2^{7s+5j-3} + 93 \cdot 2^{6s+6j-3})
\end{aligned}$$

Lemma 10.8. *We have for $0 \leq j \leq s-1$.*

$$\begin{aligned}
 (10.96) \quad \Gamma_{2s+1+j}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+1+j)} &= 16^{3j} \cdot \Gamma_{2(s-j)+1}^{\left[\begin{smallmatrix} s-j \\ s-j \\ s-j \end{smallmatrix} \right] \times (2(s-j)+1)} \\
 &= 2^{12j} \cdot (2^{9(s-j)} - 7 \cdot 2^{8(s-j)-2} + 7 \cdot 2^{7(s-j)-3} - 2^{6(s-j)-3}) \\
 &= 2^{9s+3j} - 7 \cdot 2^{8s+4j-2} + 7 \cdot 2^{7s+5j-3} - 2^{6s+6j-3}
 \end{aligned}$$

We have for $k \geq 3s$.

$$\begin{aligned}
 (10.97) \quad \Gamma_{3s}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times k} &= 16^{3(s-1)} \cdot \Gamma_3^{\left[\begin{smallmatrix} 1 \\ 1 \\ 1 \end{smallmatrix} \right] \times (k-3(s-1))} \\
 &= 2^{12s-12} \cdot (2^{3k-9s+9} - 7 \cdot 2^{2k-6s+6} + 7 \cdot 2^{k-3s+4} - 2^3) \\
 &= 2^{3k+3s-3} - 7 \cdot 2^{2k+6s-6} + 7 \cdot 2^{k+9s-8} - 2^{12s-9}
 \end{aligned}$$

Proof. Proof of (10.95) and (10.96)

The proof is by strong induction, that is:

If

- (H_0) is true, and
- for all $j \geq 1$, $(H_0) \wedge (H_1) \wedge \dots \wedge (H_j)$ implies (H_{j+1})

then (H_j) is true for all $j \geq 1$.

(H_0) is true
Obviously

$(H_0) \Rightarrow (H_1)$ Immediately by the proof of (10.76)

$(H_0) \wedge (H_1) \wedge \dots \wedge (H_j)$ implies (H_{j+1})

We are going to show that (H_{j+1}) is true assuming that (H_q) is true for all $0 \leq q \leq j$

$$\begin{aligned}
 (H_{j+1}) \quad \Gamma_{2s+1+(j+1)}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times k} &= 16^{3(j+1)} \cdot \Gamma_{2(s-(j+1))+1}^{\left[\begin{smallmatrix} s-(j+1) \\ s-(j+1) \\ s-(j+1) \end{smallmatrix} \right] \times (k-3(j+1))} \\
 &= 2^{12j+12} \cdot 105 \cdot (2^{2(k-3(j+1))+2(s-(j+1))-2} + 7 \cdot 2^{k-3(j+1)+5(s-(j+1))-3} - 31 \cdot 2^{k-3(j+1)+4(s-(j+1))-3}) \\
 &\quad + 2^{12j+12} \cdot 105 \cdot (2^{8(s-(j+1))-2} - 31 \cdot 2^{7(s-(j+1))-3} + 93 \cdot 2^{6(s-(j+1))-3}) \\
 &= 105 \cdot (2^{2k+2s+4j+2} + 7 \cdot 2^{k+5s+4j+1} - 31 \cdot 2^{k+4s+5j+2}) \\
 &\quad + 105 \cdot (2^{8s+4j+2} - 31 \cdot 2^{7s+5j+2} + 93 \cdot 2^{6s+6j+3})
 \end{aligned}$$

We get respectively from (10.1) and (10.2) for $k = 2s+j+3$

$$(10.98) \quad \sum_{i=0}^{2s+j+3} \Gamma_i^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3)} = 2^{9s+3j+6}$$

and

$$(10.99) \quad \sum_{i=0}^{2s+j+3} \Gamma_i^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3)} \cdot 2^{-i} = 2^{7s+2j+3} + 2^{6s+3j+6} - 2^{4s+2j+3}.$$

We have from (10.3), (10.4) with $k = 2s+j+3$

$$(10.100) \quad \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3)} = 7 \cdot 2^{4s-6} - 3 \cdot 2^{3s-5}$$

$$(10.101) \quad \sum_{i=0}^{s-1} \Gamma_i \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \cdot 2^{-i} = 15 \cdot 2^{3s-6} - 7 \cdot 2^{2s-5}$$

By (10.5) with $k=2s+j+3$

$$(10.102) \quad \Gamma_s \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) = 7 \cdot 2^{3s+j+2} - 7 \cdot 2^{2s} + 105 \cdot 2^{4s-6} - 21 \cdot 2^{3s-5}$$

$$(10.103) \quad \Gamma_s \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \cdot 2^{-s} = 7 \cdot 2^{2s+j+2} - 7 \cdot 2^s + 105 \cdot 2^{3s-6} - 21 \cdot 2^{2s-5}$$

From (10.62) and (10.63) with $k = 2k+j+3$ we obtain

$$(10.104) \quad \sum_{q=1}^{s-1} \Gamma_{s+q} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \\ = 49 \cdot 2^{7s+j-4} - 7 \cdot 2^{3s+j+2} - 21 \cdot 2^{6s+j-3} + 7 \cdot 2^{8s-6} - 105 \cdot 2^{7s-8} + 37 \cdot 2^{6s-7} - 7 \cdot 2^{4s-2} + 3 \cdot 2^{3s-2} + 7 \cdot 2^{2s}$$

$$(10.105) \quad \sum_{q=1}^{s-1} \Gamma_{s+q} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \cdot 2^{-(s+q)} \\ = 105 \cdot 2^{5s+j-4} - 49 \cdot 2^{4s+j-3} - 7 \cdot 2^{2s+j+2} + 15 \cdot 2^{6s-6} - 217 \cdot 2^{5s-8} + 77 \cdot 2^{4s-7} - 15 \cdot 2^{3s-3} + 7 \cdot 2^{2s-3} + 7 \cdot 2^s$$

We get from (10.48) with $k = 2s+j+3$

$$(10.106) \quad \Gamma_{2s} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) = 7 \cdot 2^{6s+2j+4} + 735 \cdot 2^{7s+j-4} - 819 \cdot 2^{6s+j-3} + 105 \cdot 2^{8s-6} - 3255 \cdot 2^{7s-8} + 2443 \cdot 2^{6s-7}$$

$$(10.107) \quad \Gamma_{2s} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \cdot 2^{-2s} = 7 \cdot 2^{4s+2j+4} + 735 \cdot 2^{5s+j-4} - 819 \cdot 2^{4s+j-3} + 105 \cdot 2^{6s-6} - 3255 \cdot 2^{5s-8} + 2443 \cdot 2^{4s-7}$$

From the induction hypothesis (H_q) for $0 \leq q \leq j$ we get by (10.95) with $k = 2s+j+3$.

$$(10.108) \quad \sum_{q=0}^j \Gamma_{2s+1+q} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \\ = \sum_{q=0}^j 105 \cdot (2^{2(2s+j+3)+2s+4q-2} + 7 \cdot 2^{(2s+j+3)+5s+4q-3} - 31 \cdot 2^{(2s+j+3)+4s+5q-3}) \\ + \sum_{q=0}^j 105 \cdot (2^{8s+4q-2} - 31 \cdot 2^{7s+5q-3} + 93 \cdot 2^{6s+6q-3}) \\ = 91 \cdot 2^{7s+5j+2} - 41 \cdot 2^{6s+6j+3} + 7 \cdot 2^{8s+4j+2} - 7 \cdot 2^{6s+2j+4} \\ - 49 \cdot 2^{7s+j} + 105 \cdot 2^{6s+j} - 7 \cdot 2^{8s-2} + 105 \cdot 2^{7s-3} - 155 \cdot 2^{6s-3}$$

$$(10.109) \quad \sum_{q=0}^j \Gamma_{2s+1+q} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \cdot 2^{-(2s+1+q)} \\ = \sum_{q=0}^j 105 \cdot (2^{2(2s+j+3)+2s+4q-2-(2s+1+q)} + 7 \cdot 2^{(2s+j+3)+5s+4q-3-(2s+1+q)} - 31 \cdot 2^{(2s+j+3)+4s+5q-3-(2s+1+q)})$$

$$\begin{aligned}
& + \sum_{q=0}^j 105 \cdot (2^{8s+4q-2-(2s+1+q)} - 31 \cdot 2^{7s+5q-3-(2s+1+q)} + 93 \cdot 2^{6s+6q-3-(2s+1+q)}) \\
& = \sum_{q=0}^j 105 \cdot (2^{4s+2j+3+3q} + 7 \cdot 2^{5s+j+3q-1} - 31 \cdot 2^{4s+j+4q-1}) + \sum_{q=0}^j 105 \cdot (2^{6s+3q-3} - 31 \cdot 2^{5s+4q-4} + 93 \cdot 2^{4s+5q-4}) \\
& = 203 \cdot 2^{5s+4j} - 73 \cdot 2^{4s+5j+1} - 15 \cdot 2^{4s+2j+3} + 15 \cdot 2^{6s+3j} - 15 \cdot 2^{6s-3} \\
& - 105 \cdot 2^{5s+j-1} + 217 \cdot 2^{4s+j-1} + 217 \cdot 2^{5s-4} - 315 \cdot 2^{4s-4}
\end{aligned}$$

We obtain from (10.98), (10.100), (10.102), (10.104), (10.106) and (10.108).

$$\begin{aligned}
& \sum_{i=0}^{2s+j+3} \Gamma_i \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) = \sum_{i=0}^{s-1} \Gamma_i \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) + \Gamma_s \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) + \sum_{q=1}^{s-1} \Gamma_{s+q} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \\
& + \Gamma_{2s} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) + \sum_{q=0}^j \Gamma_{2s+1+q} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) + \Gamma_{2s+j+2} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) + \Gamma_{2s+j+3} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \\
& = (7 \cdot 2^{4s-6} - 3 \cdot 2^{3s-5}) + (7 \cdot 2^{3s+j+2} - 7 \cdot 2^{2s} + 105 \cdot 2^{4s-6} - 21 \cdot 2^{3s-5}) \\
& + (287 \cdot 2^{7s-8} - 53 \cdot 2^{3s-2} - 131 \cdot 2^{6s-7} + 7 \cdot 2^{8s-6} - 7 \cdot 2^{4s-2} + 7 \cdot 2^{2s}) \\
& + (49 \cdot 2^{7s+j-4} - 7 \cdot 2^{3s+j+2} - 21 \cdot 2^{6s+j-3} + 7 \cdot 2^{8s-6} - 105 \cdot 2^{7s-8} + 37 \cdot 2^{6s-7} - 7 \cdot 2^{4s-2} + 3 \cdot 2^{3s-2} + 7 \cdot 2^{2s}) \\
& + (7 \cdot 2^{6s+2j+4} + 735 \cdot 2^{7s+j-4} - 819 \cdot 2^{6s+j-3} + 105 \cdot 2^{8s-6} - 3255 \cdot 2^{7s-8} + 2443 \cdot 2^{6s-7}) \\
& + (91 \cdot 2^{7s+5j+2} - 41 \cdot 2^{6s+6j+3} + 7 \cdot 2^{8s+4j+2} - 7 \cdot 2^{6s+2j+4} \\
& - 49 \cdot 2^{7s+j} + 105 \cdot 2^{6s+j} - 7 \cdot 2^{8s-2} + 105 \cdot 2^{7s-3} - 155 \cdot 2^{6s-3}) + \Gamma_{2s+j+2} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) + \Gamma_{2s+j+3} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) = 2^{9s+3j+6} \\
& \Downarrow
\end{aligned}$$

(10.110)

$$\Gamma_{2s+j+2} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) + \Gamma_{2s+j+3} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) = 2^{9s+3j+6} - 7 \cdot 2^{8s+4j+2} - 91 \cdot 2^{7s+5j+2} + 41 \cdot 2^{6s+6j+3}$$

We get in the same way from (10.99), (10.101), (10.103), (10.105), (10.107) and (10.109).

$$\begin{aligned}
& \sum_{i=0}^{2s+j+3} \Gamma_i \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \cdot 2^{-i} = \sum_{i=0}^{s-1} \Gamma_i \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \cdot 2^{-i} + \Gamma_s \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \cdot 2^{-s} + \sum_{q=1}^{s-1} \Gamma_{s+q} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \cdot 2^{-(s+q)} \\
& + \Gamma_{2s} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \cdot 2^{-2s} + \sum_{q=0}^j \Gamma_{2s+1+q} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \cdot 2^{-(2s+1+q)} + \Gamma_{2s+j+2} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \cdot 2^{-(2s+j+2)} + \Gamma_{2s+j+3} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \cdot 2^{-(2s+j+3)} \\
& = (15 \cdot 2^{3s-6} - 7 \cdot 2^{2s-5}) + (7 \cdot 2^{2s+j+2} - 7 \cdot 2^s + 105 \cdot 2^{3s-6} - 21 \cdot 2^{2s-5}) \\
& + (105 \cdot 2^{5s+j-4} - 49 \cdot 2^{4s+j-3} - 7 \cdot 2^{2s+j+2} + 15 \cdot 2^{6s-6} - 217 \cdot 2^{5s-8} + 77 \cdot 2^{4s-7} - 15 \cdot 2^{3s-3} + 7 \cdot 2^{2s-3} + 7 \cdot 2^s) \\
& + (7 \cdot 2^{4s+2j+4} + 735 \cdot 2^{5s+j-4} - 819 \cdot 2^{4s+j-3} + 105 \cdot 2^{6s-6} - 3255 \cdot 2^{5s-8} + 2443 \cdot 2^{4s-7}) \\
& + (203 \cdot 2^{5s+4j} - 73 \cdot 2^{4s+5j+1} - 15 \cdot 2^{4s+2j+3} + 15 \cdot 2^{6s+3j} - 15 \cdot 2^{6s-3} \\
& - 105 \cdot 2^{5s+j-1} + 217 \cdot 2^{4s+j-1} + 217 \cdot 2^{5s-4} - 315 \cdot 2^{4s-4}) \\
& + \Gamma_{2s+j+2} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \cdot 2^{-(2s+j+2)} + \Gamma_{2s+j+3} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \cdot 2^{-(2s+j+3)} = 2^{7s+2j+3} + 2^{6s+3j+6} - 2^{4s+2j+3} \\
& \Downarrow
\end{aligned}$$

$$\Gamma_{2s+j+2} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \cdot 2^{-(2s+j+2)} + \Gamma_{2s+j+3} \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times (2s+j+3) \cdot 2^{-(2s+j+3)} = 2^{7s+2j+3} + 73 \cdot 2^{4s+5j+1} - 203 \cdot 2^{5s+4j} + 49 \cdot 2^{6s+3j}$$

\Downarrow

(10.111)

$$2 \cdot \Gamma_{2s+j+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+j+3)} + \Gamma_{2s+j+3}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+j+3)} = 2^{9s+3j+6} + 73 \cdot 2^{6s+6j+4} - 203 \cdot 2^{7s+5j+3} + 49 \cdot 2^{8s+4j+3}$$

Hence by (10.110), (10.111)

$$(10.112) \quad \Gamma_{2s+j+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+j+3)} = 105 \cdot (2^{6s+6j+3} - 3 \cdot 2^{7s+5j+2} + 2^{8s+4j+2})$$

$$(10.113) \quad \Gamma_{2s+j+3}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+j+3)} = 2^{9s+3j+6} - 7 \cdot 2^{8s+4j+6} + 7 \cdot 2^{7s+5j+7} - 2^{6s+6j+9}$$

We have from (9.7) with $j \rightarrow j+2$, $j+2 \leq s-1$ and $k \geq 2s+j+3$

(10.114)

$$\Delta_k \Gamma_{2s+j+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = \Gamma_{2s+j+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (k+1)} - \Gamma_{2s+j+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times k} = 105 \cdot (3 \cdot 2^{2k+2s+4j+2} + 7 \cdot 2^{k+5s+4j+1} - 31 \cdot 2^{k+4s+5j+2})$$

We obtain from (10.114) for $k \geq 2s+j+3$ and (10.112)

$$\begin{aligned} \sum_{i=2s+j+3}^k (\Gamma_{2s+j+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (i+1)} - \Gamma_{2s+j+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times i}) &= \sum_{i=2s+j+3}^k 105 \cdot (3 \cdot 2^{2i+2s+4j+2} + 7 \cdot 2^{i+5s+4j+1} - 31 \cdot 2^{i+4s+5j+2}) \\ \Downarrow \\ \Gamma_{2s+j+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (k+1)} - \Gamma_{2s+j+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (2s+j+3)} &= 315 \cdot 2^{2s+4j+2} \cdot \sum_{i=2s+j+3}^k 2^{2i} + (735 \cdot 2^{5s+4j+1} - 3255 \cdot 2^{4s+5j+2}) \cdot \sum_{i=2s+j+3}^k 2^i \\ &= 105 \cdot 2^{2k+2s+4j+4} + 735 \cdot 2^{k+5s+4j+2} - 3255 \cdot 2^{k+4s+5j+3} - 735 \cdot 2^{7s+5j+4} + 2415 \cdot 2^{6s+6j+5} \\ \Downarrow \end{aligned}$$

$$\begin{aligned} \Gamma_{2s+j+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times k} &= 105 \cdot 2^{2k+2s+4j+2} + 735 \cdot 2^{k+5s+4j+1} - 3255 \cdot 2^{k+4s+5j+2} - 735 \cdot 2^{7s+5j+4} + 2415 \cdot 2^{6s+6j+5} \\ &+ 105 \cdot (2^{6s+6j+3} - 3 \cdot 2^{7s+5j+2} + 2^{8s+4j+2}) \quad \text{for } k \geq 2s+j+4 \end{aligned}$$

 $\Downarrow$

(10.115)

$$\begin{aligned} \Gamma_{2s+j+2}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times k} &= 105 \cdot (2^{2k+2s+4j+2} + 7 \cdot 2^{k+5s+4j+1} - 31 \cdot 2^{k+4s+5j+2}) \\ &+ 105 \cdot (2^{8s+4j+2} - 31 \cdot 2^{7s+5j+2} + 93 \cdot 2^{6s+6j+3}) \quad \text{for } k \geq 2s+j+4 \end{aligned}$$

From (10.112) we see that (10.115) is valid for $k = 2s+j+3$.Proof of (10.97)We then obtain from (10.110) with $j = s-2$

(10.116)

$$\Gamma_{3s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (3s+1)} = 2^{12s} - 7 \cdot 2^{8s+4(s-2)+2} - 91 \cdot 2^{7s+5(s-2)+2} + 41 \cdot 2^{6s+6(s-2)+3} = 315 \cdot 2^{12s-9}$$

We have from (9.7) with $j = s$ and $k \geq 3s+1$

$$(10.117) \quad \Delta_k \Gamma_{3s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix}} = \Gamma_{3s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (k+1)} - \Gamma_{3s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times k} = 7 \cdot 2^{3k+3s-3} - 21 \cdot 2^{2k+6s-6} + 7 \cdot 2^{k+9s-8}$$

We obtain from (10.117) for $k \geq 3s+1$ and (10.116)

$$\begin{aligned} \sum_{i=3s+1}^k (\Gamma_{3s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times (i+1)} - \Gamma_{3s}^{\begin{bmatrix} s \\ s \\ s \end{bmatrix} \times i}) &= \sum_{i=3s+1}^k 7 \cdot 2^{3i+3s-3} - 21 \cdot 2^{2i+6s-6} + 7 \cdot 2^{i+9s-8} \\ \Downarrow \end{aligned}$$

$$\begin{aligned} \Gamma_{3s}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (k+1)} - \Gamma_{3s}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (3s+1)} &= 7 \cdot 2^{3s-3} \cdot \sum_{i=3s+1}^k 2^{3i} - 21 \cdot 2^{6s-6} \cdot \sum_{i=3s+1}^k 2^{2i} + 7 \cdot 2^{9s-8} \cdot \sum_{i=3s+1}^k 2^i \\ &= 2^{3k+3s} - 7 \cdot 2^{2k+6s-4} + 7 \cdot 2^{k+9s-7} - 79 \cdot 2^{12s-7} \end{aligned}$$

$\Updownarrow$

$$\Gamma_{3s}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} = 2^{3k+3s-3} - 7 \cdot 2^{2k+6s-6} + 7 \cdot 2^{k+9s-8} - 79 \cdot 2^{12s-7} + 315 \cdot 2^{12s-9} \quad \text{for } k \geq 3s+2$$

$\Updownarrow$

(10.118)

$$\Gamma_{3s}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} = 2^{3k+3s-3} - 7 \cdot 2^{2k+6s-6} + 7 \cdot 2^{k+9s-8} - 2^{12s-9}$$

From (10.116) we see that (10.118) is valid for $k = 3s+1$. $\square$

Theorem 10.9. *We have*

(10.119)

$$\Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} = \begin{cases} 1 & \text{if } i = 0, \\ 105 \cdot 2^{4i-6} - 21 \cdot 2^{3i-5} & \text{if } 1 \leq i \leq s-1, k \geq i+1, \\ 7 \cdot 2^{k+s-1} - 7 \cdot 2^{2s} + 105 \cdot 2^{4s-6} - 21 \cdot 2^{3s-5} & \text{if } i = s, k \geq s+1, s \geq 1, \\ 147 \cdot (5 \cdot 2^{j-1} - 1) \cdot 2^{k+s+3j-6} & \\ + 21 \cdot [5 \cdot 2^{4s+4j-6} - 2^{3s+3j-5} - (155 \cdot 2^{j-1} - 35) \cdot 2^{2s+4j-7}] & \text{if } i = s+j, 1 \leq j \leq s-1, k \geq s+j+1, \\ 7 \cdot 2^{2k+2s-2} + 21 \cdot [35 \cdot 2^{k+5s-7} - 39 \cdot 2^{k+4s-6}] & \\ + 7 \cdot [15 \cdot 2^{8s-6} - 465 \cdot 2^{7s-8} + 349 \cdot 2^{6s-7}] & \text{if } i = 2s, k \geq 2s+1, \\ 105 \cdot (2^{2k+2s+4j-2} + 7 \cdot 2^{k+5s+4j-3} - 31 \cdot 2^{k+4s+5j-3}) & \\ + 105 \cdot (2^{8s+4j-2} - 31 \cdot 2^{7s+5j-3} + 93 \cdot 2^{6s+6j-3}) & \text{if } i = 2s+1+j, k \geq 2s+2+j, \\ & 0 \leq j \leq s-2, \\ 2^{3k+3s-3} - 7 \cdot 2^{2k+6s-6} + 7 \cdot 2^{k+9s-8} - 2^{12s-9} & \text{if } i = 3s, k \geq 3s \end{cases}$$

(10.120)

$$\Gamma_i^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times i} = \begin{cases} 2^{3s+3i-3} - 7 \cdot 2^{4i-6} + 3 \cdot 2^{3i-5} & \text{if } 1 \leq i \leq s+1, \\ 2^{6s+3j-3} + 7 \cdot 2^{2s+5j-8} - 7 \cdot 2^{2s+4j-7} - 7 \cdot 2^{4s+4j-6} + 3 \cdot 2^{3s+3j-5} & \text{if } i = s+j, 1 \leq j \leq s+1, \\ 2^{9s+3j} - 7 \cdot 2^{8s+4j-2} + 7 \cdot 2^{7s+5j-3} - 2^{6s+6j-3} & \text{if } i = 2s+1+j, 0 \leq j \leq s-1, \end{cases}$$

We have for $0 \leq j \leq s-2, k \geq 2s+2+j$

(10.121)

$$\begin{aligned} \Gamma_{2s+1+j}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times k} &= 16^{3j} \cdot \Gamma_{2(s-j)+1}^{\left[\begin{smallmatrix} s-j & s-j \\ s-j & s-j \end{smallmatrix} \right] \times (k-3j)} \\ &= 2^{12j} \cdot 105 \cdot (2^{2(k-3j)+2(s-j)-2} + 7 \cdot 2^{k-3j+5(s-j)-3} - 31 \cdot 2^{k-3j+4(s-j)-3}) \\ &\quad + 2^{12j} \cdot 105 \cdot (2^{8(s-j)-2} - 31 \cdot 2^{7(s-j)-3} + 93 \cdot 2^{6(s-j)-3}) \\ &= 105 \cdot (2^{2k+2s+4j-2} + 7 \cdot 2^{k+5s+4j-3} - 31 \cdot 2^{k+4s+5j-3}) \\ &\quad + 105 \cdot (2^{8s+4j-2} - 31 \cdot 2^{7s+5j-3} + 93 \cdot 2^{6s+6j-3}) \end{aligned}$$

We have for $0 \leq j \leq s-1$.

(10.122)

$$\begin{aligned} \Gamma_{2s+1+j}^{\left[\begin{smallmatrix} s & s \\ s & s \end{smallmatrix} \right] \times (2s+1+j)} &= 16^{3j} \cdot \Gamma_{2(s-j)+1}^{\left[\begin{smallmatrix} s-j & s-j \\ s-j & s-j \end{smallmatrix} \right] \times (2(s-j)+1)} \\ &= 2^{12j} \cdot (2^{9(s-j)} - 7 \cdot 2^{8(s-j)-2} + 7 \cdot 2^{7(s-j)-3} - 2^{6(s-j)-3}) \\ &= 2^{9s+3j} - 7 \cdot 2^{8s+4j-2} + 7 \cdot 2^{7s+5j-3} - 2^{6s+6j-3} \end{aligned}$$

We have for $k \geq 3s$.

$$\begin{aligned}
 (10.123) \quad \Gamma_{3s}^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times k} &= 16^{3(s-1)} \cdot \Gamma_3^{\left[\begin{smallmatrix} 1 \\ 1 \\ 1 \end{smallmatrix} \right] \times (k-3(s-1))} \\
 &= 2^{12s-12} \cdot (2^{3k-9s+9} - 7 \cdot 2^{2k-6s+6} + 7 \cdot 2^{k-3s+4} - 2^3) \\
 &= 2^{3k+3s-3} - 7 \cdot 2^{2k+6s-6} + 7 \cdot 2^{k+9s-8} - 2^{12s-9}
 \end{aligned}$$

Theorem 10.10. Let q be a rational integer ≥ 1 , then

$$(10.124) \quad g_{k,s}(t, \eta, \xi) = g(t, \eta, \xi) = \sum_{\deg Y \leq k-1} \sum_{\deg Z \leq s-1} E(tYZ) \sum_{\deg U \leq s-1} E(\eta YU) \sum_{\deg V \leq s-1} E(\eta YV) = 2^{3s+k-r(D \left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times k)}(t, \eta, \xi),$$

$$(10.125) \quad \int_{\mathbb{P} \times \mathbb{P} \times \mathbb{P}} g^q(t, \eta, \xi) dt d\eta d\xi = 2^{(3s+k)q} \cdot 2^{-3k-3s+3} \cdot \sum_{i=0}^{\inf(3s,k)} \Gamma_i^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times k} \cdot 2^{-qi}.$$

Theorem 10.11. We denote by $R_q(k, s)$ the number of solutions $(Y_1, Z_1, U_1, V_1, \dots, Y_q, Z_q, U_q, V_q)$ of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0, \\ Y_1 U_1 + Y_2 U_2 + \dots + Y_q U_q = 0, \\ Y_1 V_1 + Y_2 V_2 + \dots + Y_q V_q = 0, \end{cases}$$

satisfying the degree conditions

$$\deg Y_i \leq k-1, \quad \deg Z_i \leq s-1, \quad \deg U_i \leq s-1, \quad \deg V_i \leq s-1 \quad \text{for } 1 \leq i \leq q.$$

Then

$$(10.126) \quad R_q(k, s) = \int_{\mathbb{P} \times \mathbb{P} \times \mathbb{P}} g_{k,s}^q(t, \eta, \xi) dt d\eta d\xi = 2^{(3s+k)q} \cdot 2^{-3k-3s+3} \cdot \sum_{i=0}^{\inf(3s,k)} \Gamma_i^{\left[\begin{smallmatrix} s \\ s \\ s \end{smallmatrix} \right] \times k} \cdot 2^{-qi}.$$

Example. $s = 1$, $k \geq i + 1$ for $0 \leq i \leq 2$

$$\Gamma_i^{\left[\begin{smallmatrix} 1 \\ 1 \\ 1 \end{smallmatrix} \right] \times k} = \begin{cases} 1 & \text{if } i = 0 \\ 7 \cdot (2^k - 1) & \text{if } i = 1 \\ 7 \cdot (2^k - 1) \cdot (2^k - 2) & \text{if } i = 2 \\ 2^{3k} - 7 \cdot 2^{2k} + 7 \cdot 2^{k+1} - 2^3 & \text{if } i = 3, k \geq 3 \end{cases}$$

Example. $s = 2$, $k \geq i + 1$ for $0 \leq i \leq 5$

$$\Gamma_i^{\left[\begin{smallmatrix} 2 \\ 2 \\ 2 \end{smallmatrix} \right] \times k} = \begin{cases} 1 & \text{if } i = 0 \\ 21 & \text{if } i = 1 \\ 7 \cdot 2^{k+1} + 266 & \text{if } i = 2 \\ 147 \cdot 2^{k+1} + 1344 & \text{if } i = 3 \\ 7 \cdot 2^{2k+2} + 651 \cdot 2^{k+2} - 22624 & \text{if } i = 4 \\ 105 \cdot 2^{2k+2} - 315 \cdot 2^{k+5} + 53760 & \text{if } i = 5 \\ 2^{3k+3} - 7 \cdot 2^{2k+6} + 7 \cdot 2^{k+10} - 32768 & \text{if } i = 6, k \geq 6 \end{cases}$$

Example. $s = 2, k = 6$.

The number $\Gamma_i^{\left[\frac{2}{2}\right] \times 6}$ of rank i matrices of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 \\ \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 & \alpha_7 \\ \beta_1 & \beta_2 & \beta_3 & \beta_4 & \beta_5 & \beta_6 \\ \beta_2 & \beta_3 & \beta_4 & \beta_5 & \beta_6 & \beta_7 \\ \gamma_1 & \gamma_2 & \gamma_3 & \gamma_4 & \gamma_5 & \gamma_6 \\ \gamma_2 & \gamma_3 & \gamma_4 & \gamma_5 & \gamma_6 & \gamma_7 \end{pmatrix}$$

is equal to

$$\begin{cases} 1 & \text{if } i = 0 \\ 21 & \text{if } i = 1 \\ 1162 & \text{if } i = 2 \\ 20160 & \text{if } i = 3 \\ 258720 & \text{if } i = 4 \\ 1128960 & \text{if } i = 5 \\ 688128 & \text{if } i = 6 \end{cases}$$

The number of solutions

$(Y_1, Z_1, U_1, V_1, \dots, Y_q, Z_q, U_q, V_q)$ of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0, \\ Y_1 U_1 + Y_2 U_2 + \dots + Y_q U_q = 0, \\ Y_1 V_1 + Y_2 V_2 + \dots + Y_q V_q = 0, \end{cases}$$

satisfying the degree conditions

$$\deg Y_i \leq 5, \quad \deg Z_i \leq 1, \quad \deg U_i \leq 1, \quad \deg V_i \leq 1 \quad \text{for } 1 \leq i \leq q.$$

is equal to

$$\begin{aligned} R_q(6, 2) &= \int_{\mathbb{P} \times \mathbb{P} \times \mathbb{P}} g_{6,2}^q(t, \eta, \xi) dt d\eta d\xi = 2^{12q-21} \cdot \sum_{i=0}^6 \Gamma_i^{\left[\frac{2}{2}\right] \times 6} \cdot 2^{-qi} \\ &= 2^{12q-21} \cdot (1 + 21 \cdot 2^{-q} + 1162 \cdot 2^{-2q} + 20160 \cdot 2^{-3q} + 258720 \cdot 2^{-4q} + 1128960 \cdot 2^{-5q} + 688128 \cdot 2^{-6q}) \\ &= 2^{6q-21} \cdot (2^{6q} + 21 \cdot 2^{5q} + 1162 \cdot 2^{4q} + 20160 \cdot 2^{3q} + 258720 \cdot 2^{2q} + 1128960 \cdot 2^q + 688128) \end{aligned}$$

Example. $s = 3, k \geq i + 1$ for $0 \leq i \leq 8$

$$\Gamma_i^{\left[\frac{3}{3}\right] \times k} = \begin{cases} 1 & \text{if } i = 0 \\ 21 & \text{if } i = 1 \\ 378 & \text{if } i = 2 \\ 7 \cdot 2^{k+2} + 5936 & \text{if } i = 3 \\ 147 \cdot 2^{k+2} + 84672 & \text{if } i = 4 \\ 147 \cdot 9 \cdot 2^{k+3} + 959616 & \text{if } i = 5 \\ 7 \cdot 2^{2k+4} + 2121 \cdot 2^{k+6} + 5863424 & \text{if } i = 6 \\ 105 \cdot 2^{2k+4} + 2625 \cdot 2^{k+9} - 92897280 & \text{if } i = 7 \\ 105 \cdot 2^{2k+8} - 315 \cdot 2^{k+14} + 220200960 & \text{if } i = 8 \\ 2^{3k+6} - 7 \cdot 2^{2k+12} + 7 \cdot 2^{k+19} - 134217728 & \text{if } i = 9, k \geq 9 \end{cases}$$

Example. $s = 3, k = 5, q = 3$.

The number $\Gamma_i^{\left[\begin{smallmatrix} 3 \\ 3 \\ 3 \end{smallmatrix}\right] \times 5}$ of rank i matrices of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 \\ \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 \\ \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 & \alpha_7 \\ \beta_1 & \beta_2 & \beta_3 & \beta_4 & \beta_5 \\ \beta_2 & \beta_3 & \beta_4 & \beta_5 & \beta_6 \\ \beta_3 & \beta_4 & \beta_5 & \beta_6 & \beta_7 \\ \gamma_1 & \gamma_2 & \gamma_3 & \gamma_4 & \gamma_5 \\ \gamma_2 & \gamma_3 & \gamma_4 & \gamma_5 & \gamma_6 \\ \gamma_3 & \gamma_4 & \gamma_5 & \gamma_6 & \gamma_7 \end{pmatrix}$$

is equal to

$$\begin{cases} 1 & \text{if } i = 0 \\ 21 & \text{if } i = 1 \\ 378 & \text{if } i = 2 \\ 6832 & \text{if } i = 3 \\ 103488 & \text{if } i = 4 \\ 1986432 & \text{if } i = 5 \end{cases}$$

The number of solutions $(Y_1, Z_1, U_1, V_1, Y_2, Z_2, U_2, V_2, Y_3, Z_3, U_3, V_3)$ of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + Y_3 Z_3 = 0, \\ Y_1 U_1 + Y_2 U_2 + Y_3 U_3 = 0, \\ Y_1 V_1 + Y_2 V_2 + Y_3 V_3 = 0, \end{cases}$$

satisfying the degree conditions

$$\deg Y_i \leq 4, \quad \deg Z_i \leq 2, \quad \deg U_i \leq 2, \quad \deg V_i \leq 2 \quad \text{for } 1 \leq i \leq 3.$$

is equal to

$$\begin{aligned} R_3(5, 3) &= \int_{\mathbb{P} \times \mathbb{P} \times \mathbb{P}} g_{5,3}^3(t, \eta, \xi) dt d\eta d\xi = 2^{33} \cdot \sum_{i=0}^5 \Gamma_i^{\left[\begin{smallmatrix} 3 \\ 3 \\ 3 \end{smallmatrix}\right] \times 5} \cdot 2^{-3i} \\ &= 2^{33} \cdot (1 + 21 \cdot 2^{-3} + 378 \cdot 2^{-6} + 6832 \cdot 2^{-9} + 103488 \cdot 2^{-12} + 1986432 \cdot 2^{-15}) = 3563904 \times 2^{18} \end{aligned}$$

11. Computation of $\Delta_k \Gamma_i \left[\begin{smallmatrix} s \\ s+m \\ s+m \end{smallmatrix} \right]$ for $0 \leq i \leq 3s+2m$

11.1. Notation.

Definition 11.1. In this section we define γ_j to be equal to $(21 \cdot 2^{3j-4} - 3 \cdot 2^{2j-3})$ for $1 \leq j \leq m-1$.

11.2. Introduction. We adapt the method in Section 9 to compute explicitly $\Delta_k \Gamma_i \left[\begin{smallmatrix} s \\ s+m \\ s+m \end{smallmatrix} \right]$ for $s \leq i \leq 3s+2m$

11.3. Computation of $\Delta_k \Gamma_{s+j} \left[\begin{smallmatrix} s \\ s+m+l \\ s+m+l \end{smallmatrix} \right]$ for $0 \leq j \leq m-1, l \geq 0$.

Lemma 11.2. We have the following reduction formula for $0 \leq j \leq m, l \geq 0$

$$(11.1) \quad \Delta_k \Gamma_{s+j} \left[\begin{smallmatrix} s \\ s+m+l \\ s+m+l \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_{s+j-1} \left[\begin{smallmatrix} s \\ s+m+l-1 \\ s+m+l-1 \end{smallmatrix} \right] = 4^j \cdot \Delta_k \Gamma_s \left[\begin{smallmatrix} s+(m-j) \\ s+m+l \end{smallmatrix} \right] + \sum_{q=0}^{j-1} 4^q \cdot \Delta_k \omega_{s+j-q}(s, m-q, l+q)$$

From (11.1) we deduce successively for $j = 0, 1, \dots, m-1, m$

$$(11.2) \quad \Delta_k \Gamma_{s+j} \left[\begin{smallmatrix} s \\ s+m+l \\ s+m+l \end{smallmatrix} \right] = \begin{cases} 2^{k+s-1} & \text{if } j = 0 \\ \gamma_j \cdot 2^{k+s-1} = (21 \cdot 2^{3j-4} - 3 \cdot 2^{2j-3}) \cdot 2^{k+s-1} & \text{if } 1 \leq j \leq m-1, l \geq 0 \\ 21 \cdot 2^{k+s+3m-5} + 13 \cdot 2^{k+s+2m-4} & \text{if } j = m \geq 1, l \geq 1 \\ 21 \cdot 2^{k+s+3m-5} + 45 \cdot 2^{k+s+2m-4} & \text{if } j = m \geq 1, l = 0 \end{cases}$$

Proof. From (8.3) we obtain successively the following equations

$$\begin{aligned} \Delta_k \Gamma_{s+j} \left[\begin{smallmatrix} s \\ s+m+l \\ s+m+l \end{smallmatrix} \right] - 4 \cdot \Delta_k \Gamma_{s+j-1} \left[\begin{smallmatrix} s \\ s+m+l \\ s+m+l \end{smallmatrix} \right] &= 8 \cdot \left[\Delta_k \Gamma_{s+j-1} \left[\begin{smallmatrix} s \\ s+m+l-1 \\ s+m+l-1 \end{smallmatrix} \right] - 4 \cdot \Delta_k \Gamma_{s+j-2} \left[\begin{smallmatrix} s \\ s+m+l-1 \\ s+m+l-1 \end{smallmatrix} \right] \right] + \Delta_k \omega_{s+j}(s, m, l) \\ 4 \cdot \left[\Delta_k \Gamma_{s+j-1} \left[\begin{smallmatrix} s \\ s+m+l \\ s+m+l \end{smallmatrix} \right] - 4 \cdot \Delta_k \Gamma_{s+j-2} \left[\begin{smallmatrix} s \\ s+m+l \\ s+m+l \end{smallmatrix} \right] \right] &= 4 \cdot \left[8 \cdot \left[\Delta_k \Gamma_{s+j-2} \left[\begin{smallmatrix} s \\ s+m+l-1 \\ s+m+l-1 \end{smallmatrix} \right] - 4 \cdot \Delta_k \Gamma_{s+j-3} \left[\begin{smallmatrix} s \\ s+m+l-1 \\ s+m+l-1 \end{smallmatrix} \right] \right] + \Delta_k \omega_{s+j-1}(s, m-1, l+1) \right] \\ 4^2 \cdot \left[\Delta_k \Gamma_{s+j-2} \left[\begin{smallmatrix} s \\ s+m+l \\ s+m+l \end{smallmatrix} \right] - 4 \cdot \Delta_k \Gamma_{s+j-3} \left[\begin{smallmatrix} s \\ s+m+l \\ s+m+l \end{smallmatrix} \right] \right] &= 4^2 \cdot \left[8 \cdot \left[\Delta_k \Gamma_{s+j-3} \left[\begin{smallmatrix} s \\ s+m+l-1 \\ s+m+l-1 \end{smallmatrix} \right] - 4 \cdot \Delta_k \Gamma_{s+j-4} \left[\begin{smallmatrix} s \\ s+m+l-1 \\ s+m+l-1 \end{smallmatrix} \right] \right] + \Delta_k \omega_{s+j-2}(s, m-2, l+2) \right] \\ &\vdots \\ 4^{j-1} \cdot \left[\Delta_k \Gamma_{s+j-(j-1)} \left[\begin{smallmatrix} s \\ s+m+l \\ s+m+l \end{smallmatrix} \right] - 4 \cdot \Delta_k \Gamma_{s+j-j} \left[\begin{smallmatrix} s \\ s+m+l \\ s+m+l \end{smallmatrix} \right] \right] &= 4^{j-1} \cdot \left[8 \cdot \left[\Delta_k \Gamma_{s+j-j} \left[\begin{smallmatrix} s \\ s+m+l-1 \\ s+m+l-1 \end{smallmatrix} \right] - 4 \cdot \Delta_k \Gamma_{s+j-(j+1)} \left[\begin{smallmatrix} s \\ s+m+l-1 \\ s+m+l-1 \end{smallmatrix} \right] \right] + \Delta_k \omega_{s+j-(j-1)}(s, m-(j-1), l+(j-1)) \right] \end{aligned}$$

By summing the left-hand side and the right-hand side of the above equations we obtain the following equality

$$\begin{aligned} \Delta_k \Gamma_{s+j} \left[\begin{smallmatrix} s \\ s+m+l \\ s+m+l \end{smallmatrix} \right] - 4^j \cdot \Delta_k \Gamma_s \left[\begin{smallmatrix} s+(m-j) \\ s+m+l \end{smallmatrix} \right] &= 8 \cdot \left[\Delta_k \Gamma_{s+j-1} \left[\begin{smallmatrix} s \\ s+m+l-1 \\ s+m+l-1 \end{smallmatrix} \right] - 4^j \cdot \Delta_k \Gamma_{s-1} \left[\begin{smallmatrix} s+(m-j) \\ s+m+l-1 \end{smallmatrix} \right] \right] + \sum_{q=0}^{j-1} 4^q \cdot \Delta_k \omega_{s+j-q}(s, m-q, l+q) \end{aligned}$$

$\Downarrow$

$$\Delta_k \Gamma_{s+j} \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_{s+j-1} \left[\begin{smallmatrix} s \\ s+m+l-1 \end{smallmatrix} \right] = 4^j \cdot \left[\Delta_k \Gamma_s \left[\begin{smallmatrix} s+(m-j) \\ s+m+l \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_{s-1} \left[\begin{smallmatrix} s+(m-j) \\ s+m+l-1 \end{smallmatrix} \right] \right] + \sum_{q=0}^{j-1} 4^q \cdot \Delta_k \omega_{s+j-q}(s, m-q, l+q)$$

$\Downarrow$

$$\Delta_k \Gamma_{s+j} \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_{s+j-1} \left[\begin{smallmatrix} s \\ s+m+l-1 \end{smallmatrix} \right] = 4^j \cdot \Delta_k \Gamma_s \left[\begin{smallmatrix} s+(m-j) \\ s+m+l \end{smallmatrix} \right] + \sum_{q=0}^{j-1} 4^q \cdot \Delta_k \omega_{s+j-q}(s, m-q, l+q)$$

To prove (11.2) we shall apply Lemma 7.4.

The case $j = 1$

$$\begin{aligned} \Delta_k \Gamma_{s+1} \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] &= 8 \cdot \Delta_k \Gamma_s \left[\begin{smallmatrix} s \\ s+m+l-1 \end{smallmatrix} \right] + 4 \cdot \Delta_k \Gamma_s \left[\begin{smallmatrix} s+(m-1) \\ s+m+l \end{smallmatrix} \right] + \Delta_k \omega_{s+1}(s, m, l) \\ &= 8 \cdot 2^{k+s-1} + 4 \cdot 2^{k+s-1} - 3 \cdot 2^{k+s-1} = 9 \cdot 2^{k+s-1} = \gamma_1 \cdot 2^{k+s-1} \quad \text{if } m \geq 2, \quad l \geq 0 \end{aligned}$$

The case $j = 2$

$$\begin{aligned} \Delta_k \Gamma_{s+2} \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] &= 8 \cdot \Delta_k \Gamma_{s+1} \left[\begin{smallmatrix} s \\ s+m+l-1 \end{smallmatrix} \right] + 4^2 \cdot \Delta_k \Gamma_s \left[\begin{smallmatrix} s+(m-2) \\ s+m+l \end{smallmatrix} \right] + \Delta_k \omega_{s+2}(s, m, l) + 4 \cdot \Delta_k \omega_{s+1}(s, m-1, l+1) \\ &= 8 \cdot 9 \cdot 2^{k+s-1} + 4^2 \cdot 2^{k+s-1} + 2^{k+s} - 4 \cdot 3 \cdot 2^{k+s-1} = 78 \cdot 2^{k+s-1} = \gamma_2 \cdot 2^{k+s-1} \quad \text{if } m \geq 3, \quad l \geq 0 \end{aligned}$$

The case $j = 3$

$$\begin{aligned} \Delta_k \Gamma_{s+3} \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] &= 8 \cdot \Delta_k \Gamma_{s+2} \left[\begin{smallmatrix} s \\ s+m+l-1 \end{smallmatrix} \right] + 4^3 \cdot \Delta_k \Gamma_s \left[\begin{smallmatrix} s+(m-3) \\ s+m+l \end{smallmatrix} \right] \\ &+ \Delta_k \omega_{s+3}(s, m, l) + 4 \cdot \Delta_k \omega_{s+2}(s, m-1, l+1) + 4^2 \cdot \Delta_k \omega_{s+1}(s, m-2, l+2) \\ &= 8 \cdot 78 \cdot 2^{k+s-1} + 4^3 \cdot 2^{k+s-1} + 4 \cdot 2^{k+s} - 4^2 \cdot 3 \cdot 2^{k+s-1} = 648 \cdot 2^{k+s-1} = \gamma_3 \cdot 2^{k+s-1} \quad \text{if } m \geq 4, \quad l \geq 0 \end{aligned}$$

The case general $1 \leq j \leq m-1$

Proof by induction on $j \leq m-1$. Assume that $\Delta_k \Gamma_{s+j-1} \left[\begin{smallmatrix} s \\ s+m+l-1 \end{smallmatrix} \right] = \gamma_{j-1} \cdot 2^{k+s-1}$ for $1 \leq j-1 \leq m-2$.

We shall prove that $\Delta_k \Gamma_{s+j} \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] = \gamma_j \cdot 2^{k+s-1}$ for $1 \leq j \leq m-1$.

$$\begin{aligned} \Delta_k \Gamma_{s+j} \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] &= 8 \cdot \Delta_k \Gamma_{s+j-1} \left[\begin{smallmatrix} s \\ s+m+l-1 \end{smallmatrix} \right] + 4^j \cdot \Delta_k \Gamma_s \left[\begin{smallmatrix} s+(m-j) \\ s+m+l \end{smallmatrix} \right] + \sum_{q=0}^{j-1} 4^q \cdot \Delta_k \omega_{s+j-q}(s, m-q, l+q) \\ &= 8 \cdot \gamma_{j-1} \cdot 2^{k+s-1} + 2^{2j} \cdot 2^{k+s-1} + \sum_{q=j-2}^{j-1} 4^q \cdot \Delta_k \omega_{s+j-q}(s, m-q, l+q) \\ &= 8 \cdot (21 \cdot 2^{3(j-1)} - 3 \cdot 2^{2(j-1)-3}) \cdot 2^{k+s-1} + 2^{2j} \cdot 2^{k+s-1} \\ &+ 4^{j-2} \cdot \Delta_k \omega_{s+j-(j-2)}(s, m-(j-2), l+(j-2)) + 4^{j-1} \cdot \Delta_k \omega_{s+j-(j-1)}(s, m-(j-1), l+(j-1)) \\ &= 8 \cdot [21 \cdot 2^{3(j-1)} - 3 \cdot 2^{2(j-1)-3}] \cdot 2^{k+s-1} + 2^{2j} \cdot 2^{k+s-1} + 2^{2j-4} \cdot 2^{k+s} + 2^{2j-2} \cdot (-3 \cdot 2^{k+s-1}) \\ &= [21 \cdot 2^{3j-4} - 3 \cdot 2^{2j-3}] \cdot 2^{k+s-1} = \gamma_j \cdot 2^{k+s-1} \quad \text{if } m \geq j+1, \quad l \geq 0 \end{aligned}$$

The case $j = m$, $l \geq 1$

$$\begin{aligned}
\Delta_k \Gamma_{s+m}^{\left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{s+m-1}^{\left[\begin{smallmatrix} s \\ s+m+l-1 \end{smallmatrix} \right]} &= 4^m \cdot \Delta_k \Gamma_s^{\left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right]} + \sum_{q=0}^{m-1} 4^q \cdot \Delta_k \omega_{s+m-q}(s, m-q, l+q) \\
\Delta_k \Gamma_{s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right]} &= 2^{2m} \cdot 3 \cdot 2^{k+s-1} + 8 \cdot \gamma_{m-1} \cdot 2^{k+s-1} + \sum_{q=0}^{m-1} 4^q \cdot \Delta_k \omega_{s+m-q}(s, m-q, l+q) \\
&= 2^{2m} \cdot 3 \cdot 2^{k+s-1} + 8 \cdot \gamma_{m-1} \cdot 2^{k+s-1} + \sum_{q=m-2}^{m-1} 4^q \cdot \Delta_k \omega_{s+m-q}(s, m-q, l+q) \\
&= 3 \cdot 2^{k+s+2m-1} + 8 \cdot [21 \cdot 2^{3m-7} - 3 \cdot 2^{2m-5}] \cdot 2^{k+s-1} \\
&\quad + 2^{2m-4} \cdot \Delta_k \omega_{s+2}(s, 2, l+(m-2)) + 2^{2m-2} \cdot \Delta_k \omega_{s+1}(s, 1, l+(m-1)) \\
&= 3 \cdot 2^{k+s+2m-1} + [21 \cdot 2^{3m-7} - 3 \cdot 2^{2m-5}] \cdot 2^{k+s+2} + 2^{2m-4} \cdot 2^{k+s} + 2^{2m-2} \cdot (-3 \cdot 2^{k+s-1}) \\
&= 21 \cdot 2^{k+s+3m-5} + 13 \cdot 2^{k+s+2m-4}
\end{aligned}$$

The case $j = m$, $l = 0$

$$\begin{aligned}
\Delta_k \Gamma_{s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right]} &= 2^{2m} \cdot 3 \cdot 2^{k+s-1} + 8 \cdot \Delta_k \Gamma_{s+m-1}^{\left[\begin{smallmatrix} s \\ s+m-1 \end{smallmatrix} \right]} + \sum_{q=0}^{m-1} 4^q \cdot \Delta_k \omega_{s+m-q}(s, m-q, q) \\
&= 2^{2m} \cdot 3 \cdot 2^{k+s-1} + 8 \cdot [21 \cdot 2^{k+s+3m-8} + 13 \cdot 2^{k+s+2m-6}] + \sum_{q=m-2}^{m-1} 4^q \cdot \Delta_k \omega_{s+m-q}(s, m-q, q) \\
&= 2^{2m} \cdot 3 \cdot 2^{k+s-1} + 8 \cdot [21 \cdot 2^{k+s+3m-8} + 13 \cdot 2^{k+s+2m-6}] + 2^{2m-4} \cdot \Delta_k \omega_{s+2}(s, 2, m-2) + 2^{2m-2} \cdot \Delta_k \omega_{s+1}(s, 1, m-1) \\
&= 2^{2m} \cdot 3 \cdot 2^{k+s-1} + 8 \cdot [21 \cdot 2^{k+s+3m-8} + 13 \cdot 2^{k+s+2m-6}] + 2^{2m-4} \cdot 2^{k+s} + 2^{2m-2} \cdot (-3 \cdot 2^{k+s-1}) \\
&= 21 \cdot 2^{k+s+3m-5} + 45 \cdot 2^{k+s+2m-4}
\end{aligned}$$

□

11.4. Computation of $\Delta_k \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{s+j-1}^{\left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right]}$ for $0 \leq j \leq s-1$, $l \geq 0$.

Lemma 11.3. *Let j and l be integers such that $2 \leq j \leq s-1$, $l \geq 0$.*

We have the following reduction formula for $2 \leq q \leq s-2$

(11.3)

$$\begin{aligned}
\Delta_k \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{s+j-1}^{\left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right]} &= 4^{2q+2} \cdot \Delta_k \Gamma_{s+j-2q-2}^{\left[\begin{smallmatrix} s-q-1 \\ s-q-1 \end{smallmatrix} \right]} - 8 \cdot 4^{2q+2} \cdot \Delta_k \Gamma_{s+j-2q-3}^{\left[\begin{smallmatrix} s-q-1 \\ s-q-1 \end{smallmatrix} \right]} \\
&\quad + \sum_{i=0}^q 4^{2i} \cdot \Delta_k \omega_{s-i+(j-i)}(s-i, 0, l+i) + \sum_{i=0}^q 4^{2i+1} \cdot \Delta_k \omega_{s-i-1+(j-i)}(s-i-1, 1, l+i)
\end{aligned}$$

$$\text{Setting } \begin{cases} q = j & \text{if } 2 \leq j \leq s-3, \\ q = s-3 & \text{if } j = s-2, \\ q = s-2 & \text{if } j = s-1. \end{cases}$$

in (11.3) we obtain

$$(11.4) \quad \Delta_k \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{s+j-1}^{\left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right]} = \begin{cases} 3 \cdot 2^{k+s-1} & \text{if } j = 0, l \geq 0. \\ 15 \cdot 2^{k+s-1} & \text{if } j = 1, l \geq 0. \\ 63 \cdot 2^{k+s+3j-6} & \text{if } 2 \leq j \leq s-1, l \geq 0. \end{cases}$$

Proof. From (8.3) we obtain successively the followings equations

$$\begin{aligned}
& \Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s+l \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s+l \end{bmatrix}} = 8 \cdot \left[\Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s \\ s+l-1 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-2}^{\begin{bmatrix} s-1 \\ s+l-1 \end{bmatrix}} \right] + \Delta_k \omega_{s+j}(s, 0, l) \\
& 4 \cdot \left[\Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s-1 \\ s+l \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-2}^{\begin{bmatrix} s-1 \\ s+l \end{bmatrix}} \right] = 4 \cdot \left[8 \cdot \left[\Delta_k \Gamma_{s+j-2}^{\begin{bmatrix} s-1 \\ s+l-1 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-3}^{\begin{bmatrix} s-1 \\ s+l-1 \end{bmatrix}} \right] + \Delta_k \omega_{s+j-1}(s-1, 1, l) \right] \\
& 4^2 \cdot \left[\Delta_k \Gamma_{s+j-2}^{\begin{bmatrix} s-1 \\ s+l \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-3}^{\begin{bmatrix} s-2 \\ s+l \end{bmatrix}} \right] = 4^2 \cdot \left[8 \cdot \left[\Delta_k \Gamma_{s+j-3}^{\begin{bmatrix} s-1 \\ s+l-1 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-4}^{\begin{bmatrix} s-2 \\ s+l-1 \end{bmatrix}} \right] + \Delta_k \omega_{s+j-2}(s-1, 0, l+1) \right] \\
& \vdots \\
& 4^{2q} \cdot \left[\Delta_k \Gamma_{s+j-2q}^{\begin{bmatrix} s-q \\ s+l \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-2q-1}^{\begin{bmatrix} s-q-1 \\ s+l \end{bmatrix}} \right] = 4^{2q} \cdot \left[8 \cdot \left[\Delta_k \Gamma_{s+j-2q-1}^{\begin{bmatrix} s-q \\ s+l-1 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-2q-2}^{\begin{bmatrix} s-q-1 \\ s+l-1 \end{bmatrix}} \right] + \Delta_k \omega_{s+j-2q}(s-q, 0, l+q) \right] \\
& 4^{2q+1} \cdot \left[\Delta_k \Gamma_{s+j-2q-1}^{\begin{bmatrix} s-q-1 \\ s+l \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-2q-2}^{\begin{bmatrix} s-q-1 \\ s+l \end{bmatrix}} \right] \\
& = 4^{2q+1} \cdot \left[8 \cdot \left[\Delta_k \Gamma_{s+j-2q-2}^{\begin{bmatrix} s-q-1 \\ s+l-1 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{s+j-2q-3}^{\begin{bmatrix} s-q-1 \\ s+l-1 \end{bmatrix}} \right] + \Delta_k \omega_{s+j-2q-1}(s-q-1, 1, l+q) \right]
\end{aligned}$$

By summing the above equations we obtain after reduction (11.3).

$$\begin{aligned}
& \Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s+l \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s \\ s+l-1 \end{bmatrix}} = 4^{2q+2} \cdot \Delta_k \Gamma_{s+j-2q-2}^{\begin{bmatrix} s-q-1 \\ s+l \end{bmatrix}} - 8 \cdot 4^{2q+2} \cdot \Delta_k \Gamma_{s+j-2q-3}^{\begin{bmatrix} s-q-1 \\ s+l-1 \end{bmatrix}} \\
& + \sum_{i=0}^q 4^{2i} \cdot \Delta_k \omega_{s-i+(j-i)}(s-i, 0, l+i) + \sum_{i=0}^q 4^{2i+1} \cdot \Delta_k \omega_{s-i-1+(j-i)}(s-i-1, 1, l+i)
\end{aligned}$$

The two first cases $j=0, l \geq 0$ and $j=1, l \geq 0$ follows from (9.1) and (9.2)

The case $2 \leq j \leq s-4$

Setting $q=j$ **in the reduction formula** (11.3) **we obtain using Lemma 6.3 and results in Lemma 7.4**

$$\begin{aligned}
& \Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s+l \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s \\ s+l-1 \end{bmatrix}} - 4^{2j+2} \cdot \Delta_k \Gamma_{s-j-2}^{\begin{bmatrix} s-j-1 \\ s+l \end{bmatrix}} \\
& = -8 \cdot 4^{2j+2} \cdot \Delta_k \Gamma_{s-j-3}^{\begin{bmatrix} s-j-1 \\ s+l-1 \end{bmatrix}} + \sum_{i=0}^j 4^{2i} \cdot \Delta_k \omega_{s-i+(j-i)}(s-i, 0, l+i) + \sum_{i=0}^j 4^{2i+1} \cdot \Delta_k \omega_{s-i-1+(j-i)}(s-i-1, 1, l+i)
\end{aligned}$$

we have

$$\Delta_k \omega_{s-i+(j-i)}(s-i, 0, l+i) = 0 \Leftrightarrow s-i+3 \leq s-i+(j-i) \leq 2 \cdot (s-i) - 1 \Leftrightarrow i \leq j-3$$

$$\Delta_k \omega_{s-i-1+(j-i)}(s-i-1, 1, l+i) = 0 \Leftrightarrow s-i-1+3 \leq s-i-1+(j-i) \leq 1+2 \cdot (s-i-1) - 1 \Leftrightarrow i \leq j-3$$

We then deduce

$$\begin{aligned}
& \Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s+l \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+j-1}^{\begin{bmatrix} s \\ s+l-1 \end{bmatrix}} \\
& = \sum_{i=j-2}^j 4^{2i} \cdot \Delta_k \omega_{s-i+(j-i)}(s-i, 0, l+i) + \sum_{i=j-2}^j 4^{2i+1} \cdot \Delta_k \omega_{s-i-1+(j-i)}(s-i-1, 1, l+i) \\
& = 2^{4(j-2)} \cdot \Delta_k \omega_{s-j+2+2}(s-j+2, 0, l+j-2) + 2^{4(j-1)} \cdot \Delta_k \omega_{s-j+1+1}(s-j+1, 0, l+j-1)
\end{aligned}$$

$$\begin{aligned}
& + 2^{4j} \cdot \Delta_k \omega_{s-j}(s-j, 0, l+j) + 2^{4(j-2)+2} \cdot \Delta_k \omega_{s-j+1+2}(s-j+1, 1, l+j-2) \\
& 2^{4(j-1)+2} \cdot \Delta_k \omega_{s-j+1}(s-j, 1, l+j-1) + 2^{4j+2} \cdot \Delta_k \omega_{s-j-1}(s-j-1, 1, l+j) \\
& = 2^{4j-8} \cdot 2^{k+s-j+2} + (-3 \cdot 2^{k+(s-j+1)-1}) \cdot 2^{4j-4} + 2^{4j} \cdot 2^{k+s-j-1} + 2^{4j-6} \cdot 2^{k+s-j+1} + (-3 \cdot 2^{k+(s-j)-1}) \cdot 2^{4j-2} \\
& + 2^{4j+2} \cdot 2^{k+(s-j-1)-1} = 21 \cdot 2^{k+s+3j-6} + 21 \cdot 2^{k+s+3j-5} = 63 \cdot 2^{k+s+3j-6}
\end{aligned}$$

The case $j = s - 3$

Setting $q = s - 3$ in the reduction formula (11.3) we obtain similarly

$$\begin{aligned}
& \Delta_k \Gamma_{2s-3} \left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_{2s-4} \left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right] - 4^{2s-4} \cdot \Delta_k \Gamma_1 \left[\begin{smallmatrix} 2 \\ s+l \end{smallmatrix} \right] \\
& = -8 \cdot 4^{2s-4} \cdot \Delta_k \Gamma_0 \left[\begin{smallmatrix} 2 \\ s+l-1 \end{smallmatrix} \right] + \sum_{i=0}^{s-3} 4^{2i} \cdot \Delta_k \omega_{2(s-i)-3}(s-i, 0, l+i) + \sum_{i=0}^{s-3} 4^{2i+1} \cdot \Delta_k \omega_{2(s-i-1)-2}(s-i-1, 1, l+i)
\end{aligned}$$

we have

$$\Delta_k \omega_{2(s-i)-3}(s-i, 0, l+i) = 0 \Leftrightarrow s-i+3 \leq 2(s-i) - 3 \leq 2 \cdot (s-i) - 1 \Leftrightarrow i \leq s-6$$

$$\Delta_k \omega_{2(s-i-1)-2}(s-i-1, 1, l+i) = 0 \Leftrightarrow s-i-1+3 \leq 2(s-i-1) - 2 \leq 1+2 \cdot (s-i-1) - 1 \Leftrightarrow i \leq s-6$$

We then deduce

$$\begin{aligned}
& \Delta_k \Gamma_{2s-3} \left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_{2s-4} \left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right] \\
& = \sum_{i=s-5}^{s-3} 4^{2i} \cdot \Delta_k \omega_{2(s-i)-3}(s-i, 0, l+i) + \sum_{i=s-5}^{s-3} 4^{2i+1} \cdot \Delta_k \omega_{2(s-i-1)-2}(s-i-1, 1, l+i) \\
& = 2^{4(s-5)} \cdot \Delta_k \omega_7(5, 0, l+s-5) + 2^{4(s-4)} \cdot \Delta_k \omega_5(4, 0, l+s-4) + 2^{4(s-3)} \cdot \Delta_k \omega_3(3, 0, l+s-3) \\
& + 2^{4(s-5)+2} \cdot \Delta_k \omega_6(4, 1, l+s-5) + 2^{4(s-4)+2} \cdot \Delta_k \omega_4(3, 1, l+s-4) \\
& + 2^{4(s-3)+2} \cdot \Delta_k \omega_2(2, 1, l+s-3) \\
& = 2^{4s-20} \cdot 2^{k+5} + (-3 \cdot 2^{k+3}) \cdot 2^{4s-16} + 2^{4s-12} \cdot 2^{k+2} + 2^{4s-18} \cdot 2^{k+4} + (-3 \cdot 2^{k+2}) \cdot 2^{4s-14} + 2^{4s-10} \cdot 2^{k+1} \\
& = 21 \cdot 2^{k+4s-15} + 21 \cdot 2^{k+4s-14} = 63 \cdot 2^{k+4s-15}
\end{aligned}$$

The case $j = s - 2$

Setting $q = s - 3$ in the reduction formula (11.3) we obtain similarly

$$\begin{aligned}
& \Delta_k \Gamma_{2s-2} \left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_{2s-3} \left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right] - 4^{2s-4} \cdot \Delta_k \Gamma_2 \left[\begin{smallmatrix} 2 \\ s+l \end{smallmatrix} \right] \\
& = -8 \cdot 4^{2s-4} \cdot \Delta_k \Gamma_1 \left[\begin{smallmatrix} 2 \\ s+l-1 \end{smallmatrix} \right] + \sum_{i=0}^{s-3} 4^{2i} \cdot \Delta_k \omega_{2(s-i)-2}(s-i, 0, l+i) + \sum_{i=0}^{s-3} 4^{2i+1} \cdot \Delta_k \omega_{2(s-i-1)-1}(s-i-1, 1, l+i)
\end{aligned}$$

we have

$$\Delta_k \omega_{2(s-i)-2}(s-i, 0, l+i) = 0 \Leftrightarrow s-i+3 \leq 2(s-i) - 2 \leq 2 \cdot (s-i) - 1 \Leftrightarrow i \leq s-5$$

$$\Delta_k \omega_{2(s-i-1)-1}(s-i-1, 1, l+i) = 0 \Leftrightarrow s-i-1+3 \leq 2(s-i-1) - 1 \leq 1+2 \cdot (s-i-1) - 1 \Leftrightarrow i \leq s-5$$

$$\Delta_k \Gamma_2 \left[\begin{smallmatrix} 2 \\ s+l \end{smallmatrix} \right] = 3 \cdot 2^{k+1}$$

We then deduce

$$\begin{aligned}
& \Delta_k \Gamma_{2s-2} \left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_{2s-3} \left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right] \\
& = 4^{2s-4} \cdot \Delta_k \Gamma_2 \left[\begin{smallmatrix} 2 \\ s+l \end{smallmatrix} \right] + \sum_{i=s-4}^{s-3} 4^{2i} \cdot \Delta_k \omega_{2(s-i)-2}(s-i, 0, l+i) + \sum_{i=s-4}^{s-3} 4^{2i+1} \cdot \Delta_k \omega_{2(s-i-1)-1}(s-i-1, 1, l+i) \\
& = 2^{4s-8} \cdot 3 \cdot 2^{k+1} + 2^{4(s-4)} \cdot \Delta_k \omega_6(4, 0, l+s-4) + 2^{4(s-3)} \cdot \Delta_k \omega_4(3, 0, l+s-3)
\end{aligned}$$

$$\begin{aligned}
& + 2^{4(s-4)+2} \cdot \Delta_k \omega_5(3, 1, l + s - 4) + 2^{4(s-3)+2} \cdot \Delta_k \omega_3(2, 1, l + s - 3) \\
& = 2^{4s-8} \cdot 3 \cdot 2^{k+1} + 2^{4s-16} \cdot 2^{k+4} + (-3 \cdot 2^{k+2}) \cdot 2^{4s-12} + 2^{4s-14} \cdot 2^{k+3} + (-3 \cdot 2^{k+1}) \cdot 2^{4s-10} \\
& = 63 \cdot 2^{k+4s-12}
\end{aligned}$$

The case $j = s - 1$

Setting $q = s - 2$ in the reduction formula (11.3) we obtain similarly

$$\begin{aligned}
& \Delta_k \Gamma_{2s-1} \left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_{2s-2} \left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right] - 4^{2s-2} \cdot \Delta_k \Gamma_1 \left[\begin{smallmatrix} 1 \\ s+l \end{smallmatrix} \right] \\
& = -8 \cdot 4^{2s-2} \cdot \Delta_k \Gamma_0 \left[\begin{smallmatrix} 1 \\ s+l-1 \end{smallmatrix} \right] + \sum_{i=0}^{s-2} 4^{2i} \cdot \Delta_k \omega_{2(s-i)-1}(s-i, 0, l+i) + \sum_{i=0}^{s-2} 4^{2i+1} \cdot \Delta_k \omega_{2(s-i-1)}(s-i-1, 1, l+i)
\end{aligned}$$

we have

$$\Delta_k \omega_{2(s-i)-1}(s-i, 0, l+i) = 0 \Leftrightarrow s-i+3 \leq 2(s-i)-1 \leq 2 \cdot (s-i)-1 \Leftrightarrow i \leq s-4$$

$$\Delta_k \omega_{2(s-i-1)}(s-i-1, 1, l+i) = 0 \Leftrightarrow s-i-1+3 \leq 2(s-i-1) \leq 1+2 \cdot (s-i-1)-1 \Leftrightarrow i \leq s-4, s-i-1 \geq 4$$

$$\Delta_k \Gamma_1 \left[\begin{smallmatrix} 1 \\ s+l \end{smallmatrix} \right] = 3 \cdot 2^k$$

We then deduce

$$\begin{aligned}
& \Delta_k \Gamma_{2s-1} \left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_{2s-2} \left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right] \\
& = 4^{2s-2} \cdot \Delta_k \Gamma_1 \left[\begin{smallmatrix} 1 \\ s+l \end{smallmatrix} \right] + \sum_{i=s-3}^{s-2} 4^{2i} \cdot \Delta_k \omega_{2(s-i)-1}(s-i, 0, l+i) + \sum_{i=s-4}^{s-2} 4^{2i+1} \cdot \Delta_k \omega_{2(s-i-1)}(s-i-1, 1, l+i) \\
& = 2^{4s-4} \cdot 3 \cdot 2^k + 2^{4(s-3)} \cdot \Delta_k \omega_5(3, 0, l+s-3) + 2^{4(s-2)} \cdot \Delta_k \omega_3(2, 0, l+s-2) \\
& + 2^{4(s-4)+2} \cdot \Delta_k \omega_6(3, 1, l+s-4) + 2^{4(s-3)+2} \cdot \Delta_k \omega_4(2, 1, l+s-3) + 2^{4(s-2)+2} \cdot \Delta_k \omega_2(1, 1, l+s-2) \\
& = 2^{4s-4} \cdot 3 \cdot 2^k + 2^{4s-12} \cdot 2^{k+3} + (-3 \cdot 2^{k+1}) \cdot 2^{4s-8} + 2^{4s-14} \cdot 0 + 2^{4s-10} \cdot 2^{k+2} + (-3 \cdot 2^k) \cdot 2^{4s-6} \\
& = 2^{k+4s-9} \cdot [1 + 3 \cdot 2^5 - 3 \cdot 2^2 + 2 - 3 \cdot 2^3] = 63 \cdot 2^{k+4s-9}
\end{aligned}$$

□

$$11.5. \text{ Computation of } \Delta_k \Gamma_{s+m+j} \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_{s+m+j-1} \left[\begin{smallmatrix} s \\ s+m+l-1 \end{smallmatrix} \right] \quad \text{for } 0 \leq j \leq s-1, m \geq 1, l \geq 0.$$

Lemma 11.4. *We have for $m \geq 1, l \geq 0$ the following reduction formula*

(11.5)

$$\begin{aligned}
& \Delta_k \Gamma_{s+m+j} \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_{s+m+j-1} \left[\begin{smallmatrix} s \\ s+m+l-1 \end{smallmatrix} \right] \\
& = 2^{2m} \cdot \left[\Delta_k \Gamma_{s+j} \left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_{s+j-1} \left[\begin{smallmatrix} s \\ s+m+l-1 \end{smallmatrix} \right] \right] + \sum_{i=0}^{m-1} 2^{2i} \cdot \Delta_k \omega_{s+m+j-i}(s, m-i, l+i)
\end{aligned}$$

where

(11.6)

$$\sum_{i=0}^{m-1} 2^{2i} \cdot \Delta_k \omega_{s+m+j-i}(s, m-i, l+i) = \begin{cases} (-3) \cdot 2^{k+s-1} & \text{if } j = 0, m = 1, l \geq 0 \\ (-5) \cdot 2^{k+s+2m-4} & \text{if } j = 0, m \geq 2, l \geq 0 \\ 2^{k+s+2m-2} & \text{if } j = 1, m \geq 1, l \geq 0 \\ 0 & \text{if } 2 \leq j \leq s-1, m \geq 1, l \geq 0. \end{cases}$$

We get from (11.4) applying the reduction formula (11.5):
(11.7)

$$\Delta_k \Gamma_{s+m+j}^{\left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{s+m+j-1}^{\left[\begin{smallmatrix} s \\ s+m+l-1 \end{smallmatrix} \right]} = \begin{cases} 9 \cdot 2^{k+s-1} & \text{if } j = 0, m = 1, l \geq 0 \\ 19 \cdot 2^{k+s+2m-4} & \text{if } j = 0, m \geq 2, l \geq 0 \\ 31 \cdot 2^{k+s+2m-2} & \text{if } j = 1, m \geq 1, l \geq 0 \\ 63 \cdot 2^{k+s+2m+3j-6} & \text{if } 2 \leq j \leq s-1, m \geq 0, l \geq 0. \end{cases}$$

Proof. Let q be a rational integer between one and $m-1$. From (8.3) we obtain successively the following equations

$$\begin{aligned} \Delta_k \Gamma_{s+m+j}^{\left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{s+m+j-1}^{\left[\begin{smallmatrix} s \\ s+m+l-1 \end{smallmatrix} \right]} &= 8 \cdot \left[\Delta_k \Gamma_{s+m+j-1}^{\left[\begin{smallmatrix} s \\ s+m+l-1 \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{s+m+j-2}^{\left[\begin{smallmatrix} s \\ s+m+l-2 \end{smallmatrix} \right]} \right] + \Delta_k \omega_{s+m+j}(s, m, l) \\ 4 \cdot \left[\Delta_k \Gamma_{s+m+j-1}^{\left[\begin{smallmatrix} s \\ s+m+l-1 \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{s+m+j-2}^{\left[\begin{smallmatrix} s \\ s+m+l-2 \end{smallmatrix} \right]} \right] &= 4 \cdot \left[8 \cdot \left[\Delta_k \Gamma_{s+m+j-2}^{\left[\begin{smallmatrix} s \\ s+m+l-2 \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{s+m+j-3}^{\left[\begin{smallmatrix} s \\ s+m+l-3 \end{smallmatrix} \right]} \right] + \Delta_k \omega_{s+m+j-1}(s, m-1, l+1) \right] \\ 4^2 \cdot \left[\Delta_k \Gamma_{s+m+j-2}^{\left[\begin{smallmatrix} s \\ s+m+l-2 \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{s+m+j-3}^{\left[\begin{smallmatrix} s \\ s+m+l-3 \end{smallmatrix} \right]} \right] &= 4^2 \cdot \left[8 \cdot \left[\Delta_k \Gamma_{s+m+j-3}^{\left[\begin{smallmatrix} s \\ s+m+l-3 \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{s+m+j-4}^{\left[\begin{smallmatrix} s \\ s+m+l-4 \end{smallmatrix} \right]} \right] + \Delta_k \omega_{s+m+j-2}(s, m-2, l+2) \right] \\ &\vdots \\ 4^q \cdot \left[\Delta_k \Gamma_{s+m+j-q}^{\left[\begin{smallmatrix} s \\ s+m+l-q \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{s+m+j-(q+1)}^{\left[\begin{smallmatrix} s \\ s+m+l-(q+1) \end{smallmatrix} \right]} \right] \\ &= 4^q \cdot \left[8 \cdot \left[\Delta_k \Gamma_{s+m+j-(q+1)}^{\left[\begin{smallmatrix} s \\ s+m+l-(q+1) \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{s+m+j-(q+2)}^{\left[\begin{smallmatrix} s \\ s+m+l-(q+2) \end{smallmatrix} \right]} \right] + \Delta_k \omega_{s+m+j-q}(s, m-q, l+q) \right] \end{aligned}$$

By summing the above equations we obtain after reduction the following formula

$$\begin{aligned} \Delta_k \Gamma_{s+m+j}^{\left[\begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{s+m+j-1}^{\left[\begin{smallmatrix} s \\ s+m+l-1 \end{smallmatrix} \right]} \\ = 4^{q+1} \cdot \Delta_k \Gamma_{s+m+j-(q+1)}^{\left[\begin{smallmatrix} s \\ s+m+l-(q+1) \end{smallmatrix} \right]} - 8 \cdot 4^{q+1} \cdot \Delta_k \Gamma_{s+m+j-(q+2)}^{\left[\begin{smallmatrix} s \\ s+m+l-(q+2) \end{smallmatrix} \right]} + \sum_{i=0}^q 4^i \cdot \Delta_k \omega_{s+m+j-i}(s, m-i, l+i) \end{aligned}$$

Setting $q = m-1$ in the above formula we get the reduction formula (11.5)

Proof of (11.6) The proof is based on results in Lemma 7.4

The case $j = 0, m = 1, l \geq 0$

$$\sum_{i=0}^{m-1} 2^{2i} \cdot \Delta_k \omega_{s+m+j-i}(s, m-i, l+i) = \Delta_k \omega_{s+1}(s, 1, l) = (-3) \cdot 2^{k+s-1}$$

The case $j = 0, m \geq 2, l \geq 0$

$$\begin{aligned} \sum_{i=0}^{m-1} 2^{2i} \cdot \Delta_k \omega_{s+m+j-i}(s, m-i, l+i) &= \sum_{i=m-2}^{m-1} 2^{2i} \cdot \Delta_k \omega_{s+m-i}(s, m-i, l+i) \\ &= 2^{2(m-2)} \cdot \Delta_k \omega_{s+2}(s, 2, l+m-2) + 2^{2(m-1)} \cdot \Delta_k \omega_{s+1}(s, 1, l+m-1) \\ &= 2^{2(m-2)} \cdot 2^{k+s} + 2^{2(m-1)} \cdot (-3 \cdot 2^{k+s-1}) = (-5) \cdot 2^{k+s+2m-4} \end{aligned}$$

Since $\Delta_k \omega_{s+m-i}(s, m-i, l+i) = 0 \iff i \leq m-3$

The case $j = 1, m \geq 1, l \geq 0$

$$\begin{aligned} \sum_{i=0}^{m-1} 2^{2i} \cdot \Delta_k \omega_{s+m+j-i}(s, m-i, l+i) &= \sum_{i=m-1}^{m-1} 2^{2i} \cdot \Delta_k \omega_{s+m+1-i}(s, m-i, l+i) \\ &= 2^{2(m-1)} \cdot \Delta_k \omega_{s+2}(s, 1, l+m-1) = 2^{2(m-1)} \cdot 2^{k+s} = 2^{k+s+2m-2} \end{aligned}$$

Since $\Delta_k \omega_{s+m+1-i}(s, m-i, l+i) = 0 \iff i \leq m-2$

The case $2 \leq j \leq s-1, m \geq 1, l \geq 0$

$$\sum_{i=0}^{m-1} 2^{2i} \cdot \Delta_k \omega_{s+m+j-i}(s, m-i, l+i) = 0$$

Since $\Delta_k \omega_{s+m+j-i}(s, m-i, l+i) = 0 \iff 0 \leq i \leq m-1$

Proof of (11.7)

We deduce easily (11.7) from the reduction formula (11.5) using (11.6) and (11.4) with $l \longrightarrow m+l$ $\square$

11.6. **Computation of $\Delta_k \Gamma_{s+m+1}^{\begin{bmatrix} s \\ s+m \end{bmatrix}}$ for $m \geq 0$.**

Lemma 11.5. *We have*

$$(11.8) \quad \Delta_k \Gamma_{s+m+1}^{\begin{bmatrix} s \\ s+m \end{bmatrix}} = \begin{cases} 147 \cdot 2^{k+s-1} & \text{if } m = 0 \\ 315 \cdot 2^{k+s} & \text{if } m = 1 \\ 21 \cdot 2^{k+s+3m-2} + 273 \cdot 2^{k+s+2m-2} & \text{if } m \geq 2 \text{ (holds for } m \geq 0) \end{cases}$$

Proof. From (8.3) and, applying (11.7) in the case $j = 1, m \geq 1, l = 0$:

$$\begin{aligned} &\Delta_k \Gamma_{s+m+1}^{\begin{bmatrix} s \\ s+m \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+m}^{\begin{bmatrix} s \\ s+m-1 \end{bmatrix}} \\ &= 4 \cdot \left[\Delta_k \Gamma_{s+m}^{\begin{bmatrix} s \\ s+m-1 \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+m-1}^{\begin{bmatrix} s \\ s+m-1 \end{bmatrix}} \right] + \Delta_k \omega_{s+m+1}(s, m, 0) = 31 \cdot 2^{k+s+2m-2} \\ &\Delta_k \Gamma_{s+m}^{\begin{bmatrix} s \\ s+m-1 \end{bmatrix}} = 8 \cdot \Delta_k \Gamma_{s+m-1}^{\begin{bmatrix} s \\ s+m-1 \end{bmatrix}} + \frac{1}{4} \cdot (31 \cdot 2^{k+s+2m-2} - \Delta_k \omega_{s+m+1}(s, m, 0)) \end{aligned}$$

Combining the above equations we have by (11.2) with $m \longrightarrow m-1$ and (7.48) in the case $m \geq 2$

$$\begin{aligned} \Delta_k \Gamma_{s+m+1}^{\begin{bmatrix} s \\ s+m \end{bmatrix}} &= 8 \cdot \Delta_k \Gamma_{s+m}^{\begin{bmatrix} s \\ s+m-1 \end{bmatrix}} + 31 \cdot 2^{k+s+2m-2} \\ &= 8 \cdot \left(8 \cdot \Delta_k \Gamma_{s+m-1}^{\begin{bmatrix} s \\ s+m-1 \end{bmatrix}} + \frac{1}{4} \cdot (31 \cdot 2^{k+s+2m-2} - \Delta_k \omega_{s+m+1}(s, m, 0)) \right) + 31 \cdot 2^{k+s+2m-2} \\ &= 2^6 \cdot [21 \cdot 2^{k+s+3(m-1)-5} + 45 \cdot 2^{k+s+2(m-1)-4}] + 93 \cdot 2^{k+s+2m-2} \quad \text{since } \Delta_k \omega_{s+m+1}(s, m, 0) = 0 \\ &= 21 \cdot 2^{k+s+3m-2} + 273 \cdot 2^{k+s+2m-2} \end{aligned}$$

similarly in the case $m = 1$

$$\begin{aligned} \Delta_k \Gamma_{s+2}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}} &= 8 \cdot \Delta_k \Gamma_{s+1}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}} + 31 \cdot 2^{k+s} \\ &= 8 \cdot \left(8 \cdot \Delta_k \Gamma_s^{\begin{bmatrix} s \\ s \end{bmatrix}} + \frac{1}{4} \cdot (31 \cdot 2^{k+s} - \Delta_k \omega_{s+2}(s, 1, 0)) \right) + 31 \cdot 2^{k+s} \end{aligned}$$

$$= 2^6 \cdot 7 \cdot 2^{k+s-1} + 91 \cdot 2^{k+s} = 315 \cdot 2^{k+s} \quad \text{since } \Delta_k \omega_{s+2}(s, 1, 0) = 2^{k+s}$$

□

11.7. **Computation of** $\Delta_k \Gamma_{2s+j}^{\left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{2s+j-1}^{\left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right]}$ for $0 \leq j \leq l$.

Lemma 11.6. We have for $l \geq 3$ the following reduction formula holding for $0 \leq j \leq l$.

(11.9)

$$\begin{aligned} \Delta_k \Gamma_{2s+j}^{\left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{2s+j-1}^{\left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right]} &= 4^{2s-2} \cdot \Delta_k \Gamma_{2+j}^{\left[\begin{smallmatrix} 1 \\ s+l \end{smallmatrix} \right]} - 8 \cdot 4^{2s-2} \cdot \Delta_k \Gamma_{1+j}^{\left[\begin{smallmatrix} 1 \\ s+l-1 \end{smallmatrix} \right]} \\ &+ \sum_{i=0}^{s-2} 4^{2i} \cdot \Delta_k \omega_{2(s-i)+j}(s-i, 0, l+i) + \sum_{i=0}^{s-2} 4^{2i+1} \cdot \Delta_k \omega_{2(s-i-1)+(j+1)}(s-i-1, 1, l+i) \end{aligned}$$

where

$$(11.10) \quad \Delta_k \Gamma_{2+j}^{\left[\begin{smallmatrix} 1 \\ s+l \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{1+j}^{\left[\begin{smallmatrix} 1 \\ s+l-1 \end{smallmatrix} \right]} = \begin{cases} 3 \cdot 2^{2k} - 9 \cdot 2^k & \text{if } j = 0, \\ -15 \cdot 2^{2k} + 3 \cdot 2^{k+1} & \text{if } j = 1, \\ -9 \cdot 2^{2k+2j-2} & \text{if } 2 \leq j \leq l, \end{cases}$$

(11.11)

$$\sum_{i=0}^{s-2} 4^{2i} \cdot \Delta_k \omega_{2(s-i)+j}(s-i, 0, l+i) = \begin{cases} -2^{2k+4s-4} + 2^{2k+2s-2} + 2^{k+4s-6} & \text{if } j = 0, \\ 5 \cdot 2^{2k+4s-4} - 5 \cdot 2^{2k+2s-2} & \text{if } j = 1, \\ 3 \cdot 2^{2k+4s+2j-6} - 3 \cdot 2^{2k+2s+2j-4} & \text{if } 2 \leq j \leq l-1, \\ 3 \cdot 2^{2k+4s+2l-6} - 27 \cdot 2^{2k+2s+2l-4} & \text{if } j = l, \end{cases}$$

(11.12)

$$\sum_{i=0}^{s-2} 4^{2i+1} \cdot \Delta_k \omega_{2(s-i-1)+j+1}(s-i-1, 1, l+i) = \begin{cases} -2^{2k+4s-3} + 2^{2k+2s-1} + 2^{k+4s-5} & \text{if } j = 0, \\ 5 \cdot 2^{2k+4s-3} - 5 \cdot 2^{2k+2s-1} & \text{if } j = 1, \\ 3 \cdot 2^{2k+4s+2j-5} - 3 \cdot 2^{2k+2s+2j-3} & \text{if } 2 \leq j \leq l-1, \\ 3 \cdot 2^{2k+4s+2l-5} - 27 \cdot 2^{2k+2s+2l-3} & \text{if } j = l. \end{cases}$$

Combining (11.9), (11.10), (11.11) and (11.12) we obtain

$$(11.13) \quad \Delta_k \Gamma_{2s+j}^{\left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{2s+j-1}^{\left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right]} = \begin{cases} 3 \cdot 2^{2k+2s-2} - 33 \cdot 2^{k+4s-6} & \text{if } j = 0 \\ -15 \cdot 2^{2k+2s-2} + 3 \cdot 2^{k+4s-3} & \text{if } j = 1 \\ -9 \cdot 2^{2k+2s+2j-4} & \text{if } 2 \leq j \leq l-1 \\ -81 \cdot 2^{2k+2s+2j-4} & \text{if } j = l \end{cases}$$

Proof. Proof of (11.9)

Let q be a rational integer between zero and $s-2$. From (8.3) we obtain successively the following equations.

$$\begin{aligned} \Delta_k \Gamma_{2s+j}^{\left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{2s+j-1}^{\left[\begin{smallmatrix} s-1 \\ s+l \end{smallmatrix} \right]} &= 8 \cdot \left[\Delta_k \Gamma_{2s+j-1}^{\left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{2s+j-2}^{\left[\begin{smallmatrix} s-1 \\ s+l-1 \end{smallmatrix} \right]} \right] + \Delta_k \omega_{2s+j}(s, 0, l) \\ 4 \cdot \left[\Delta_k \Gamma_{2s+j-1}^{\left[\begin{smallmatrix} s-1 \\ s+l \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{2s+j-2}^{\left[\begin{smallmatrix} s-1 \\ s+l \end{smallmatrix} \right]} \right] &= 4 \cdot \left[8 \cdot \left[\Delta_k \Gamma_{2s+j-2}^{\left[\begin{smallmatrix} s-1 \\ s+l-1 \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{2s+j-3}^{\left[\begin{smallmatrix} s-1 \\ s+l-1 \end{smallmatrix} \right]} \right] + \Delta_k \omega_{2s+j-1}(s-1, 1, l) \right] \end{aligned}$$

$$\begin{aligned}
& 4^2 \cdot \left[\Delta_k \Gamma_{2s+j-2}^{\begin{bmatrix} s-1 \\ s-1 \\ s+l \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{2s+j-3}^{\begin{bmatrix} s-2 \\ s-2 \\ s+l \end{bmatrix}} \right] = 4^2 \cdot \left[8 \cdot \left[\Delta_k \Gamma_{2s+j-3}^{\begin{bmatrix} s-1 \\ s-1 \\ s+l-1 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{2s+j-4}^{\begin{bmatrix} s-2 \\ s-1 \\ s+l-1 \end{bmatrix}} \right] + \Delta_k \omega_{2s+j-2}(s-1, 0, l+1) \right] \\
& \vdots \\
& 4^{2q} \cdot \left[\Delta_k \Gamma_{2s+j-2q}^{\begin{bmatrix} s-q \\ s-q \\ s+l \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{2s+j-2q-1}^{\begin{bmatrix} s-q \\ s-q \\ s+l \end{bmatrix}} \right] \\
& = 4^{2q} \cdot \left[8 \cdot \left[\Delta_k \Gamma_{2s+j-2q-1}^{\begin{bmatrix} s-q \\ s-q \\ s+l-1 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{2s+j-2q-2}^{\begin{bmatrix} s-q-1 \\ s-q \\ s+l-1 \end{bmatrix}} \right] + \Delta_k \omega_{2s+j-2q}(s-q, 0, l+q) \right] \\
& 4^{2q+1} \cdot \left[\Delta_k \Gamma_{2s+j-2q-1}^{\begin{bmatrix} s-q-1 \\ s-q \\ s+l \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{2s+j-2q-2}^{\begin{bmatrix} s-q-1 \\ s-q-1 \\ s+l \end{bmatrix}} \right] \\
& = 4^{2q+1} \cdot \left[8 \cdot \left[\Delta_k \Gamma_{2s+j-2q-2}^{\begin{bmatrix} s-q-1 \\ s-q \\ s+l-1 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{2s+j-2q-3}^{\begin{bmatrix} s-q-1 \\ s-q-1 \\ s+l-1 \end{bmatrix}} \right] + \Delta_k \omega_{2s+j-2q-1}(s-q-1, 1, l+q) \right]
\end{aligned}$$

By summing the above equations we obtain after reduction the following formula

$$\begin{aligned}
& \Delta_k \Gamma_{2s+j}^{\begin{bmatrix} s \\ s \\ s+l \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+j-1}^{\begin{bmatrix} s \\ s \\ s+l-1 \end{bmatrix}} \\
& = 4^{2q+2} \cdot \Delta_k \Gamma_{2(s-q-1)+j}^{\begin{bmatrix} s-q-1 \\ s-q-1 \\ s+l \end{bmatrix}} - 8 \cdot 4^{2q+2} \cdot \Delta_k \Gamma_{2(s-q-1)+(j-1)}^{\begin{bmatrix} s-q-1 \\ s-q-1 \\ s+l-1 \end{bmatrix}} \\
& + \sum_{i=0}^q 4^{2i} \cdot \Delta_k \omega_{2(s-i)+j}(s-i, 0, l+i) + \sum_{i=0}^q 4^{2i+1} \cdot \Delta_k \omega_{2(s-i-1)+(j+1)}(s-i-1, 1, l+i)
\end{aligned}$$

Setting $q = s - 2$ in the above formula we obtain the reduction formula (11.9)

Proof of (11.10) Let $2 + j < \inf(k, 2 + s + l)$. We obtain from Corollary 3.10 for $n = 2$ applying Theorem 3.1 (see [2] section 3)

$$(11.14) \quad \Gamma_{2+j}^{\begin{bmatrix} 1 \\ 1 \\ s+l \end{bmatrix} \times k} = \begin{cases} 2^{2k} + 15 \cdot 2^k + 158 & \text{if } j = 0 \\ 3 \cdot 2^{2k+2j-2} + 63 \cdot [2^{k+3j-2} + 5 \cdot 2^{4j-1}] & \text{if } 1 \leq j \leq l \end{cases}$$

$$(11.15) \quad \Gamma_{1+j}^{\begin{bmatrix} 1 \\ 1 \\ s+l-1 \end{bmatrix} \times k} = \begin{cases} 3 \cdot 2^k + 9 & \text{if } j = 0 \\ 2^{2k} + 15 \cdot 2^k + 158 & \text{if } j = 1 \\ 3 \cdot 2^{2k+2j-4} + 63 \cdot [2^{k+3j-5} + 5 \cdot 2^{4j-5}] & \text{if } 2 \leq j \leq l \end{cases}$$

We deduce (11.10) from (11.14) and (11.15).

Proof of (11.11) The proof is based on results in Lemma 7.4
The case $j = 0$,

$$\begin{aligned}
& \Delta_k \omega_{2(s-i)}(s-i, 0, l+i) = -3 \cdot 2^{2k+2(s-i)-2} \quad \text{if } 0 \leq i \leq s-3, \\
& \Delta_k \omega_4(2, 0, l+(s-2)) = -3 \cdot 2^{2k+2} + 2^{k+2} \quad \text{if } i = s-2.
\end{aligned}$$

It follows that

$$\begin{aligned} \sum_{i=0}^{s-2} 4^{2i} \cdot \Delta_k \omega_{2(s-i)}(s-i, 0, l+i) &= \sum_{i=0}^{s-2} 2^{4i} \cdot (-3 \cdot 2^{2k+2(s-i)-2}) + 2^{k+2} \cdot 2^{4(s-2)} \\ &= -2^{2k+4s-4} + 2^{2k+2s-2} + 2^{k+4s-6}. \end{aligned}$$

The case $j = 1$,

$$\Delta_k \omega_{2(s-i)+1}(s-i, 0, l+i) = 15 \cdot 2^{2k+2(s-i)-2} \quad \text{if } 0 \leq i \leq s-2,$$

It follows that

$$\begin{aligned} \sum_{i=0}^{s-2} 4^{2i} \cdot \Delta_k \omega_{2(s-i)+1}(s-i, 0, l+i) &= \sum_{i=0}^{s-2} 2^{4i} \cdot (15 \cdot 2^{2k+2(s-i)-2}) \\ &= 5 \cdot 2^{2k+4s-4} - 5 \cdot 2^{2k+2s-2}. \end{aligned}$$

The case $2 \leq j \leq l-1$,

$$\Delta_k \omega_{2(s-i)+j}(s-i, 0, l+i) = 9 \cdot 2^{2k+2(s-i)+2j-4} \quad \text{if } 0 \leq i \leq s-2,$$

It follows that

$$\begin{aligned} \sum_{i=0}^{s-2} 4^{2i} \cdot \Delta_k \omega_{2(s-i)+j}(s-i, 0, l+i) &= \sum_{i=0}^{s-2} 2^{4i} \cdot (9 \cdot 2^{2k+2(s-i)+2j-4}) \\ &= 3 \cdot 2^{2k+4s+2j-6} - 3 \cdot 2^{2k+2s+2j-4}. \end{aligned}$$

The case $j = l$,

$$\Delta_k \omega_{2(s-i)+l}(s-i, 0, l+i) = 9 \cdot 2^{2k+2(s-i)+2l-4} \quad \text{if } 1 \leq i \leq s-2,$$

$$\Delta_k \omega_{2s+l}(s, 0, l) = -15 \cdot 2^{2k+2s+2l-4} \quad \text{if } i = 0,$$

It follows that

$$\begin{aligned} \sum_{i=0}^{s-2} 4^{2i} \cdot \Delta_k \omega_{2(s-i)+j}(s-i, 0, l+i) &= \sum_{i=1}^{s-2} 2^{4i} \cdot (9 \cdot 2^{2k+2(s-i)+2l-4}) + (-15 \cdot 2^{2k+2s+2l-4}) \\ &= 3 \cdot 2^{2k+4s+2l-6} - 27 \cdot 2^{2k+2s+2l-4}. \end{aligned}$$

Proof of (11.12)

Similarly to the proof of (11.11) applying (11.14), (11.15) and Lemma 1.22.

The case $j = 0$,

$$\Delta_k \omega_{2(s-i-1)+1}(s-i-1, 1, l+i) = -3 \cdot 2^{2k+2(s-i-1)-1} \quad \text{if } 0 \leq i \leq s-3,$$

$$\begin{aligned} \Delta_k \omega_3(1, 1, l+(s-2)) &= \Delta_k \Gamma_3 \left[\begin{smallmatrix} \frac{1}{2} \\ s+l \end{smallmatrix} \right] - 4 \cdot \Delta_k \Gamma_2 \left[\begin{smallmatrix} \frac{1}{1} \\ s+l \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_2 \left[\begin{smallmatrix} \frac{1}{2} \\ s+l-1 \end{smallmatrix} \right] + 32 \cdot \Delta_k \Gamma_1 \left[\begin{smallmatrix} \frac{1}{1} \\ s+l-1 \end{smallmatrix} \right] \\ &= -3 \cdot 2^{2k+1} + 2^{k+1} \quad \text{if } i = s-2. \end{aligned}$$

It follows that

$$\begin{aligned} \sum_{i=0}^{s-2} 4^{2i+1} \cdot \Delta_k \omega_{2(s-i-1)+1}(s-i-1, 1, l+i) &= \sum_{i=0}^{s-2} 2^{4i+2} \cdot (-3 \cdot 2^{2k+2(s-i-1)-1}) + 2^{k+1} \cdot 2^{4(s-2)+2} \\ &= -2^{2k+4s-3} + 2^{2k+2s-1} + 2^{k+4s-5}. \end{aligned}$$

The case $j = 1$,

$$\Delta_k \omega_{2(s-i-1)+2}(s-i-1, 1, l+i) = 15 \cdot 2^{2k+2(s-i-1)-1} \quad \text{if } 0 \leq i \leq s-2,$$

$$\Delta_k \omega_4(1, 1, l+s-2) = \Delta_k \Gamma_4 \left[\begin{smallmatrix} \frac{1}{2} \\ s+l \end{smallmatrix} \right] - 4 \cdot \Delta_k \Gamma_3 \left[\begin{smallmatrix} \frac{1}{1} \\ s+l \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_3 \left[\begin{smallmatrix} \frac{1}{2} \\ s+l-1 \end{smallmatrix} \right] + 32 \cdot \Delta_k \Gamma_2 \left[\begin{smallmatrix} \frac{1}{1} \\ s+l-1 \end{smallmatrix} \right]$$

$$= [9 \cdot 2^{2k+1} + 105 \cdot 2^{k+3}] - 4 \cdot [9 \cdot 2^{2k} + 63 \cdot 2^{k+1}] - 8 \cdot [3 \cdot 2^{2k+1} + 102 \cdot 2^k] + 32 \cdot [3 \cdot 2^{2k} + 15 \cdot 2^k] = 15 \cdot 2^{2k+1}$$

It follows that

$$\begin{aligned} \sum_{i=0}^{s-2} 4^{2i+1} \cdot \Delta_k \omega_{2(s-i-1)+2}(s-i-1, 1, l+i) &= \sum_{i=0}^{s-2} 2^{4i+2} \cdot (15 \cdot 2^{2k+2(s-i-1)-1}) \\ &= 5 \cdot 2^{2k+4s-3} - 5 \cdot 2^{2k+2s-1}. \end{aligned}$$

The case $2 \leq j \leq l-1$,

$$\begin{aligned} \Delta_k \omega_{2(s-i-1)+j+1}(s-i-1, 1, l+i) &= 9 \cdot 2^{2k+2(s-i)+2j-5} \quad \text{if } 0 \leq i \leq s-2, \\ \Delta_k \omega_{j+3}(1, 1, l+s-2) &= \Delta_k \Gamma_{j+3}^{\left[\begin{smallmatrix} \frac{1}{2} \\ s+l \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{j+2}^{\left[\begin{smallmatrix} \frac{1}{2} \\ s+l \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{j+2}^{\left[\begin{smallmatrix} \frac{1}{2} \\ s+l-1 \end{smallmatrix} \right]} + 32 \cdot \Delta_k \Gamma_{j+1}^{\left[\begin{smallmatrix} \frac{1}{2} \\ s+l-1 \end{smallmatrix} \right]} \\ &= [9 \cdot 2^{2k+2j-1} + 105 \cdot 2^{k+3j}] - 4 \cdot [9 \cdot 2^{2k+2j-2} + 63 \cdot 2^{k+3j-2}] \\ &\quad - 8 \cdot [9 \cdot 2^{2k+2j-3} + 105 \cdot 2^{k+3j-3}] + 32 \cdot [9 \cdot 2^{2k+2j-4} + 63 \cdot 2^{k+3j-5}] = 9 \cdot 2^{2k+2j-1} \end{aligned}$$

It follows that

$$\begin{aligned} \sum_{i=0}^{s-2} 4^{2i+1} \cdot \Delta_k \omega_{2(s-i-1)+j+1}(s-i-1, 1, l+i) &= \sum_{i=0}^{s-2} 2^{4i+2} \cdot (9 \cdot 2^{2k+2(s-i)+2j-5}) \\ &= 3 \cdot 2^{2k+4s+2j-5} - 3 \cdot 2^{2k+2s+2j-3}. \end{aligned}$$

The case $j = l$,

$$\begin{aligned} \Delta_k \omega_{2(s-i-1)+l+1}(s-i-1, 1, l+i) &= 9 \cdot 2^{2k+2(s-i)+2l-5} \quad \text{if } 1 \leq i \leq s-2, \\ \Delta_k \omega_{2(s-1)+l+1}(s-1, 1, l) &= -15 \cdot 2^{2k+2s+2l-5} \quad \text{if } i = 0, \\ \Delta_k \omega_{l+3}(1, 1, l+s-2) &= \Delta_k \Gamma_{l+3}^{\left[\begin{smallmatrix} \frac{1}{2} \\ s+l \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{l+2}^{\left[\begin{smallmatrix} \frac{1}{2} \\ s+l \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{l+2}^{\left[\begin{smallmatrix} \frac{1}{2} \\ s+l-1 \end{smallmatrix} \right]} + 32 \cdot \Delta_k \Gamma_{l+1}^{\left[\begin{smallmatrix} \frac{1}{2} \\ s+l-1 \end{smallmatrix} \right]} \\ &= [9 \cdot 2^{2k+2l-1} + 105 \cdot 2^{k+3l}] - 4 \cdot [9 \cdot 2^{2k+2l-2} + 63 \cdot 2^{k+3l-2}] \\ &\quad - 8 \cdot [9 \cdot 2^{2k+2l-3} + 105 \cdot 2^{k+3l-3}] + 32 \cdot [9 \cdot 2^{2k+2l-4} + 63 \cdot 2^{k+3l-5}] = 9 \cdot 2^{2k+2l-1} \end{aligned}$$

It follows that

$$\begin{aligned} \sum_{i=0}^{s-2} 4^{2i+1} \cdot \Delta_k \omega_{2(s-i-1)+l+1}(s-i-1, 1, l+i) &= \sum_{i=1}^{s-2} 2^{4i+2} \cdot (9 \cdot 2^{2k+2(s-i)+2l-5}) + (-15 \cdot 2^{2k+2s+2l-5}) \\ &= 3 \cdot 2^{2k+4s+2l-5} - 27 \cdot 2^{2k+2s+2l-3}. \end{aligned}$$

(11.13) is obtained as indicated in Lemma 11.6 □

$$11.8. \text{ Computation of } \Delta_k \Gamma_{2s+l+1+j}^{\left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{2s+l+j}^{\left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right]} \quad \text{for } 0 \leq j \leq s-1.$$

Lemma 11.7. *We have for $l \geq 3$ the following reduction formula holding for $0 \leq j \leq s-3$.*

(11.16)

$$\begin{aligned} \Delta_k \Gamma_{2s+l+1+j}^{\left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{2s+l+j}^{\left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right]} &= 2^{4j+8} \cdot \Delta_k \Gamma_{2(s-j-2)+l+j+1}^{\left[\begin{smallmatrix} s-j-2 \\ s-j-2 \\ s-j-2+(j+2+l) \end{smallmatrix} \right]} - 8 \cdot 2^{4j+8} \cdot \Delta_k \Gamma_{2(s-j-2)+l+j}^{\left[\begin{smallmatrix} s-j-2 \\ s-j-2 \\ s-j-2+(j+1+l) \end{smallmatrix} \right]} \\ &+ \sum_{i=0}^{j+1} 2^{4i} \cdot \Delta_k \omega_{2(s-i)+l+j+1}(s-i, 0, l+i) + \sum_{i=0}^{j+1} 2^{4i+2} \cdot \Delta_k \omega_{2(s-i-1)+l+j+2}(s-i-1, 1, l+i) \end{aligned}$$

where

$$(11.17) \quad \Delta_k \Gamma_{2(s-j-2)+l+j+1}^{\left[\begin{smallmatrix} s-j-2 \\ s-j-2 \\ s-j-2+(j+2+l) \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{2(s-j-2)+l+j}^{\left[\begin{smallmatrix} s-j-2 \\ s-j-2 \\ s-j-2+(j+1+l) \end{smallmatrix} \right]} = -9 \cdot 2^{2k+2s+2l-6},$$

(11.18)

$$\sum_{i=0}^{j+1} 2^{4i} \cdot \Delta_k \omega_{2(s-i)+l+j+1}(s-i, 0, l+i) = 3 \cdot 2^{2k+2s+2l+4j-2} - 15 \cdot 2^{2k+2s+2l+4j},$$

(11.19)

$$\sum_{i=0}^{j+1} 2^{4i+2} \cdot \Delta_k \omega_{2(s-i-1)+l+j+2}(s-i-1, 1, l+i) = 3 \cdot 2^{2k+2s+2l+4j-1} - 15 \cdot 2^{2k+2s+2l+4j+1},$$

Combining (11.16), (11.17), (11.18) and (11.19) we obtain

$$(11.20) \quad \Delta_k \Gamma_{2s+l+1+j}^{\left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{2s+l+j}^{\left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right]} = -315 \cdot 2^{2k+2s+2l+4j-2} \quad \text{for } 0 \leq j \leq s-3$$

Proof. proof of (11.16)

Let q be a rational integer between zero and $s-2$. From (8.3) we obtain successively the following equations.

$$\begin{aligned} & \Delta_k \Gamma_{2s+l+1+j}^{\left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{2s+l+j}^{\left[\begin{smallmatrix} s-1 \\ s+l \end{smallmatrix} \right]} = 8 \cdot \left[\Delta_k \Gamma_{2s+l+j}^{\left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{2s+l+j-1}^{\left[\begin{smallmatrix} s-1 \\ s+l-1 \end{smallmatrix} \right]} \right] + \Delta_k \omega_{2s+l+1+j}(s, 0, l) \\ & 4 \cdot \left[\Delta_k \Gamma_{2s+l+j}^{\left[\begin{smallmatrix} s-1 \\ s+l \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{2s+l+j-1}^{\left[\begin{smallmatrix} s-1 \\ s+l \end{smallmatrix} \right]} \right] = 4 \cdot \left[8 \cdot \left[\Delta_k \Gamma_{2s+l+j-1}^{\left[\begin{smallmatrix} s-1 \\ s+l-1 \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{2s+l+j-2}^{\left[\begin{smallmatrix} s-1 \\ s+l-1 \end{smallmatrix} \right]} \right] + \Delta_k \omega_{2s+l+j}(s-1, 1, l) \right] \\ & 4^2 \cdot \left[\Delta_k \Gamma_{2s+l+j-1}^{\left[\begin{smallmatrix} s-1 \\ s+l \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{2s+l+j-2}^{\left[\begin{smallmatrix} s-2 \\ s+l \end{smallmatrix} \right]} \right] = 4^2 \cdot \left[8 \cdot \left[\Delta_k \Gamma_{2s+l+j-2}^{\left[\begin{smallmatrix} s-1 \\ s+l-1 \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{2s+l+j-3}^{\left[\begin{smallmatrix} s-2 \\ s+l-1 \end{smallmatrix} \right]} \right] + \Delta_k \omega_{2s+l+j-1}(s-1, 0, l+1) \right] \\ & \vdots \\ & 4^{2q} \cdot \left[\Delta_k \Gamma_{2s+l+1+j-2q}^{\left[\begin{smallmatrix} s-q \\ s+l \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{2s+l+j-2q}^{\left[\begin{smallmatrix} s-q-1 \\ s+l \end{smallmatrix} \right]} \right] \\ & = 4^{2q} \cdot \left[8 \cdot \left[\Delta_k \Gamma_{2s+l+j-2q}^{\left[\begin{smallmatrix} s-q \\ s+l-1 \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{2s+l+j-1-2q}^{\left[\begin{smallmatrix} s-q-1 \\ s+l-1 \end{smallmatrix} \right]} \right] + \Delta_k \omega_{2s+l+1+j-2q}(s-q, 0, l+q) \right] \\ & 4^{2q+1} \cdot \left[\Delta_k \Gamma_{2s+l+j-2q}^{\left[\begin{smallmatrix} s-q-1 \\ s+l \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{2s+l+j-1-2q}^{\left[\begin{smallmatrix} s-q-1 \\ s+l \end{smallmatrix} \right]} \right] \\ & = 4^{2q+1} \cdot \left[8 \cdot \left[\Delta_k \Gamma_{2s+l+j-1-2q}^{\left[\begin{smallmatrix} s-q-1 \\ s+l-1 \end{smallmatrix} \right]} - 4 \cdot \Delta_k \Gamma_{2s+l+j-2-2q}^{\left[\begin{smallmatrix} s-q-1 \\ s+l-1 \end{smallmatrix} \right]} \right] + \Delta_k \omega_{2s+l+j-2q}(s-q-1, 1, l+q) \right] \end{aligned}$$

By summing the above equations we obtain the following formula holding for $0 \leq q \leq s-2$

(11.21)

$$\begin{aligned} & \Delta_k \Gamma_{2s+l+1+j}^{\left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{2s+l+j}^{\left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right]} = 4^{2q+2} \cdot \Delta_k \Gamma_{2s+l+j-1-2q}^{\left[\begin{smallmatrix} s-q-1 \\ s+l \end{smallmatrix} \right]} - 8 \cdot 4^{2q+2} \cdot \Delta_k \Gamma_{2s+l+j-2-2q}^{\left[\begin{smallmatrix} s-q-1 \\ s+l-1 \end{smallmatrix} \right]} \\ & + \sum_{i=0}^q 4^{2i} \cdot \Delta_k \omega_{2(s-i)+l+j+1}(s-i, 0, l+i) + \sum_{i=0}^q 4^{2i+1} \cdot \Delta_k \omega_{2(s-i-1)+l+j+2}(s-i-1, 1, l+i) \end{aligned}$$

Setting $q = j+1$ for $0 \leq j \leq s-3$ in (11.21) we get the reduction formula (11.16)

Proof of (11.17)

We get from (11.13) with $s \rightarrow s-j-2$, $l \rightarrow j+2+l$, $j \rightarrow l+j+1$:

$$\Delta_k \Gamma_{2(s-j-2)+l+j+1}^{\begin{bmatrix} s-j-2 \\ s-j-2 \\ s-j-2+(j+2+l) \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2(s-j-2)+l+j}^{\begin{bmatrix} s-j-2 \\ s-j-2 \\ s-j-2+(j+1+l) \end{bmatrix}} = -9 \cdot 2^{2k+2(s-j-2)+2(j+2+l)-4} = -9 \cdot 2^{2k+2s+2l-6},$$

observing that $2 \leq l+j+1 \leq j+2+l-1$ and $s-j-2 \geq s-(s-3)-2=1$

Remark that in the case $j = s-3$ we have from Lemma 1.23

$$\Delta_k \Gamma_{1+(l+s-1)}^{\begin{bmatrix} 1 \\ 1+(s+l-1) \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{1+(l+s-2)}^{\begin{bmatrix} 1 \\ 1+(s+l-2) \end{bmatrix}} = -9 \cdot 2^{2k+2s+2l-6}$$

Proof of (11.18)

We obtain from Lemma 7.4 (see (7.51))

$$\begin{aligned} \Delta_k \omega_{2(s-i)+l+j+1}(s-i, 0, l+i) &= 0 \\ \Updownarrow \\ 2(s-i) + l + i + 2 &\leq 2(s-i) + l + j + 1 \leq 3 \cdot (s-i) + l + i - 1 \\ \Updownarrow \\ i &\leq j-1 \text{ and } j \leq s-2 \end{aligned}$$

We then deduce :

$$\begin{aligned} \sum_{i=0}^{j+1} 2^{4i} \cdot \Delta_k \omega_{2(s-i)+l+j+1}(s-i, 0, l+i) &= \sum_{i=j}^{j+1} 2^{4i} \cdot \Delta_k \omega_{2(s-i)+(l+j+1)}(s-i, 0, l+i) = \\ &= 2^{4j} \cdot \Delta_k \omega_{2(s-j)+l+j+1}(s-j, 0, l+j) + 2^{4(j+1)} \cdot \Delta_k \omega_{2(s-j-1)+l+j+1}(s-j-1, 0, l+j+1) \\ &= 2^{4j} \cdot 3 \cdot 2^{2k+2(s-j)+2(l+j)-2} + 2^{4(j+1)} \cdot (-15) \cdot 2^{2k+2(s-j-1)+2(l+j+1)-4} \\ &= 3 \cdot 2^{2k+2s+2l+4j-2} - 15 \cdot 2^{2k+2s+2l+4j} \end{aligned}$$

Proof of (11.19)

Similar to the proof of (11.18).

$$\begin{aligned} \Delta_k \omega_{2(s-i-1)+l+j+2}(s-i-1, 1, l+i) &= 0 \\ \Updownarrow \\ 1 + 2(s-i-1) + l + i + 2 &\leq 2(s-i-1) + l + j + 2 \leq 2 + 3 \cdot (s-i-1) + l + i - 1 \\ \Updownarrow \\ i &\leq j-1 \text{ and } j \leq s-2 \end{aligned}$$

We then deduce :

$$\sum_{i=0}^{j+1} 2^{4i+2} \cdot \Delta_k \omega_{2(s-i-1)+l+j+2}(s-i-1, 1, l+i) = \sum_{i=j}^{j+1} 2^{4i+2} \cdot \Delta_k \omega_{2(s-i-1)+(l+j+2)}(s-i-1, 1, l+i) =$$

$$\begin{aligned}
&= 2^{4j+2} \cdot \Delta_k \omega_{2(s-j-1)+l+j+2}(s-j-1, 1, l+j) + 2^{4(j+1)+2} \cdot \Delta_k \omega_{2(s-j-1)+l+j+2}(s-j-2, 1, l+j+1) \\
&= 2^{4j+2} \cdot 3 \cdot 2^{2k+2(s-j-1)+1+2(l+j)-2} + 2^{4j+6} \cdot (-15) \cdot 2^{2k+2(s-j-2)+1+2(l+j+1)-4} \\
&= 3 \cdot 2^{2k+2s+2l+4j-1} - 15 \cdot 2^{2k+2s+2l+4j+1}
\end{aligned}$$

(11.19) is obtained as indicated in Lemma 11.7 □

Lemma 11.8. *We have for $l \geq 3$ the following reduction formula*

(11.22)

$$\begin{aligned}
&\Delta_k \Gamma_{3s+l-1}^{\left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{3s+l-2}^{\left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right]} = 2^{4s-4} \cdot \Delta_k \Gamma_{s+l+1}^{\left[\begin{smallmatrix} 1 \\ s+l \end{smallmatrix} \right]} - 8 \cdot 2^{4s-4} \cdot \Delta_k \Gamma_{s+l}^{\left[\begin{smallmatrix} 1 \\ s+l-1 \end{smallmatrix} \right]} \\
&+ \sum_{i=0}^{s-2} 2^{4i} \cdot \Delta_k \omega_{2(s-i)+l+s-1}(s-i, 0, l+i) + \sum_{i=0}^{s-2} 2^{4i+2} \cdot \Delta_k \omega_{2(s-i-1)+l+s}(s-i-1, 1, l+i)
\end{aligned}$$

where

$$(11.23) \quad \Delta_k \Gamma_{s+l+1}^{\left[\begin{smallmatrix} 1 \\ s+l \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{s+l}^{\left[\begin{smallmatrix} 1 \\ s+l-1 \end{smallmatrix} \right]} = -81 \cdot 2^{2k+2s+2l-4},$$

$$(11.24) \quad \sum_{i=0}^{s-2} 2^{4i} \cdot \Delta_k \omega_{2(s-i)+l+s-1}(s-i, 0, l+i) = 3 \cdot 2^{2k+2l+6s-10},$$

$$(11.25) \quad \sum_{i=0}^{s-2} 2^{4i+2} \cdot \Delta_k \omega_{2(s-i-1)+l+s}(s-i-1, 1, l+i) = 3 \cdot 2^{2k+2s+2l+6s-9},$$

Combining (11.22), (11.23), (11.24) and (11.25) we obtain

$$(11.26) \quad \Delta_k \Gamma_{3s+l-1}^{\left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{3s+l-2}^{\left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right]} = -315 \cdot 2^{2k+2l+6s-10}$$

Proof.

Proof of (11.22)

Setting $q = j = s - 2$ in (11.21) we get the reduction formula (11.22)

Proof of (11.23)

From Lemma 1.23 we obtain :

$$\begin{aligned}
&\Delta_k \Gamma_{s+l+1}^{\left[\begin{smallmatrix} 1 \\ 1+s+l-1 \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{s+l}^{\left[\begin{smallmatrix} 1 \\ 1+s+l-2 \end{smallmatrix} \right]} \\
&= 81 \cdot (2^{2k+2s+2l-4} - 2^{k+3s+3l-5}) - 8 \cdot (81 \cdot (2^{2k+2s+2l-6} - 2^{k+3s+3l-8})) = -81 \cdot 2^{2k+2s+2l-4}.
\end{aligned}$$

Proof of (11.24) We obtain from Lemma 7.4 (see (7.37))

$$\begin{aligned}
&\Delta_k \omega_{2(s-i)+l+s-1}(s-i, 0, l+i) = 0 \\
&\Updownarrow \\
&2(s-i) + l + i + 2 \leq 2(s-i) + l + s - 1 \leq 3 \cdot (s-i) + l + i - 1 \\
&\Updownarrow \\
&i \leq s - 3
\end{aligned}$$

We then deduce :

$$\begin{aligned} \sum_{i=0}^{s-2} 2^{4i} \cdot \Delta_k \omega_{2(s-i)+l+s-1}(s-i, 0, l+i) &= \sum_{i=s-2}^{s-2} 2^{4i} \cdot \Delta_k \omega_{2(s-i)+(l+s-1)}(s-i, 0, l+i) = \\ &= 2^{4(s-2)} \cdot \Delta_k \omega_{l+s+3}(2, 0, l+s-2) = 2^{4(s-2)} \cdot 3 \cdot 2^{2k+2(l+s-2)+2} = 3 \cdot 2^{2k+2l+6s-10} \end{aligned}$$

Proof of (11.25) We obtain from Lemmas 1.22, 1.23

$$\begin{aligned} \Delta_k \omega_{2(s-i-1)+l+s}(s-i-1, 1, l+i) &= 0 \\ \Downarrow \\ 1 + 2(s-i-1) + l + i + 2 &\leq 2(s-i-1) + l + s \leq 2 + 3 \cdot (s-i-1) + l + i - 1 \\ \Downarrow \\ i &\leq s-3 \end{aligned}$$

We then deduce :

$$\begin{aligned} \sum_{i=0}^{s-2} 2^{4i+2} \cdot \Delta_k \omega_{2(s-i-1)+l+s}(s-i-1, 1, l+i) &= \sum_{i=s-2}^{s-2} 2^{4i+2} \cdot \Delta_k \omega_{2(s-i-1)+l+s}(s-i-1, 1, l+i) = \\ &= 2^{4s-6} \cdot \Delta_k \omega_{l+s+2}(1, 1, l+s-2) \\ &= 2^{4s-6} \cdot \left(\Delta_k \Gamma_{l+s+2}^{\left[\frac{1}{2} \right]} - 4 \cdot \Delta_k \Gamma_{l+s+1}^{\left[\frac{1}{s+l} \right]} - 8 \cdot \Delta_k \Gamma_{l+s+1}^{\left[\frac{1}{s+l-1} \right]} + 32 \cdot \Delta_k \Gamma_{l+s}^{\left[\frac{1}{s+l-1} \right]} \right) \\ &= 2^{4s-6} \cdot \left(\left(\Delta_k \Gamma_{l+s+2}^{\left[\frac{1}{1+1+s+l-2} \right]} - 8 \cdot \Delta_k \Gamma_{l+s+1}^{\left[\frac{1}{1+1+s+l-3} \right]} \right) - 4 \cdot \left(\Delta_k \Gamma_{l+s+1}^{\left[\frac{1}{1+s+l-1} \right]} - 8 \cdot \Delta_k \Gamma_{l+s}^{\left[\frac{1}{1+s+l-2} \right]} \right) \right) \\ &= 2^{4s-6} \cdot \left(-159 \cdot 2^{2k+2s+2l-3} - 4 \cdot (-81) \cdot 2^{2k+2l+2s-4} \right) = 2^{4s-6} \cdot 3 \cdot 2^{2k+2l+2s-3} \\ &= 3 \cdot 2^{2k+2l+6s-9}, \end{aligned}$$

or otherwise using (5.2) and (7.37)

$$\Delta_k \omega_{l+s+2}(1, 1, l+s-2) = \frac{1}{2} \cdot \Delta_k \omega_{l+s+3}(2, 0, l+s-2) = 3 \cdot 2^{2k+2l+2s-3}$$

□

Lemma 11.9. *We have for $l \geq 3$ the following reduction formula*

(11.27)

$$\begin{aligned} \Delta_k \Gamma_{3s+l}^{\left[\frac{s}{s+l} \right]} - 8 \cdot \Delta_k \Gamma_{3s+l-1}^{\left[\frac{s}{s+l-1} \right]} &= 2^{4s-4} \cdot \Delta_k \Gamma_{s+l+2}^{\left[\frac{1}{s+l} \right]} - 8 \cdot 2^{4s-4} \cdot \Delta_k \Gamma_{s+l+1}^{\left[\frac{1}{s+l-1} \right]} \\ &+ \sum_{i=0}^{s-2} 2^{4i} \cdot \Delta_k \omega_{2(s-i)+l+s}(s-i, 0, l+i) + \sum_{i=0}^{s-2} 2^{4i+2} \cdot \Delta_k \omega_{2(s-i-1)+l+s+1}(s-i-1, 1, l+i) \end{aligned}$$

where

$$(11.28) \quad \Delta_k \Gamma_{s+l+2}^{\left[\frac{1}{1+s+l-1} \right]} - 8 \cdot \Delta_k \Gamma_{s+l+1}^{\left[\frac{1}{1+s+l-2} \right]} = -21 \cdot 2^{3k+s+l-1} + 21 \cdot 2^{2k+2s+2l-2},$$

$$(11.29) \quad \sum_{i=0}^{s-2} 2^{4i} \cdot \Delta_k \omega_{2(s-i)+l+s}(s-i, 0, l+i) = 7 \cdot 2^{3k+5s+l-5} - 7 \cdot 2^{3k+3s+l-3},$$

$$(11.30) \quad \sum_{i=0}^{s-2} 2^{4i+2} \cdot \Delta_k \omega_{2(s-i-1)+l+s+1}(s-i-1, 1, l+i) = 7 \cdot 2^{3k+5s+l-6} - 7 \cdot 2^{3k+3s+l-4},$$

Combining (11.27), (11.28), (11.29) and (11.30) we obtain

$$(11.31) \quad \Delta_k \Gamma_{3s+l} \left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_{3s+l-1} \left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right] = -21 \cdot 2^{3k+3s+l-3} + 21 \cdot 2^{2k+6s+2l-6}$$

Proof. Proof of (11.27)

Setting $q = s - 2$, $j = s - 1$ in (11.21) we get the reduction formula (11.27)

Proof of (11.28) From Lemma 1.23 we obtain :

$$\begin{aligned} & \Delta_k \Gamma_{s+l+2} \left[\begin{smallmatrix} 1 \\ 1+s+l-1 \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_{s+l+1} \left[\begin{smallmatrix} 1 \\ 1+s+l-2 \end{smallmatrix} \right] \\ &= (7 \cdot 2^{3k+s+l-1} - 21 \cdot 2^{2k+2s+2l-2} + 7 \cdot 2^{k+3s+3l-2}) \\ & \quad - 8 \cdot (7 \cdot 2^{3k+s+l-2} - 21 \cdot 2^{2k+2s+2l-4} + 7 \cdot 2^{k+3s+3l-5}) \\ &= -21 \cdot 2^{3k+s+l-1} + 21 \cdot 2^{2k+2s+2l-2} \end{aligned}$$

Proof of (11.29) We obtain from Lemma 7.4 (see (7.37), (7.46) and (7.51))

$$\Delta_k \omega_{2(s-i)+l+s}(s-i, 0, l+i) = 21 \cdot 2^{3k+3s-2i+l-3} \quad \text{for } 0 \leq i \leq s-2$$

We then deduce :

$$\sum_{i=0}^{s-2} 2^{4i} \cdot \Delta_k \omega_{2(s-i)+l+s}(s-i, 0, l+i) = \sum_{i=0}^{s-2} 2^{4i} \cdot 21 \cdot 2^{3k+3s-2i+l-3} = 7 \cdot 2^{3k+5s+l-5} - 7 \cdot 2^{3k+3s+l-3}$$

Proof of (11.30) We obtain from Lemma 7.4 (see (7.38), (7.47) and (7.51))

$$\begin{aligned} & \Delta_k \omega_{2(s-i-1)+l+s+1}(s-i-1, 1, l+i) = 21 \cdot 2^{3k+3s-2i+l-4} \quad \text{for } 0 \leq i \leq s-3 \\ & \Delta_k \omega_{l+s+3}(1, 1, l+s-2) = \frac{1}{2} \cdot \Delta_k \omega_{l+s+4}(2, 0, l+s-2) \\ &= \frac{1}{2} \cdot 21 \cdot 2^{3k+l+s+1} = 21 \cdot 2^{3k+l+s} \quad \text{for } i = s-2 \end{aligned}$$

We then deduce :

$$\begin{aligned} & \sum_{i=0}^{s-2} 2^{4i+2} \cdot \Delta_k \omega_{2(s-i-1)+l+s+1}(s-i-1, 1, l+i) = \sum_{i=0}^{s-2} 2^{4i+2} \cdot 21 \cdot 2^{3k+3s-2i+l-4} \\ &= 7 \cdot 2^{3k+5s+l-6} - 7 \cdot 2^{3k+3s+l-4} \end{aligned}$$

□

$$11.9. \text{ Computation of } \Delta_k \Gamma_{2s+m+j} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_{2s+m+j-1} \left[\begin{smallmatrix} s \\ s+m-1 \end{smallmatrix} \right] \quad \text{for } 0 \leq j \leq m.$$

Lemma 11.10. *We have for $m \geq 3$ the following reduction formula holding for $0 \leq j \leq s+m$.*

$$(11.32) \quad \begin{aligned} & \Delta_k \Gamma_{2s+m+j} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_{2s+m+j-1} \left[\begin{smallmatrix} s \\ s+m-1 \end{smallmatrix} \right] \\ &= 2^{2m} \cdot \left[\Delta_k \Gamma_{2s+j} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] - 8 \cdot \Delta_k \Gamma_{2s+j-1} \left[\begin{smallmatrix} s \\ s+m-1 \end{smallmatrix} \right] \right] + \sum_{i=0}^{m-1} 2^{2i} \cdot \Delta_k \omega_{2s+m+j-i}(s, m-i, i), \end{aligned}$$

where

$$(11.33) \quad \Delta_k \Gamma_{2s+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+j-1}^{\begin{bmatrix} s \\ s+m-1 \\ s+m \end{bmatrix}} = \begin{cases} 3 \cdot 2^{2k+2s-2} - 33 \cdot 2^{k+4s-6} & \text{if } j = 0, \\ -15 \cdot 2^{2k+2s-2} + 3 \cdot 2^{k+4s-3} & \text{if } j = 1, \\ -9 \cdot 2^{2k+2s+2j-4} & \text{if } 2 \leq j \leq m-1, \\ -81 \cdot 2^{2k+2s+2m-4} & \text{if } j = m, \end{cases}$$

(11.34)

$$\sum_{i=0}^{m-1} 2^{2i} \cdot \Delta_k \omega_{2s+m+j-i}(s, m-i, i) = \begin{cases} -3 \cdot 2^{2k+2s+m-2} - 3 \cdot 2^{2k+2s+m-2} & \text{if } j = 0, \\ 15 \cdot 2^{2k+2s+2m-2} - 69 \cdot 2^{2k+2s+m-2} & \text{if } j = 1, \\ -63 \cdot 2^{2k+2s+m+3j-5} + 9 \cdot 2^{2k+2s+2m+2j-4} & \text{if } 2 \leq j \leq m-1, \\ 3 \cdot 2^{2k+2s+4m-5} & \text{if } j = m. \end{cases}$$

Combining (11.32), (11.33) and (11.34) we obtain

(11.35)

$$\Delta_k \Gamma_{2s+m+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+m+j-1}^{\begin{bmatrix} s \\ s+m-1 \\ s+m \end{bmatrix}} = \begin{cases} -3 \cdot 2^{2k+2s+m-2} - 33 \cdot 2^{k+4s+2m-6} & \text{if } j = 0 \\ -69 \cdot 2^{2k+2s+m-2} + 3 \cdot 2^{k+4s+2m-3} & \text{if } j = 1 \\ -63 \cdot 2^{2k+2s+m+3j-5} & \text{if } 2 \leq j \leq m-1 \\ -159 \cdot 2^{2k+2s+4m-5} & \text{if } j = m \end{cases}$$

Proof. Proof of (11.32)

Let q be a rational integer between zero and $m-1$. From (8.3) we obtain successively the following equations.

$$\begin{aligned} \Delta_k \Gamma_{2s+m+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{2s+m+j-1}^{\begin{bmatrix} s \\ s+m-1 \\ s+m \end{bmatrix}} &= 8 \cdot \left[\Delta_k \Gamma_{2s+m+j-1}^{\begin{bmatrix} s \\ s+m-1 \\ s+m \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{2s+m+j-2}^{\begin{bmatrix} s \\ s+m-1 \\ s+m-1 \end{bmatrix}} \right] + \Delta_k \omega_{2s+m+j}(s, m, 0) \\ 4 \cdot \left[\Delta_k \Gamma_{2s+m+j-1}^{\begin{bmatrix} s \\ s+m-1 \\ s+m \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{2s+m+j-2}^{\begin{bmatrix} s \\ s+m-2 \\ s+m \end{bmatrix}} \right] &= 4 \cdot \left[8 \cdot \left[\Delta_k \Gamma_{2s+m+j-2}^{\begin{bmatrix} s \\ s+m-1 \\ s+m-1 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{2s+m+j-3}^{\begin{bmatrix} s \\ s+m-2 \\ s+m-1 \end{bmatrix}} \right] + \Delta_k \omega_{2s+m+j-1}(s, m-1, 1) \right] \\ 4^2 \cdot \left[\Delta_k \Gamma_{2s+m+j-2}^{\begin{bmatrix} s \\ s+m-2 \\ s+m \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{2s+m+j-3}^{\begin{bmatrix} s \\ s+m-3 \\ s+m \end{bmatrix}} \right] &= 4^2 \cdot \left[8 \cdot \left[\Delta_k \Gamma_{2s+m+j-3}^{\begin{bmatrix} s \\ s+m-2 \\ s+m-1 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{2s+m+j-4}^{\begin{bmatrix} s \\ s+m-3 \\ s+m-1 \end{bmatrix}} \right] + \Delta_k \omega_{2s+m+j-2}(s, m-2, 2) \right] \\ &\vdots \\ 4^q \cdot \left[\Delta_k \Gamma_{2s+m+j-q}^{\begin{bmatrix} s \\ s+m-q \\ s+m \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{2s+m+j-(q+1)}^{\begin{bmatrix} s \\ s+m-(q+1) \\ s+m \end{bmatrix}} \right] \\ &= 4^q \cdot \left[8 \cdot \left[\Delta_k \Gamma_{2s+m+j-(q+1)}^{\begin{bmatrix} s \\ s+m-q \\ s+m-1 \end{bmatrix}} - 4 \cdot \Delta_k \Gamma_{2s+m+j-(q+2)}^{\begin{bmatrix} s \\ s+m-(q+1) \\ s+m-1 \end{bmatrix}} \right] + \Delta_k \omega_{2s+m+j-q}(s, m-q, q) \right] \end{aligned}$$

By summing the above equations we obtain after reduction the following formula

$$\begin{aligned} \Delta_k \Gamma_{2s+m+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+m+j-1}^{\begin{bmatrix} s \\ s+m-1 \\ s+m \end{bmatrix}} \\ = 4^{q+1} \cdot \Delta_k \Gamma_{2s+m+j-(q+1)}^{\begin{bmatrix} s+m-(q+1) \\ s+m \end{bmatrix}} - 8 \cdot 4^{q+1} \cdot \Delta_k \Gamma_{2s+m+j-(q+2)}^{\begin{bmatrix} s+m-(q+1) \\ s+m-1 \end{bmatrix}} + \sum_{i=0}^q 4^i \cdot \Delta_k \omega_{2s+m+j-i}(s, m-i, i) \end{aligned}$$

Setting $q = m-1$ in the above formula we get (11.32)

Proof of (11.33)

(11.33) is obtained by (11.13) with $l \mapsto m$.

Proof of (11.34)

The proof is based on results in Lemma 7.4

The case $j = 0$, we have

$$\Delta_k \omega_{2s+(m-i)}(s, m-i, i) = -3 \cdot 2^{2k+2s+(m-i)-2} \quad \text{if } 1 \leq i \leq s-3,$$

$$\Delta_k \omega_{2s+m}(s, m, 0) = -9 \cdot 2^{2k+2s+m-2} \quad \text{if } i = 0.$$

It follows that

$$\begin{aligned} \sum_{i=0}^{m-1} 2^{2i} \cdot \Delta_k \omega_{2s+(m-i)}(s, m-i, i) &= (-9 \cdot 2^{2k+2s+m-2}) + \sum_{i=1}^{m-1} 2^{2i} \cdot (-3 \cdot 2^{2k+2s+(m-i)-2}) \\ &= -3 \cdot 2^{2k+2s+m-2} - 3 \cdot 2^{2k+2s+m-2} \end{aligned}$$

The case $j = 1$, we obtain similarly

$$\Delta_k \omega_{2s+(m-i)+1}(s, m-i, i) = 15 \cdot 2^{2k+2s+(m-i)-2} \quad \text{if } 2 \leq i \leq m-1,$$

$$\Delta_k \omega_{2s+(m-1)+1}(s, m-1, 1) = -9 \cdot 2^{2k+2s+(m-1)-2} \quad \text{if } i = 1,$$

$$\Delta_k \omega_{2s+m+1}(s, m, 0) = 9 \cdot 2^{2k+2s+m-2} \quad \text{if } i = 0,$$

It follows that

$$\begin{aligned} \sum_{i=0}^{m-1} 2^{2i} \cdot \Delta_k \omega_{2s+(m-i)+1}(s, m-i, i) &= 9 \cdot 2^{2k+2s+m-2} - 9 \cdot 2^{2k+2s+(m-1)-2} + \sum_{i=2}^{m-1} 2^{2i} \cdot (15 \cdot 2^{2k+2s+(m-i)-2}) \\ &= 15 \cdot 2^{2k+2s+m-2} - 69 \cdot 2^{2k+2s+m-2} \end{aligned}$$

The case $2 \leq j \leq m-1$, similarly

$$\Delta_k \omega_{2s+(m-i)+j}(s, m-i, i) = 0 \quad \text{if } 0 \leq i \leq j-2$$

$$\Delta_k \omega_{2s+(m-i)+j}(s, m-i, i) = 3 \cdot 2^{2k+2s+m+j-3} \quad \text{if } i = j-1$$

$$\Delta_k \omega_{2s+(m-i)+j}(s, m-i, i) = -15 \cdot 2^{2k+2s+m+j-4} \quad \text{if } i = j$$

$$\Delta_k \omega_{2s+(m-i)+j}(s, m-i, i) = 9 \cdot 2^{2k+2s+m+2j-4-i} \quad \text{if } i \geq j+1$$

We then deduce

$$\begin{aligned} \sum_{i=0}^{m-1} 2^{2i} \cdot \Delta_k \omega_{2s+(m-i)+j}(s, m-i, i) &= \sum_{i=j-1}^{m-1} 2^{2i} \cdot \Delta_k \omega_{2s+(m-i)+j}(s, m-i, i) \\ &= (3 \cdot 2^{2k+2s+m+j-3}) \cdot 2^{2(j-1)} + (-15 \cdot 2^{2k+2s+m+j-4}) \cdot 2^{2j} + \sum_{i=j+1}^{m-1} 2^{2i} \cdot 9 \cdot 2^{2k+2s+m+2j-4-i} \\ &= -63 \cdot 2^{2k+2s+m+3j-5} + 9 \cdot 2^{2k+2s+m+2j-4} \end{aligned}$$

The case $j = m$, similarly

$$\Delta_k \omega_{2s+(m-i)+m}(s, m-i, i) = 0 \quad \text{if } 0 \leq i \leq m-2$$

$$\Delta_k \omega_{2s+1+m}(s, 1, m-1) = 3 \cdot 2^{2k+2s+2m-3} \quad \text{if } i = m-1$$

We then deduce

$$\begin{aligned} \sum_{i=0}^{m-1} 2^{2i} \cdot \Delta_k \omega_{2s+(m-i)+m}(s, m-i, i) &= \sum_{i=m-1}^{m-1} 2^{2i} \cdot \Delta_k \omega_{2s+(m-i)+m}(s, m-i, i) \\ &= 3 \cdot 2^{2k+2s+4m-5} \end{aligned}$$

□

$$11.10. \text{ Computation of } \Delta_k \Gamma_{2s+2m+j+1}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+2m+j}^{\begin{bmatrix} s \\ s+m-1 \\ s+m \end{bmatrix}} \quad \text{for } 0 \leq j \leq s-1.$$

Lemma 11.11. *We have for $m \geq 3$ the following reduction formula holding for $0 \leq j \leq s-1$.*

(11.36)

$$\begin{aligned} &\Delta_k \Gamma_{2s+2m+j+1}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+2m+j}^{\begin{bmatrix} s \\ s+m-1 \\ s+m \end{bmatrix}} \\ &= 2^{2m} \cdot \left[\Delta_k \Gamma_{2s+m+1+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+m+j}^{\begin{bmatrix} s \\ s+m-1 \\ s+m \end{bmatrix}} \right] + \sum_{i=0}^{m-1} 2^{2i} \cdot \Delta_k \omega_{2s+2m+j+1-i}(s, m-i, i), \end{aligned}$$

where

(11.37)

$$\Delta_k \Gamma_{2s+m+1+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+m+j}^{\begin{bmatrix} s \\ s+m-1 \\ s+m \end{bmatrix}} = \begin{cases} -315 \cdot 2^{2k+2s+2m+4j-2} & \text{if } 0 \leq j \leq s-2, \\ -21 \cdot 2^{3k+3s+m-3} + 21 \cdot 2^{2k+6s+2m-6} & \text{if } j = s-1, \end{cases}$$

(11.38)

$$\sum_{i=0}^{m-1} 2^{2i} \cdot \Delta_k \omega_{2s+2m+j+1-i}(s, m-i, i) = \begin{cases} 0 & \text{if } 2 \leq j \leq s-2, \\ -21 \cdot 2^{3k+3s+2m-3} + 21 \cdot 2^{2k+3s+3m-3} & \text{if } j = s-1. \end{cases}$$

Combining (11.36), (11.37) and (11.38) we obtain

(11.39)

$$\Delta_k \Gamma_{2s+2m+1+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+2m+j}^{\begin{bmatrix} s \\ s+m-1 \\ s+m \end{bmatrix}} = \begin{cases} -315 \cdot 2^{2k+2s+4m+4j-2} & \text{if } 0 \leq j \leq s-2 \\ -21 \cdot 2^{3k+3s+2m-3} + 21 \cdot 2^{2k+6s+4m-6} & \text{if } j = s-1 \end{cases}$$

Proof. Proof of (11.36)

(11.36) follows from (11.32) with $j \longrightarrow m+j+1$.

Proof of (11.37)

(11.37) is obtained by (11.20), (11.26) and (11.31) with $l \longrightarrow m$.

Proof of (11.38)

The case $0 \leq j \leq s-2$

From (7.51)

$$\Delta_k \omega_{2s+2m+j+1-i}(s, m-i, i) = 0$$

$$\begin{aligned}
& \Updownarrow \\
& m - i + 2s + i + 2 \leq 2s + 2m + j + 1 - i \leq 2(m - i) + 3s + i - 1 \\
& \Updownarrow \\
& 3 \leq i \leq \inf(m - 1, m + j - 1) = m - 1 \\
& \text{From (7.50)} \\
& \Delta_k \omega_{2s+2m+j-1}(s, m - 2, 2) = 0 \\
& \Updownarrow \\
& m - 2 + 2s + 4 \leq 2s + 2m + j - 1 \leq 2(m - 2) + 3s + 1 \\
& \Updownarrow \\
& \sup(0, -m + 3) \leq j \leq s - 2 \quad \text{recall that } m \geq 3 \\
& \text{From (7.49)} \\
& \Delta_k \omega_{2s+2m+j}(s, m - 1, 1) = 0 \\
& \Updownarrow \\
& m - 1 + 2s + 3 \leq 2s + 2m + j \leq 2(m - 1) + 3s \\
& \Updownarrow \\
& \sup(0, -m + 2) \leq j \leq s - 2 \quad \text{recall that } m \geq 3 \\
& \text{From (7.48)} \\
& \Delta_k \omega_{2s+2m+j+1}(s, m, 0) = 0 \\
& \Updownarrow \\
& m + 2s + 2 \leq 2s + 2m + j + 1 \leq 2m + 3s - 1 \\
& \Updownarrow \\
& \sup(0, -m + 1) \leq j \leq s - 2 \quad \text{recall that } m \geq 3 \\
& \text{We then deduce} \\
& \sum_{i=0}^{m-1} 2^{2i} \cdot \Delta_k \omega_{2s+2m+j+1-i}(s, m - i, i) = 0 \quad \text{if } 0 \leq j \leq s - 2
\end{aligned}$$

The case $j = s - 1$

Similar to the proof of (11.38) in the case $0 \leq j \leq s - 2$

$$\Delta_k \omega_{3s+2m-i}(s, m - i, i) = \Delta_k \omega_{3s+2(m-i)+i}(s, m - i, i) = 21 \cdot 2^{3k+3s+2m-i-3} \quad \text{if } 0 \leq i \leq m - 1$$

We then deduce

$$\begin{aligned}
\sum_{i=0}^{m-1} 2^{2i} \cdot \Delta_k \omega_{3s+2m-i}(s, m - i, i) &= \sum_{i=0}^{m-1} 2^{2i} \cdot 21 \cdot 2^{3k+3s+2m-i-3} \\
&= 21 \cdot 2^{2k+3s+3m-3} - 21 \cdot 2^{3k+3s+2m-3}
\end{aligned}$$

□

11.11. Formulas summary.

Lemma 11.12. *We have the following formulas :*

$$(11.40) \quad \Delta_k \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+l \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{s+j-1}^{\left[\begin{smallmatrix} s \\ s+l-1 \end{smallmatrix} \right]} = \begin{cases} 3 \cdot 2^{k+s-1} & \text{if } j = 0, l \geq 0. \\ 15 \cdot 2^{k+s-1} & \text{if } j = 1, l \geq 0. \\ 63 \cdot 2^{k+s+3j-6} & \text{if } 2 \leq j \leq s - 1, l \geq 0. \end{cases}$$

We have for $l \geq 3$:

$$(11.41) \quad \Delta_k \Gamma_{2s+j}^{\begin{bmatrix} s \\ s+l \\ s+j \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+j-1}^{\begin{bmatrix} s \\ s+l-1 \\ s+j-1 \end{bmatrix}} = \begin{cases} 3 \cdot 2^{2k+2s-2} - 33 \cdot 2^{k+4s-6} & \text{if } j = 0 \\ -15 \cdot 2^{2k+2s-2} + 3 \cdot 2^{k+4s-3} & \text{if } j = 1 \\ -9 \cdot 2^{2k+2s+2j-4} & \text{if } 2 \leq j \leq l-1 \\ -81 \cdot 2^{2k+2s+2j-4} & \text{if } j = l \end{cases}$$

$$(11.42) \quad \Delta_k \Gamma_{2s+l+1+j}^{\begin{bmatrix} s \\ s+l \\ s+j \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+l+j}^{\begin{bmatrix} s \\ s+l-1 \\ s+j \end{bmatrix}} = \begin{cases} -315 \cdot 2^{2k+2s+2l+4j-2} & \text{if } 0 \leq j \leq s-2 \\ -21 \cdot 2^{3k+3s+l-3} + 21 \cdot 2^{2k+6s+2l-6} & \text{if } j = s-1 \end{cases}$$

$$(11.43) \quad \Delta_k \Gamma_{s+j}^{\begin{bmatrix} s+m \\ s+m+l \\ s+j \end{bmatrix}} = \begin{cases} 2^{k+s-1} & \text{if } j = 0 \\ \gamma_j \cdot 2^{k+s-1} = (21 \cdot 2^{3j-4} - 3 \cdot 2^{2j-3}) \cdot 2^{k+s-1} & \text{if } 1 \leq j \leq m-1, l \geq 0 \\ 21 \cdot 2^{k+s+3m-5} + 13 \cdot 2^{k+s+2m-4} & \text{if } j = m \geq 1, l \geq 1 \\ 21 \cdot 2^{k+s+3m-5} + 45 \cdot 2^{k+s+2m-4} & \text{if } j = m \geq 1, l = 0 \end{cases}$$

$$(11.44) \quad \Delta_k \Gamma_{s+m+1}^{\begin{bmatrix} s \\ s+m \\ s+m+1 \end{bmatrix}} = \begin{cases} 147 \cdot 2^{k+s-1} & \text{if } m = 0 \\ 315 \cdot 2^{k+s} & \text{if } m = 1 \\ 21 \cdot 2^{k+s+3m-2} + 273 \cdot 2^{k+s+2m-2} & \text{if } m \geq 2 \text{ (holds for } m \geq 0) \end{cases}$$

$$(11.45) \quad \Delta_k \Gamma_{s+m+j}^{\begin{bmatrix} s+m \\ s+m+l \\ s+m+j \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{s+m+j-1}^{\begin{bmatrix} s+m \\ s+m+l-1 \\ s+m+j-1 \end{bmatrix}} = \begin{cases} 9 \cdot 2^{k+s-1} & \text{if } j = 0, m = 1, l \geq 0 \\ 19 \cdot 2^{k+s+2m-4} & \text{if } j = 0, m \geq 2, l \geq 0 \\ 31 \cdot 2^{k+s+2m-2} & \text{if } j = 1, m \geq 1, l \geq 0 \\ 63 \cdot 2^{k+s+2m+3j-6} & \text{if } 2 \leq j \leq s-1, m \geq 0, l \geq 0. \end{cases}$$

We have for $m \geq 3$:

$$(11.46) \quad \Delta_k \Gamma_{2s+m+j}^{\begin{bmatrix} s+m \\ s+m \\ s+m+j \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+m+j-1}^{\begin{bmatrix} s+m \\ s+m-1 \\ s+m+j-1 \end{bmatrix}} = \begin{cases} -3 \cdot 2^{2k+2s+m-2} - 33 \cdot 2^{k+4s+2m-6} & \text{if } j = 0 \\ -69 \cdot 2^{2k+2s+m-2} + 3 \cdot 2^{k+4s+2m-3} & \text{if } j = 1 \\ -63 \cdot 2^{2k+2s+m+3j-5} & \text{if } 2 \leq j \leq m-1 \\ -159 \cdot 2^{2k+2s+4m-5} & \text{if } j = m \end{cases}$$

$$(11.47) \quad \Delta_k \Gamma_{2s+2m+1+j}^{\begin{bmatrix} s+m \\ s+m \\ s+m+j \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+2m+j}^{\begin{bmatrix} s+m \\ s+m-1 \\ s+m+j \end{bmatrix}} = \begin{cases} -315 \cdot 2^{2k+2s+4m+4j-2} & \text{if } 0 \leq j \leq s-2 \\ -21 \cdot 2^{3k+3s+2m-3} + 21 \cdot 2^{2k+6s+4m-6} & \text{if } j = s-1 \end{cases}$$

Proof. We need only to summarize the principal formulas obtained in the subsections 11.3, 11.4, ..., 11.10 $\square$

11.12. Computation of $\Delta_k \Gamma_{s+m+j}^{\begin{bmatrix} s \\ s+m \\ s+m+j \end{bmatrix}}$ for $m \geq 1, 1 \leq j \leq s-1$.

Lemma 11.13. *We have the following reduction formula holding for $m \geq 1, 2 \leq j \leq s-1$:*

$$(11.48) \quad \Delta_k \Gamma_{s+m+j}^{\begin{bmatrix} s \\ s+m \\ s+m+j \end{bmatrix}} = 2^6 \cdot \Delta_k \Gamma_{s+(m-1)+(j-1)}^{\begin{bmatrix} s \\ s+m-1 \\ s+(m-1)+(j-1) \end{bmatrix}} + 189 \cdot 2^{k+s+2m+3j-6}$$

By induction on j we deduce from (11.48)

$$(H_j) \quad \Delta_k \Gamma_{s+m+j}^{\begin{bmatrix} s \\ s+m \\ s+m+j \end{bmatrix}} = 21 \cdot [2^{k+s+3m-2} \cdot 2^{3(j-1)} + a_j \cdot 2^{k+s+2m-3} \cdot 2^{3(j-1)}] \quad \text{where} \quad a_j = 35 \cdot 2^{j-1} - 9$$

$\Updownarrow$

$$\begin{aligned}
& \Delta_k \Gamma_{s+m+j} \begin{bmatrix} s \\ s+m \\ s+m+j \end{bmatrix} \\
(11.49) \quad & = 21 \cdot [2^{k+s+3m+3j-5} + 35 \cdot 2^{k+s+2m+4j-7} - 9 \cdot 2^{k+s+2m+3j-6}] \quad \text{for } m \geq 1, 1 \leq j \leq s-1
\end{aligned}$$

Proof. **proof of** (11.48)

We have from (7.48) and (8.3), (11.45) with $l = 0$

$$\begin{aligned}
& \Delta_k \Gamma_{s+m+j} \begin{bmatrix} s \\ s+m \\ s+m+j \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{s+m+j-1} \begin{bmatrix} s \\ s+m-1 \\ s+m+j-1 \end{bmatrix} = 63 \cdot 2^{k+s+2m+3j-6} \\
& \Updownarrow \\
& 4 \cdot \Delta_k \Gamma_{s+m+j-1} \begin{bmatrix} s \\ s+m-1 \\ s+m+j-1 \end{bmatrix} - 32 \cdot \Delta_k \Gamma_{s+m+j-2} \begin{bmatrix} s \\ s+m-1 \\ s+m+j-2 \end{bmatrix} + \Delta_k \omega_{s+m+j}(s, m, 0) = 63 \cdot 2^{k+s+2m+3j-6} \\
& \Updownarrow \\
& \Delta_k \Gamma_{s+m+j-1} \begin{bmatrix} s \\ s+m-1 \\ s+m+j-1 \end{bmatrix} - 8 \cdot \Delta_k \Gamma_{s+m+j-2} \begin{bmatrix} s \\ s+m-1 \\ s+m+j-2 \end{bmatrix} = 63 \cdot 2^{k+s+2m+3j-8}
\end{aligned}$$

We then deduce

$$\begin{aligned}
& \Delta_k \Gamma_{s+m+j} \begin{bmatrix} s \\ s+m \\ s+m+j \end{bmatrix} = 8 \cdot \left[8 \cdot \Delta_k \Gamma_{s+m+j-2} \begin{bmatrix} s \\ s+m-1 \\ s+m+j-2 \end{bmatrix} + 63 \cdot 2^{k+s+2m+3j-8} \right] + 63 \cdot 2^{k+s+2m+3j-6} \\
& = 2^6 \cdot \Delta_k \Gamma_{s+(m-1)+(j-1)} \begin{bmatrix} s \\ s+(m-1) \\ s+(m-1)+(j-1) \end{bmatrix} + 189 \cdot 2^{k+s+2m+3j-6}
\end{aligned}$$

proof of (11.49)

Proof by induction on j

We obtain from (11.44) : $\Delta_k \Gamma_{s+m+1} \begin{bmatrix} s \\ s+m \\ s+m+1 \end{bmatrix} = 21 \cdot [2^{k+s+3m-2} + 26 \cdot 2^{k+s+2m-3}]$ and so (H_j) holds for $j=1$

$$\begin{aligned}
& \text{We get by (11.48) with } j=2 \text{ using } (H_1) : \Delta_k \Gamma_{s+m+2} \begin{bmatrix} s \\ s+m \\ s+m+2 \end{bmatrix} = 2^6 \cdot \Delta_k \Gamma_{s+(m-1)+1} \begin{bmatrix} s \\ s+(m-1) \\ s+(m-1)+1 \end{bmatrix} + 189 \cdot 2^{k+s+2m} \\
& = 2^6 \cdot (21 \cdot [2^{k+s+3(m-1)-2} + 26 \cdot 2^{k+s+2(m-1)-3}]) + 189 \cdot 2^{k+s+2m} \\
& = 21 \cdot [2^{k+s+3m-2} \cdot 2^{3(2-1)} + a_2 \cdot 2^{k+s+2m-3} \cdot 2^{3(2-1)}] \\
& = 21 \cdot [2^{k+s+3m+1} + 35 \cdot 2^{k+s+2m+1} - 9 \cdot 2^{k+s+2m}] \quad \text{and so } (H_j) \text{ holds for } j=2
\end{aligned}$$

Assume now that (H_{j-1}) holds, we get by (11.48)

$$\begin{aligned}
& \Delta_k \Gamma_{s+m+j} \begin{bmatrix} s \\ s+m \\ s+m+j \end{bmatrix} = 2^6 \cdot \Delta_k \Gamma_{s+(m-1)+(j-1)} \begin{bmatrix} s \\ s+(m-1) \\ s+(m-1)+(j-1) \end{bmatrix} + 189 \cdot 2^{k+s+2m+3j-6} \\
& = 2^6 \cdot 21 \cdot [2^{k+s+3(m-1)+3(j-1)-5} + 35 \cdot 2^{k+s+2(m-1)+4(j-1)-7} - 9 \cdot 2^{k+s+2(m-1)+3(j-1)-6}] \\
& + 189 \cdot 2^{k+s+2m+3j-6} = 21 \cdot [2^{k+s+3m+3j-5} + 35 \cdot 2^{k+s+2m+4j-7} - 9 \cdot 2^{k+s+2m+3j-6}]
\end{aligned}$$

□

11.13. **Computation of** $\Delta_k \Gamma_{2s+m} \begin{bmatrix} s \\ s+m \\ 2s+m \end{bmatrix}$ for $m \geq 1$.

Lemma 11.14. *We have :*

$$(11.50) \quad \Delta_k \Gamma_{2s+m}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} = 9 \cdot 2^{2k+2s+m-2} + 21 \cdot 2^{k+4s+3m-5} + 735 \cdot 2^{k+5s+2m-7} - 477 \cdot 2^{k+4s+2m-6}$$

Proof.

We obtain using (11.46) and (7.48)

$$\begin{aligned} \Delta_k \Gamma_{2s+m}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+m-1}^{\begin{bmatrix} s \\ s+m \\ s+m-1 \end{bmatrix}} &= -3 \cdot 2^{2k+2s+m-2} - 33 \cdot 2^{k+4s+2m-6} \\ \Downarrow \\ 4 \cdot \Delta_k \Gamma_{2s+m-1}^{\begin{bmatrix} s \\ s+m \\ s+m-1 \end{bmatrix}} - 32 \cdot \Delta_k \Gamma_{2s+m-2}^{\begin{bmatrix} s \\ s+m \\ s+m-1 \end{bmatrix}} + \Delta_k \omega_{2s+m}(s, m, 0) &= -3 \cdot 2^{2k+2s+m-2} - 33 \cdot 2^{k+4s+2m-6} \\ \Downarrow \\ \Delta_k \Gamma_{2s+m-1}^{\begin{bmatrix} s \\ s+m \\ s+m-1 \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+m-2}^{\begin{bmatrix} s \\ s+m \\ s+m-1 \end{bmatrix}} \\ &= \frac{1}{4} \cdot [9 \cdot 2^{2k+2s+m-2} - 3 \cdot 2^{2k+2s+m-2} - 33 \cdot 2^{k+4s+2m-6}] = 3 \cdot 2^{2k+2s+m-3} - 33 \cdot 2^{k+4s+2m-8} \end{aligned}$$

We then deduce from (11.49) with $m \rightarrow m-1 (\geq 1)$ and $j \rightarrow s-1$

$$\begin{aligned} \Delta_k \Gamma_{2s+m}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} &= 8 \cdot \left[8 \cdot \Delta_k \Gamma_{2s+m-2}^{\begin{bmatrix} s \\ s+m \\ s+m-1 \end{bmatrix}} + 3 \cdot 2^{2k+2s+m-3} - 33 \cdot 2^{k+4s+2m-8} \right] - 3 \cdot 2^{2k+2s+m-2} - 33 \cdot 2^{k+4s+2m-6} \\ &\Leftrightarrow \Delta_k \Gamma_{2s+m}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} = 2^6 \cdot \Delta_k \Gamma_{s+(m-1)+(s-1)}^{\begin{bmatrix} s \\ s+(m-1) \\ s+(m-1) \end{bmatrix}} + 9 \cdot 2^{2k+2s+m-2} - 99 \cdot 2^{k+4s+2m-6} \\ &= 2^6 \cdot 21 \cdot [2^{k+s+3(m-1)+3(s-1)-5} + 35 \cdot 2^{k+s+2(m-1)+4(s-1)-7} - 9 \cdot 2^{k+s+2(m-1)+3(s-1)-6}] \\ &\quad + 9 \cdot 2^{2k+2s+m-2} - 99 \cdot 2^{k+4s+2m-6} \\ &= 9 \cdot 2^{2k+2s+m-2} + 21 \cdot 2^{k+4s+3m-5} + 735 \cdot 2^{k+5s+2m-7} - 477 \cdot 2^{k+4s+2m-6} \end{aligned}$$

□

11.14. Computation of $\Delta_k \Gamma_{2s+m+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}}$ for $m \geq 2, 1 \leq j \leq m$.

Lemma 11.15. *We have the following reduction formulas for $2 \leq j \leq m-1$*

$$(11.51) \quad \Delta_k \Gamma_{2s+m+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} = 2^6 \cdot \Delta_k \Gamma_{2s+(m-1)+(j-1)}^{\begin{bmatrix} s \\ s+(m-1) \\ s+(m-1) \end{bmatrix}} - 189 \cdot 2^{2k+2s+m+3j-5}$$

$$(11.52) \quad (H_j) \quad \Delta_k \Gamma_{2s+m+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} = 16^{2(j-1)} \cdot \Delta_{k-2(j-1)} \Gamma_{2s+(m-j+1)+1}^{\begin{bmatrix} s \\ s+(m-j+1) \\ s+(m-j+1) \end{bmatrix}} \quad \text{holds for } 2 \leq j \leq m$$

Applying (11.51) and (11.52) we obtain by induction on j the following formula

$$(11.53) \quad \Delta_k \Gamma_{2s+m+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} = \begin{cases} 63 \cdot 2^{2k+2s+m+3j-5} + 21 \cdot 2^{k+4s+3m+3j-5} \\ + 735 \cdot 2^{k+5s+2m+4j-7} - 945 \cdot 2^{k+4s+2m+4j-7} & \text{if } 1 \leq j \leq m-1 \\ 159 \cdot 2^{2k+2s+4m-5} + 735 \cdot 2^{k+5s+6m-7} - 1629 \cdot 2^{k+4s+6m-7} & \text{if } j = m \end{cases}$$

Proof. Proof of (11.53) in the case $j = 1$

We have from (11.46) and (7.48)

$$\Delta_k \Gamma_{2s+m+1}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+m}^{\begin{bmatrix} s \\ s+m \\ s+m-1 \end{bmatrix}} = -69 \cdot 2^{2k+2s+m-2} + 3 \cdot 2^{k+4s+2m-3}$$

$\Updownarrow$

$$4 \cdot \Delta_k \Gamma_{2s+m}^{\left[\begin{smallmatrix} s \\ s+m \\ s+m-1 \end{smallmatrix} \right]} - 32 \cdot \Delta_k \Gamma_{2s+m-1}^{\left[\begin{smallmatrix} s \\ s+m-1 \\ s+m-1 \end{smallmatrix} \right]} + \Delta_k \omega_{2s+m+1}(s, m, 0) = -69 \cdot 2^{2k+2s+m-2} + 3 \cdot 2^{k+4s+2m-3}$$

$\Updownarrow$

$$\begin{aligned} & \Delta_k \Gamma_{2s+m}^{\left[\begin{smallmatrix} s \\ s+m \\ s+m-1 \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{2s+m-1}^{\left[\begin{smallmatrix} s \\ s+m-1 \\ s+m-1 \end{smallmatrix} \right]} \\ &= \frac{1}{4} \cdot [-9 \cdot 2^{2k+2s+m-2} - 69 \cdot 2^{2k+2s+m-2} + 3 \cdot 2^{k+4s+2m-3}] = -39 \cdot 2^{2k+2s+m-3} + 3 \cdot 2^{k+4s+2m-5} \end{aligned}$$

We then deduce from (11.50) with $m \rightarrow m-1 (\geq 1)$ and $j = s$

$$\Delta_k \Gamma_{2s+m+1}^{\left[\begin{smallmatrix} s \\ s+m \\ s+m \end{smallmatrix} \right]} = 8 \cdot \left[8 \cdot \Delta_k \Gamma_{2s+(m-1)}^{\left[\begin{smallmatrix} s \\ s+m-1 \\ s+m-1 \end{smallmatrix} \right]} - 39 \cdot 2^{2k+2s+m-3} + 3 \cdot 2^{k+4s+2m-5} \right] - 69 \cdot 2^{2k+2s+m-2} + 3 \cdot 2^{k+4s+2m-3}$$

$\Updownarrow$

$$\begin{aligned} & \Delta_k \Gamma_{2s+m+1}^{\left[\begin{smallmatrix} s \\ s+m \\ s+m \end{smallmatrix} \right]} = 2^6 \cdot \Delta_k \Gamma_{2s+(m-1)}^{\left[\begin{smallmatrix} s \\ s+m-1 \\ s+m-1 \end{smallmatrix} \right]} - 225 \cdot 2^{2k+2s+m-2} + 9 \cdot 2^{k+4s+2m-3} \\ &= 2^6 \cdot [9 \cdot 2^{2k+2s+(m-1)-2} + 21 \cdot 2^{k+4s+3(m-1)-5} + 735 \cdot 2^{k+5s+2(m-1)-7} - 477 \cdot 2^{k+4s+2(m-1)-6}] \\ &\quad - 225 \cdot 2^{2k+2s+m-2} + 9 \cdot 2^{k+4s+2m-3} \\ &= 63 \cdot 2^{2k+2s+m-2} + 21 \cdot 2^{k+4s+3m-2} + 735 \cdot 2^{k+5s+2m-3} - 945 \cdot 2^{k+4s+2m-3} \end{aligned}$$

Proof of (11.53) in the case $2 \leq j \leq m-1$

We have from (11.46) and (8.3)

$$\begin{aligned} & \Delta_k \Gamma_{2s+m+j}^{\left[\begin{smallmatrix} s \\ s+m \\ s+m \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{2s+m+j-1}^{\left[\begin{smallmatrix} s \\ s+m-1 \\ s+m-1 \end{smallmatrix} \right]} = -63 \cdot 2^{2k+2s+m+3j-5} \\ & \Leftrightarrow 4 \cdot \Delta_k \Gamma_{2s+m+j-1}^{\left[\begin{smallmatrix} s \\ s+m-1 \\ s+m-1 \end{smallmatrix} \right]} - 32 \cdot \Delta_k \Gamma_{2s+m+j-2}^{\left[\begin{smallmatrix} s \\ s+m-1 \\ s+m-1 \end{smallmatrix} \right]} + \Delta_k \omega_{2s+m+j}(s, m, 0) = -63 \cdot 2^{2k+2s+m+3j-5} \end{aligned}$$

We get from (7.48)

$$\Delta_k \omega_{2s+m+j}(s, m, 0) = 0 \Leftrightarrow m + 2s + 2 \leq 2s + m + j \leq 2m + 3s - 1 \Leftrightarrow 2 \leq j \leq m-1$$

$$\text{Then } \Delta_k \Gamma_{2s+m+j-1}^{\left[\begin{smallmatrix} s \\ s+m-1 \\ s+m-1 \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{2s+m+j-2}^{\left[\begin{smallmatrix} s \\ s+m-1 \\ s+m-1 \end{smallmatrix} \right]} = \frac{1}{4} \cdot [-63 \cdot 2^{2k+2s+m+3j-5}] = -63 \cdot 2^{2k+2s+m+3j-7}$$

We then deduce (11.51)

$$\Delta_k \Gamma_{2s+m+j}^{\left[\begin{smallmatrix} s \\ s+m \\ s+m \end{smallmatrix} \right]} = 8 \cdot \left[8 \cdot \Delta_k \Gamma_{2s+m+j-2}^{\left[\begin{smallmatrix} s \\ s+m-1 \\ s+m-1 \end{smallmatrix} \right]} - 63 \cdot 2^{2k+2s+m+3j-7} \right] - 63 \cdot 2^{2k+2s+m+3j-5}$$

$\Updownarrow$

$$\Delta_k \Gamma_{2s+m+j}^{\left[\begin{smallmatrix} s \\ s+m \\ s+m \end{smallmatrix} \right]} = 2^6 \cdot \Delta_k \Gamma_{2s+(m-1)+(j-1)}^{\left[\begin{smallmatrix} s \\ s+m-1 \\ s+m-1 \end{smallmatrix} \right]} - 189 \cdot 2^{2k+2s+m+3j-5}$$

We shall now prove by induction on j that (H_j) holds for $2 \leq j \leq m-1$.

$$(H_j) \quad \Delta_k \Gamma_{2s+m+j}^{\left[\begin{smallmatrix} s \\ s+m \\ s+m \end{smallmatrix} \right]} = 16^{2(j-1)} \cdot \Delta_{k-2(j-1)} \Gamma_{2s+(m-j+1)+1}^{\left[\begin{smallmatrix} s \\ s+(m-j+1) \\ s+(m-j+1) \end{smallmatrix} \right]} \quad \text{if } 2 \leq j \leq m-1$$

(H_j) holds for $j = 2$

From (11.51) with $j = 2$ and (11.53) with $j = 1$, $m \rightarrow m - 1$ we obtain:

$$\begin{aligned}
\Delta_k \Gamma_{2s+m+2}^{\begin{bmatrix} s \\ s+m \end{bmatrix}} &= 2^6 \cdot \Delta_k \Gamma_{2s+(m-1)+1}^{\begin{bmatrix} s+(m-1) \\ s+(m-1) \end{bmatrix}} - 189 \cdot 2^{2k+2s+m+1} \\
&= 2^6 \cdot [63 \cdot 2^{2k+2s+(m-1)-2} + 21 \cdot 2^{k+4s+3(m-1)-2} + 735 \cdot 2^{k+5s+2(m-1)-3} - 945 \cdot 2^{k+4s+2(m-1)-3}] - 189 \cdot 2^{2k+2s+m+1} \\
&= 63 \cdot 2^{2k+2s+m+1} + 21 \cdot 2^{k+4s+3m+1} + 735 \cdot 2^{k+5s+2m+1} - 945 \cdot 2^{k+4s+2m+1} \\
16^2 \cdot \Delta_{k-2} \Gamma_{2s+(m-1)+1}^{\begin{bmatrix} s+(m-1) \\ s+(m-1) \end{bmatrix}} & \\
&= 2^8 \cdot [63 \cdot 2^{2(k-2)+2s+(m-1)-2} + 21 \cdot 2^{(k-2)+4s+3(m-1)-2} + 735 \cdot 2^{(k-2)+5s+2(m-1)-3} - 945 \cdot 2^{(k-2)+4s+2(m-1)-3}] \\
&= 63 \cdot 2^{2k+2s+m+1} + 21 \cdot 2^{k+4s+3m+1} + 735 \cdot 2^{k+5s+2m+1} - 945 \cdot 2^{k+4s+2m+1}
\end{aligned}$$

Assume (H_j) holds for $j = r$

We shall show that (H_j) holds for $j = r + 1$

Indeed we have from (11.53) in the case $m \rightarrow m - r$, $k \rightarrow k - 2(r - 1)$, $j = 1$

(H_j) holds for $j = r$ with $m \rightarrow m - 1$

$\Updownarrow$

$$\begin{aligned}
\Delta_k \Gamma_{2s+(m-1)+r}^{\begin{bmatrix} s+(m-1) \\ s+(m-1) \end{bmatrix}} &= 16^{2(r-1)} \cdot \Delta_{k-2(r-1)} \Gamma_{2s+((m-1)-r)+1}^{\begin{bmatrix} s+((m-1)-r+1) \\ s+((m-1)-r+1) \end{bmatrix}} \\
&= 2^{8r-8} \cdot [63 \cdot 2^{2(k-2(r-1))+2s+((m-1)-r+1)-2} + 21 \cdot 2^{(k-2(r-1))+4s+3((m-1)-r+1)-2} \\
&\quad + 735 \cdot 2^{(k-2(r-1))+5s+2((m-1)-r+1)-3} - 945 \cdot 2^{(k-r+1)+4s+2((m-1)-r+1)-3}] \\
&= 63 \cdot 2^{2k+2s+m+3r-6} + 21 \cdot 2^{k+4s+3m+3r-8} + 735 \cdot 2^{k+5s+2m+4r-9} - 945 \cdot 2^{k+4s+2m+4r-9}
\end{aligned}$$

Now from (11.51) with $j = r + 1$

$$\begin{aligned}
\Delta_k \Gamma_{2s+m+r+1}^{\begin{bmatrix} s \\ s+m \end{bmatrix}} &= 2^6 \cdot \Delta_k \Gamma_{2s+(m-1)+r}^{\begin{bmatrix} s+(m-1) \\ s+(m-1) \end{bmatrix}} - 189 \cdot 2^{2k+2s+m+3(r+1)-5} \\
&= 2^6 \cdot [63 \cdot 2^{2k+2s+m+3r-6} + 21 \cdot 2^{k+4s+3m+3r-8} \\
&\quad + 735 \cdot 2^{k+5s+2m+4r-9} - 945 \cdot 2^{k+4s+2m+4r-9}] - 189 \cdot 2^{2k+2s+m+3(r+1)-5} \\
&= 63 \cdot 2^{2k+2s+m+3r-2} + 21 \cdot 2^{k+4s+3m+3r-2} + 735 \cdot 2^{k+5s+2m+4r-3} - 945 \cdot 2^{k+4s+2m+4r-3}.
\end{aligned}$$

On the other hand

$$\begin{aligned}
16^{2((r+1)-1)} \cdot \Delta_{k-2((r+1)-1)} \Gamma_{2s+(m-(r+1)+1)+1}^{\begin{bmatrix} s+(m-(r+1)+1) \\ s+(m-(r+1)+1) \end{bmatrix}} &= 2^{8r+4} \cdot \Delta_{k-2r} \Gamma_{2s+(m-r)+1}^{\begin{bmatrix} s+(m-r) \\ s+(m-r) \end{bmatrix}} \\
&= 2^{8r} \cdot [63 \cdot 2^{2(k-2r)+2s+(m-r)-2} + 21 \cdot 2^{(k-2r)+4s+3(m-r)-2} + 735 \cdot 2^{(k-2r)+5s+2(m-r)-3} - 945 \cdot 2^{(k-2r)+4s+2(m-r)-3}] \\
&= 63 \cdot 2^{2k+2s+m+3r-2} + 21 \cdot 2^{k+4s+3m+3r-2} + 735 \cdot 2^{k+5s+2m+4r-3} - 945 \cdot 2^{k+4s+2m+4r-3}
\end{aligned}$$

Then we have for $2 \leq j \leq m - 1$

$$\begin{aligned}
\Delta_k \Gamma_{2s+m+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix}} & \\
&= 63 \cdot 2^{2k+2s+m+3j-2} + 21 \cdot 2^{k+4s+3m+3j-2} + 735 \cdot 2^{k+5s+2m+4j-3} - 945 \cdot 2^{k+4s+2m+4j-3}
\end{aligned}$$

Proof of (11.53) in the case $j = m$

We have from (11.46) with $j = m$

$$\Delta_k \Gamma_{2s+2m}^{\begin{bmatrix} s \\ s+m \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+2m-1}^{\begin{bmatrix} s+m-1 \\ s+m \end{bmatrix}} = -159 \cdot 2^{2k+2s+4m-5}$$

$$\Leftrightarrow 4 \cdot \Delta_k \Gamma_{2s+2m-1}^{\begin{bmatrix} s \\ s+m \\ s+m-1 \end{bmatrix}} - 32 \cdot \Delta_k \Gamma_{2s+2m-2}^{\begin{bmatrix} s \\ s+m-1 \\ s+m-1 \end{bmatrix}} + \Delta_k \omega_{2s+m+j}(s, m, 0) = -159 \cdot 2^{2k+2s+4m-5}$$

Using (7.48) we obtain

$$\Delta_k \omega_{2s+2m}(s, m, 0) = 0 \Leftrightarrow m + 2s + 2 \leq 2s + 2m \leq 2m + 3s - 1 \Leftrightarrow 2 \leq m$$

$$\text{Then } \Delta_k \Gamma_{2s+2m-1}^{\begin{bmatrix} s \\ s+m \\ s+m-1 \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+2m-2}^{\begin{bmatrix} s \\ s+m-1 \\ s+m-1 \end{bmatrix}} = \frac{1}{4} \cdot [-159 \cdot 2^{2k+2s+4m-5}] = -159 \cdot 2^{2k+2s+4m-7}$$

We then deduce

$$\begin{aligned} \Delta_k \Gamma_{2s+2m}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} &= 8 \cdot \left[8 \cdot \Delta_k \Gamma_{2s+2m-2}^{\begin{bmatrix} s \\ s+m-1 \\ s+m-1 \end{bmatrix}} - 159 \cdot 2^{2k+2s+4m-7} \right] - 159 \cdot 2^{2k+2s+4m-5} \\ \Leftrightarrow \Delta_k \Gamma_{2s+2m}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} &= 2^6 \cdot \Delta_k \Gamma_{2s+2(m-1)}^{\begin{bmatrix} s \\ s+(m-1) \\ s+(m-1) \end{bmatrix}} - 477 \cdot 2^{2k+2s+4m-5} \end{aligned}$$

We obtain successively

$$\begin{aligned} \Delta_k \Gamma_{2s+2m}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} &= 2^6 \cdot \Delta_k \Gamma_{2s+2(m-1)}^{\begin{bmatrix} s \\ s+(m-1) \\ s+(m-1) \end{bmatrix}} - 477 \cdot 2^{2k+2s+4m-5} \\ 2^6 \cdot \Delta_k \Gamma_{2s+2(m-1)}^{\begin{bmatrix} s \\ s+(m-1) \\ s+(m-1) \end{bmatrix}} &= 2^6 \cdot \left[2^6 \cdot \Delta_k \Gamma_{2s+2(m-2)}^{\begin{bmatrix} s \\ s+(m-2) \\ s+(m-2) \end{bmatrix}} - 477 \cdot 2^{2k+2s+4(m-1)-5} \right] \\ (2^6)^2 \cdot \Delta_k \Gamma_{2s+2(m-2)}^{\begin{bmatrix} s \\ s+(m-2) \\ s+(m-2) \end{bmatrix}} &= (2^6)^2 \cdot \left[2^6 \cdot \Delta_k \Gamma_{2s+2(m-3)}^{\begin{bmatrix} s \\ s+(m-3) \\ s+(m-3) \end{bmatrix}} - 477 \cdot 2^{2k+2s+4(m-2)-5} \right] \\ &\vdots \\ (2^6)^{m-2} \cdot \Delta_k \Gamma_{2s+2(m-(m-2))}^{\begin{bmatrix} s \\ s+(m-(m-2)) \\ s+(m-(m-2)) \end{bmatrix}} &= (2^6)^{m-2} \cdot \left[2^6 \cdot \Delta_k \Gamma_{2s+2(m-(m-1))}^{\begin{bmatrix} s \\ s+(m-(m-1)) \\ s+(m-(m-1)) \end{bmatrix}} - 477 \cdot 2^{2k+2s+4(m-(m-2))-5} \right] \end{aligned}$$

By summing the above equations we get

$$\Delta_k \Gamma_{2s+2m}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} = (2^6)^{m-1} \cdot \Delta_k \Gamma_{2(s+1)}^{\begin{bmatrix} s \\ s+1 \\ s+1 \end{bmatrix}} - 477 \cdot 2^{2k+2s+4m-5} \cdot \sum_{i=0}^{m-2} 2^{2i}.$$

From (9.13) with $j = s + 1$ and $s \rightarrow s + 1$ we obtain

$$\begin{aligned} \Delta_k \Gamma_{2s+2m}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} &= 2^{6m-6} \cdot [159 \cdot 2^{2k+2s-1} + 735 \cdot 2^{k+5s-1} - 1629 \cdot 2^{k+4s-1}] - 159 \cdot 2^{2k+2s+4m-5} \cdot (2^{2m-2} - 1) \\ &= 159 \cdot 2^{2k+2s+4m-5} + 735 \cdot 2^{k+5s+6m-7} - 1629 \cdot 2^{k+4s+6m-7} \end{aligned}$$

(H_j) holds for $j = m$

Indeed we have

$$\begin{aligned} &16^{2(m-1)} \cdot \Delta_{k-2(m-1)} \Gamma_{2s+2}^{\begin{bmatrix} s \\ s+1 \\ s+1 \end{bmatrix}} \\ &= 2^{8m-8} \cdot [159 \cdot 2^{2(k-2(m-1))+2s-1} + 735 \cdot 2^{(k-2(m-1))+5s-1} - 1629 \cdot 2^{(k-2(m-1))+4s-1}] \\ &= 159 \cdot 2^{2k+2s+4m-5} + 735 \cdot 2^{k+5s+6m-7} - 1629 \cdot 2^{k+4s+6m-7} \end{aligned}$$

□

11.15. **Computation of $\Delta_k \Gamma_{2s+2m+1+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}}$ for $m \geq 2$, $0 \leq j \leq s-1$.**

Lemma 11.16. *We have the following reduction formula for $0 \leq j \leq s-1$*

$$(11.54) \quad (H_j) \quad \Delta_k \Gamma_{2s+2m+j+1}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} = 16^{2m+3j} \cdot \Delta_{k-(2m+3j)} \Gamma_{2(s-j)+1}^{\begin{bmatrix} s-j \\ s-j \\ s-j \end{bmatrix}},$$

furthermore

$$(11.55) \quad \Delta_k \Gamma_{2s+2m+1+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} = \begin{cases} 315 \cdot 2^{2k+2s+4m+4j-2} + 105 \cdot (7 \cdot 2^{k+5s+6m+4j-3} - 31 \cdot 2^{k+4s+6m+5j-3}) & \text{if } 0 \leq j \leq s-2 \\ 7 \cdot 2^{3k+3s+2m-3} - 21 \cdot 2^{2k+6s+4m-6} + 7 \cdot 2^{k+9s+6m-8} & \text{if } j = s-1 \end{cases}$$

Proof. **The case $0 \leq j \leq s-2$**

We have from (11.47) and (8.3) with $l = 0$

$$\begin{aligned} \Delta_k \Gamma_{2s+2m+1+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+2m+j}^{\begin{bmatrix} s \\ s+m-1 \\ s+m \end{bmatrix}} &= -315 \cdot 2^{2k+2s+4m+4j-2} \\ \Leftrightarrow 4 \cdot \Delta_k \Gamma_{2s+2m+j}^{\begin{bmatrix} s \\ s+m-1 \\ s+m \end{bmatrix}} - 32 \cdot \Delta_k \Gamma_{2s+2m+j-1}^{\begin{bmatrix} s \\ s+m-1 \\ s+m-1 \end{bmatrix}} + \Delta_k \omega_{2s+2m+1+j}(s, m, 0) &= -315 \cdot 2^{2k+2s+4m+4j-2} \end{aligned}$$

Now (7.48) yields

$$\begin{aligned} \Delta_k \omega_{2s+2m+1+j}(s, m, 0) = 0 &\Leftrightarrow m + 2s + 2 \leq 2s + 2m + 1 + j \leq 2m + 3s - 1 \\ \Leftrightarrow \sup(1 - m, 0) \leq j &\leq s - 2 \end{aligned}$$

$$\text{Then } \Delta_k \Gamma_{2s+2m+j}^{\begin{bmatrix} s \\ s+m-1 \\ s+m \end{bmatrix}} - 8 \cdot \Delta_k \Gamma_{2s+2m+j-1}^{\begin{bmatrix} s \\ s+m-1 \\ s+m-1 \end{bmatrix}} = \frac{1}{4} \cdot [-315 \cdot 2^{2k+2s+4m+4j-2}] = -315 \cdot 2^{2k+2s+4m+4j-4}$$

We then deduce

$$\begin{aligned} \Delta_k \Gamma_{2s+2m+1+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} &= 8 \cdot \left[8 \cdot \Delta_k \Gamma_{2s+2m+j-1}^{\begin{bmatrix} s \\ s+m-1 \\ s+m-1 \end{bmatrix}} - 315 \cdot 2^{2k+2s+4m+4j-4} \right] - 315 \cdot 2^{2k+2s+4m+4j-2} \\ \Leftrightarrow \Delta_k \Gamma_{2s+2m+1+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} &= 2^6 \cdot \Delta_k \Gamma_{2s+2(m-1)+j+1}^{\begin{bmatrix} s \\ s+(m-1) \\ s+(m-1) \end{bmatrix}} - 945 \cdot 2^{2k+2s+4m+4j-2} \end{aligned}$$

We obtain successively

$$\begin{aligned} \Delta_k \Gamma_{2s+2m+1+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} &= 2^6 \cdot \Delta_k \Gamma_{2s+2(m-1)+j+1}^{\begin{bmatrix} s \\ s+(m-1) \\ s+(m-1) \end{bmatrix}} - 945 \cdot 2^{2k+2s+4m+4j-2} \\ 2^6 \cdot \Delta_k \Gamma_{2s+2(m-1)+j+1}^{\begin{bmatrix} s \\ s+(m-1) \\ s+(m-1) \end{bmatrix}} &= 2^6 \cdot \left[2^6 \cdot \Delta_k \Gamma_{2s+2(m-2)+j+1}^{\begin{bmatrix} s \\ s+(m-2) \\ s+(m-2) \end{bmatrix}} - 945 \cdot 2^{2k+2s+4(m-1)+4j-2} \right] \\ (2^6)^2 \cdot \Delta_k \Gamma_{2s+2(m-2)+j+1}^{\begin{bmatrix} s \\ s+(m-2) \\ s+(m-2) \end{bmatrix}} &= (2^6)^2 \cdot \left[2^6 \cdot \Delta_k \Gamma_{2s+2(m-3)+j+1}^{\begin{bmatrix} s \\ s+(m-3) \\ s+(m-3) \end{bmatrix}} - 945 \cdot 2^{2k+2s+4(m-2)+4j-2} \right] \\ &\vdots \end{aligned}$$

$$\begin{aligned}
& (2^6)^{m-2} \cdot \Delta_k \Gamma_{2s+2(m-(m-2))+j+1}^{\left[\begin{smallmatrix} s+(m-(m-2)) \\ s+(m-(m-2)) \end{smallmatrix} \right]} \\
&= (2^6)^{m-2} \cdot \left[2^6 \cdot \Delta_k \Gamma_{2s+2(m-(m-1))+j+1}^{\left[\begin{smallmatrix} s+(m-(m-1)) \\ s+(m-(m-1)) \end{smallmatrix} \right]} - 945 \cdot 2^{2k+2s+4(m-(m-2))+4j-2} \right]
\end{aligned}$$

By summing the above equations we get

$$\begin{aligned}
\Delta_k \Gamma_{2s+2m+j+1}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right]} &= (2^6)^{m-1} \cdot \Delta_k \Gamma_{2(s+1)+j+1}^{\left[\begin{smallmatrix} s+1 \\ s+1 \end{smallmatrix} \right]} - 945 \cdot 2^{2k+2s+4m+4j-2} \cdot \sum_{i=0}^{m-2} 2^{2i} \\
&= 2^{6m-6} \cdot [105 \cdot (3 \cdot 2^{2k+2s+4j+2} + 7 \cdot 2^{k+5s+4j+3} - 31 \cdot 2^{k+4s+5j+3}) \\
&\quad - 315 \cdot 2^{2k+2s+4m+4j-2} \cdot (2^{2m-2} - 1)] \\
&= 315 \cdot 2^{2k+2s+4m+4j-2} + 105 \cdot (7 \cdot 2^{k+5s+6m+4j-3} - 31 \cdot 2^{k+4s+6m+5j-3})
\end{aligned}$$

On the other hand we have by (9.7) with $k \longrightarrow k - (2m + 3j)$, $s \longrightarrow s - j$, $j = 1$

$$\begin{aligned}
& 16^{2m+3j} \cdot \Delta_{k-(2m+3j)} \Gamma_{2(s-j)+1}^{\left[\begin{smallmatrix} s-j \\ s-j \\ s-j \end{smallmatrix} \right]} = 2^{8m+12j} \cdot \Delta_{k-(2m+3j)} \Gamma_{2(s-j)+1}^{\left[\begin{smallmatrix} s-j \\ s-j \\ s-j \end{smallmatrix} \right]} \\
&= 2^{8m+12j} \cdot [105 \cdot (3 \cdot 2^{2(k-(2m+3j))+2(s-j)-2} + 7 \cdot 2^{(k-(2m+3j))+5(s-j)-3} \\
&\quad - 31 \cdot 2^{(k-(2m+3j))+4(s-j)-3})] \\
&= 315 \cdot 2^{2k+2s+4m+4j-2} + 105 \cdot (7 \cdot 2^{k+5s+6m+4j-3} - 31 \cdot 2^{k+4s+6m+5j-3})
\end{aligned}$$

Thus we get (11.54) and (11.55) in the case $0 \leq j \leq s - 2$

The case $j = s - 1$

We have by (11.47) and (8.3) with $l = 0$

$$\begin{aligned}
& \Delta_k \Gamma_{3s+2m}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{3s+2m-1}^{\left[\begin{smallmatrix} s+m-1 \\ s+m-1 \end{smallmatrix} \right]} = -21 \cdot 2^{3k+3s+2m-3} + 21 \cdot 2^{2k+6s+4m-6} \\
&\Leftrightarrow 4 \cdot \Delta_k \Gamma_{3s+2m-1}^{\left[\begin{smallmatrix} s+m-1 \\ s+m-1 \end{smallmatrix} \right]} - 32 \cdot \Delta_k \Gamma_{3s+2m-2}^{\left[\begin{smallmatrix} s+m-1 \\ s+m-1 \end{smallmatrix} \right]} + \Delta_k \omega_{3s+2m}(s, m, 0) = -21 \cdot 2^{3k+3s+2m-3} + 21 \cdot 2^{2k+6s+4m-6}
\end{aligned}$$

We get by (7.48)

$$\Delta_k \omega_{3s+2m}(s, m, 0) = 21 \cdot 2^{3k+3s+2m-3}$$

Then

$$\begin{aligned}
& \Delta_k \Gamma_{3s+2m-1}^{\left[\begin{smallmatrix} s+m-1 \\ s+m-1 \end{smallmatrix} \right]} - 8 \cdot \Delta_k \Gamma_{3s+2m-2}^{\left[\begin{smallmatrix} s+m-1 \\ s+m-1 \end{smallmatrix} \right]} = \frac{1}{4} \cdot [-21 \cdot 2^{3k+3s+2m-3} - 21 \cdot 2^{3k+3s+2m-3} + 21 \cdot 2^{2k+6s+4m-6}] \\
&= -21 \cdot 2^{3k+3s+2m-4} + 21 \cdot 2^{2k+6s+4m-8}
\end{aligned}$$

We then deduce

$$\begin{aligned}
& \Delta_k \Gamma_{3s+2m}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right]} = 8 \cdot \left[8 \cdot \Delta_k \Gamma_{3s+2m-2}^{\left[\begin{smallmatrix} s+m-1 \\ s+m-1 \end{smallmatrix} \right]} - 21 \cdot 2^{3k+3s+2m-4} + 21 \cdot 2^{2k+6s+4m-8} \right] \\
&\quad - 21 \cdot 2^{3k+3s+2m-3} + 21 \cdot 2^{2k+6s+4m-6} \\
&\Leftrightarrow \Delta_k \Gamma_{3s+2m}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right]} = 2^6 \cdot \Delta_k \Gamma_{3s+2(m-1)}^{\left[\begin{smallmatrix} s+(m-1) \\ s+(m-1) \end{smallmatrix} \right]} - 105 \cdot 2^{3k+3s+2m-3} + 63 \cdot 2^{2k+6s+4m-6}
\end{aligned}$$

We obtain successively

$$\begin{aligned}
\Delta_k \Gamma_{3s+2m}^{\begin{bmatrix} s \\ s+m \end{bmatrix}} &= 2^6 \cdot \Delta_k \Gamma_{3s+2(m-1)}^{\begin{bmatrix} s \\ s+(m-1) \end{bmatrix}} - 105 \cdot 2^{3k+3s+2m-3} + 63 \cdot 2^{2k+6s+4m-6} \\
2^6 \cdot \Delta_k \Gamma_{2s+2(m-1)}^{\begin{bmatrix} s \\ s+(m-1) \end{bmatrix}} &= 2^6 \cdot \left[2^6 \cdot \Delta_k \Gamma_{2s+2(m-2)}^{\begin{bmatrix} s \\ s+(m-2) \end{bmatrix}} - 105 \cdot 2^{3k+3s+2(m-1)-3} + 63 \cdot 2^{2k+6s+4(m-1)-6} \right] \\
(2^6)^2 \cdot \Delta_k \Gamma_{3s+2(m-2)}^{\begin{bmatrix} s \\ s+(m-2) \end{bmatrix}} &= (2^6)^2 \cdot \left[2^6 \cdot \Delta_k \Gamma_{3s+2(m-3)}^{\begin{bmatrix} s \\ s+(m-3) \end{bmatrix}} - 105 \cdot 2^{3k+3s+2(m-1)-3} + 63 \cdot 2^{2k+6s+4(m-1)-6} \right] \\
&\vdots \\
(2^6)^{m-2} \cdot \Delta_k \Gamma_{3s+2(m-(m-2))}^{\begin{bmatrix} s \\ s+(m-(m-2)) \end{bmatrix}} \\
&= (2^6)^{m-2} \cdot \left[2^6 \cdot \Delta_k \Gamma_{3s+2(m-(m-1))}^{\begin{bmatrix} s \\ s+(m-(m-1)) \end{bmatrix}} - 105 \cdot 2^{3k+3s+2(m-(m-2))-3} + 63 \cdot 2^{2k+6s+4(m-(m-2))-6} \right]
\end{aligned}$$

By summing the above equations we get

$$\begin{aligned}
\Delta_k \Gamma_{3s+2m}^{\begin{bmatrix} s \\ s+m \end{bmatrix}} &= (2^6)^{m-1} \cdot \Delta_k \Gamma_{3s+2}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}} + \sum_{i=0}^{m-2} [-105 \cdot 2^{3k+3s+2(m-i)-3} + 63 \cdot 2^{2k+6s+4(m-i)-6}] \cdot 2^{6i} \\
&= 2^{6m-6} \cdot \Delta_k \Gamma_{3s+2}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}} - 105 \cdot 2^{3k+3s+2m-3} \cdot \sum_{i=0}^{m-2} 2^{4i} + 63 \cdot 2^{2k+6s+4m-6} \cdot \sum_{i=0}^{m-2} 2^{2i} \\
&= 2^{6m-6} \cdot [7 \cdot 2^{3k+3s-1} - 21 \cdot 2^{2k+6s-2} + 7 \cdot 2^{k+9s-2}] \\
&\quad - 7 \cdot 2^{3k+3s+2m-3} \cdot (2^{4m-4} - 1) + 21 \cdot 2^{2k+6s+4m-6} \cdot (2^{2m-2} - 1) \\
&= 7 \cdot 2^{3k+3s+2m-3} - 21 \cdot 2^{2k+6s+4m-6} + 7 \cdot 2^{k+9s+6m-8}
\end{aligned}$$

$$(H_{s-1}) \text{ holds} \iff \Delta_k \Gamma_{3s+2m}^{\begin{bmatrix} s \\ s+m \end{bmatrix}} = 16^{2m+3(s-1)} \cdot \Delta_{k-(2m+3(s-1))} \Gamma_3^{\begin{bmatrix} 1 \\ 1 \end{bmatrix}}$$

Indeed we deduce from Lemma 1.23 in the case $m = 0$

$$\begin{aligned}
16^{2m+3(s-1)} \cdot \Delta_{k-(2m+3(s-1))} \Gamma_3^{\begin{bmatrix} 1 \\ 1 \end{bmatrix}} &= 2^{8m+12s-12} \cdot \Delta_{k-(2m+3(s-1))} \Gamma_3^{\begin{bmatrix} 1 \\ 1 \end{bmatrix}} \\
&= 2^{8m+12s-12} \cdot [7 \cdot 2^{3(k-2m-3s+3)} - 21 \cdot 2^{2(k-2m-3s+3)} + 7 \cdot 2^{k-2m-3s+3}] \\
&= 7 \cdot 2^{3k+3s+2m-3} - 21 \cdot 2^{2k+6s+4m-6} + 7 \cdot 2^{k+9s+6m-8}
\end{aligned}$$

□

11.16. Summary of principal results in section 11.

Lemma 11.17. *We have for $m \geq 2$:*

$$(11.56) \quad \Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix}} = \begin{cases} 2^{k+s-1} & \text{if } j = 0 \\ (21 \cdot 2^{3j-4} - 3 \cdot 2^{2j-3}) \cdot 2^{k+s-1} & \text{if } 1 \leq j \leq m-1 \\ 21 \cdot 2^{k+s+3m-5} + 45 \cdot 2^{k+s+2m-4} & \text{if } j = m \end{cases}$$

$$(11.57) \quad \Delta_k \Gamma_{s+m+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix}} = \begin{cases} 21 \cdot [2^{k+s+3m+3j-5} + 35 \cdot 2^{k+s+2m+4j-7} - 9 \cdot 2^{k+s+2m+3j-6}] & \text{if } 1 \leq j \leq s-1 \\ 9 \cdot 2^{2k+2s+m-2} + 21 \cdot 2^{k+4s+3m-5} & \\ + 735 \cdot 2^{k+5s+2m-7} - 477 \cdot 2^{k+4s+2m-6} & \text{if } j = s \end{cases}$$

(11.58)

$$\Delta_k \Gamma_{2s+m+1+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} = \begin{cases} 63 \cdot 2^{2k+2s+m+3j-2} + 21 \cdot 2^{k+4s+3m+3j-2} \\ + 735 \cdot 2^{k+5s+2m+4j-3} - 945 \cdot 2^{k+4s+2m+4j-3} & \text{if } 0 \leq j \leq m-2 \\ 159 \cdot 2^{2k+2s+4m-5} + 735 \cdot 2^{k+5s+6m-7} - 1629 \cdot 2^{k+4s+6m-7} & \text{if } j = m-1 \end{cases}$$

(11.59)

$$\Delta_k \Gamma_{2s+2m+1+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} = \begin{cases} 315 \cdot 2^{2k+2s+4m+4j-2} \\ + 105 \cdot (7 \cdot 2^{k+5s+6m+4j-3} - 31 \cdot 2^{k+4s+6m+5j-3}) & \text{if } 0 \leq j \leq s-2 \\ 7 \cdot 2^{3k+3s+2m-3} - 21 \cdot 2^{2k+6s+4m-6} + 7 \cdot 2^{k+9s+6m-8} & \text{if } j = s-1 \end{cases}$$

Further we have the following reduction formulas

$$(11.60) \quad \Delta_k \Gamma_{2s+m+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} = 16^{2(j-1)} \cdot \Delta_{k-2(j-1)} \Gamma_{2s+(m-j+1)+1}^{\begin{bmatrix} s \\ s+(m-j+1) \\ s+(m-j+1) \end{bmatrix}} \quad \text{if } 2 \leq j \leq m$$

$$(11.61) \quad \Delta_k \Gamma_{2s+2m+j+1}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} = 16^{2m+3j} \cdot \Delta_{k-(2m+3j)} \Gamma_{2(s-j)+1}^{\begin{bmatrix} s-j \\ s-j \\ s-j \end{bmatrix}} \quad \text{if } 0 \leq j \leq s-1$$

The case $m = 1$

$$(11.62) \quad \Delta_k \Gamma_{s+j}^{\begin{bmatrix} s \\ s+1 \\ s+1 \end{bmatrix}} = \begin{cases} 2^{k+s-1} & \text{if } j = 0 \\ 33 \cdot 2^{k+s-1} & \text{if } j = 1 \\ 7 \cdot (105 \cdot 2^{j-2} - 15) \cdot 2^{k+s+3j-7} & \text{if } 2 \leq j \leq s \\ 9 \cdot 2^{2k+2s-1} + 21 \cdot 35 \cdot 2^{k+5s-5} - 393 \cdot 2^{k+4s-4} & \text{if } j = s+1 \\ 159 \cdot 2^{2k+2s-1} + 735 \cdot 2^{k+5s-1} - 1629 \cdot 2^{k+4s-1} & \text{if } j = s+2 \end{cases}$$

(11.63)

$$\Delta_k \Gamma_{2s+1+j}^{\begin{bmatrix} s \\ s+1 \\ s+1 \end{bmatrix}} = \begin{cases} 105 \cdot (3 \cdot 2^{2k+2s+4j-6} + 7 \cdot 2^{k+5s+4j-5} - 31 \cdot 2^{k+4s+5j-7}) & \text{if } 2 \leq j \leq s \\ 7 \cdot 2^{3k+3s-1} - 21 \cdot 2^{2k+6s-2} + 7 \cdot 2^{k+9s-2} & \text{if } j = s+1 \end{cases}$$

Proof. We obtain (11.62) and (11.63) by combining (8.9), (8.10) and (9.5), (9.7). $\square$

12. An explicit formula for $\Gamma_i^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix} \times k}$, for $s \leq i \leq 3s+2m$, $k \geq i$

12.1. Notation.

Definition 12.1. We recall that $\Gamma_i^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix} \times k}$ denotes the number of rank i matrices of the form $\begin{bmatrix} A \\ B \\ C \end{bmatrix}$ where A is a $s \times k$ persymmetric matrix, B a $(s+m) \times k$ persymmetric matrix and C is a $(s+m) \times k$ persymmetric matrix.

12.2. Introduction. We adapt the method in Section 10 to compute explicitly the number

$$\Gamma_i^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix} \times k}.$$

12.3. Computation of $\Gamma_{s+i}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix} \times k} - \Gamma_{s+i}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix} \times (s+i+1)}$, for $0 \leq i \leq 2s+2m$, $k \geq s+i+1$.

Lemma 12.2. *We have for $m \geq 2$, $k \geq s + j + 1$:*

$$(12.1) \quad \Gamma_{s+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix} \times k} - \Gamma_{s+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+j+1)} = \begin{cases} 2^{s-1} \cdot (2^k - 2^{s+1}) & \text{if } j = 0 \\ (21 \cdot 2^{s+3j-5} - 3 \cdot 2^{s+2j-4}) \cdot (2^k - 2^{s+j+1}) & \text{if } 1 \leq j \leq m-1 \\ (21 \cdot 2^{s+3m-5} + 45 \cdot 2^{s+2m-4}) \cdot (2^k - 2^{s+m+1}) & \text{if } j = m, \end{cases}$$

for $m \geq 2$, $k \geq s + m + j + 1$:

$$(12.2) \quad \Gamma_{s+m+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix} \times k} - \Gamma_{s+m+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+m+j+1)} = \begin{cases} (2^k - 2^{s+m+j+1}) \cdot (21 \cdot 2^{s+3m+3j-5} + 735 \cdot 2^{s+2m+4j-7} - 189 \cdot 2^{s+2m+3j-6}) & \text{if } 1 \leq j \leq s-1 \\ 3 \cdot 2^{2s+m-2} \cdot (2^{2k} - 2^{4s+2m+2}) & \\ + (2^k - 2^{2s+m+1}) \cdot (21 \cdot 2^{4s+3m-5} + 735 \cdot 2^{5s+2m-7} - 477 \cdot 2^{4s+2m-6}) & \text{if } j = s, \end{cases}$$

for $m \geq 2$, $k \geq 2s + m + j + 1$:

$$(12.3) \quad \Gamma_{2s+m+1+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix} \times k} - \Gamma_{2s+m+1+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix} \times (2s+m+2+j)} = \begin{cases} (2^k - 2^{2s+m+2+j}) \cdot (21 \cdot 2^{4s+3m+3j-2} + 735 \cdot 2^{5s+2m+4j-3} - 945 \cdot 2^{4s+2m+4j-3}) \\ + 21 \cdot 2^{2s+m+3j-2} \cdot (2^{2k} - 2^{4s+2m+2j+4}) & \text{if } 0 \leq j \leq m-2 \\ 53 \cdot 2^{2s+4m-5} \cdot (2^{2k} - 2^{4s+4m+2}) \\ + (2^k - 2^{2s+2m+1}) \cdot (735 \cdot 2^{5s+6m-7} - 1629 \cdot 2^{4s+6m-7}) & \text{if } j = m-1, \end{cases}$$

for $m \geq 2$, $k \geq 2s + 2m + j + 1$:

$$(12.4) \quad \Gamma_{2s+2m+1+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix} \times k} - \Gamma_{2s+2m+1+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix} \times (2s+2m+2+j)} = \begin{cases} (2^k - 2^{2s+2m+2+j}) \cdot (735 \cdot 2^{5s+6m+4j-3} - 3255 \cdot 2^{4s+6m+5j-3}) \\ + 105 \cdot 2^{2s+4m+4j-2} \cdot (2^{2k} - 2^{4s+4m+2j+4}) & \text{if } 0 \leq j \leq s-2 \\ 2^{3s+2m-3} \cdot (2^{3k} - 2^{9s+6m+3}) - 7 \cdot 2^{6s+4m-6} \cdot (2^{2k} - 2^{6s+4m+2}) \\ + 7 \cdot 2^{9s+6m-8} \cdot (2^k - 2^{3s+2m+1}) & \text{if } j = s-1 \end{cases}$$

We have for $m = 1$, $k \geq s + j + 1$

$$(12.5) \quad \Gamma_{s+j}^{\begin{bmatrix} s \\ s+1 \end{bmatrix} \times k} - \Gamma_{s+j}^{\begin{bmatrix} s \\ s+1 \end{bmatrix} \times (s+j+1)}$$

$$= \begin{cases} 2^{s-1} \cdot (2^k - 2^{s+1}) & \text{if } j = 0 \\ 33 \cdot 2^{s-1} \cdot (2^k - 2^{s+2}) & \text{if } j = 1 \\ (735 \cdot 2^{s+4j-9} - 105 \cdot 2^{s+3j-7}) \cdot (2^k - 2^{s+j+1}) & \text{if } 2 \leq j \leq s \\ 3 \cdot 2^{2s-1} \cdot (2^{2k} - 2^{4s+4}) + (735 \cdot 2^{5s-5} - 393 \cdot 2^{4s-4}) \cdot (2^k - 2^{2s+2}) & \text{if } j = s+1, \\ 53 \cdot 2^{2s-1} \cdot (2^{2k} - 2^{4s+6}) + (735 \cdot 2^{5s-1} - 1629 \cdot 2^{4s-1}) \cdot (2^k - 2^{2s+3}) & \text{if } j = s+2, \end{cases}$$

We have for $m = 1$, $k \geq 2s + 2 + j$

(12.6)

$$\begin{aligned} & \Gamma_{2s+1+j}^{\begin{bmatrix} s+1 \\ s+1 \end{bmatrix} \times k} - \Gamma_{2s+1+j}^{\begin{bmatrix} s+1 \\ s+1 \end{bmatrix} \times (2s+2+j)} \\ &= \begin{cases} 105 \cdot 2^{2s+4j-6} \cdot (2^{2k} - 2^{4s+2j+4}) + (735 \cdot 2^{5s+4j-5} - 3255 \cdot 2^{4s+5j-7}) \cdot (2^k - 2^{2s+j+2}) & \text{if } 2 \leq j \leq s, \\ 2^{3s-1} \cdot (2^{2k} - 2^{9s+6}) - 7 \cdot 2^{6s-2} \cdot (2^{2k} - 2^{6s+6}) + 7 \cdot 2^{9s-2} \cdot (2^k - 2^{3s+3}) & \text{if } j = s+1 \end{cases} \end{aligned}$$

Proof. By way of example we prove (12.2) in the case $j = s$.

We obtain from (11.57) in the case $j = s$

$$\begin{aligned} \Delta_k \Gamma_{2s+m}^{\begin{bmatrix} s+m \\ s+m \end{bmatrix}} &= \Gamma_{2s+m}^{\begin{bmatrix} s+m \\ s+m \end{bmatrix} \times (k+1)} - \Gamma_{2s+m}^{\begin{bmatrix} s+m \\ s+m \end{bmatrix} \times k} \\ &= 9 \cdot 2^{2k+2s+m-2} + 21 \cdot 2^{k+4s+3m-5} + 735 \cdot 2^{k+5s+2m-7} - 477 \cdot 2^{k+4s+2m-6}, \end{aligned}$$

hence the above equality gives

$$\begin{aligned} & \sum_{i=2s+m+1}^k \left(\Gamma_{2s+m}^{\begin{bmatrix} s+m \\ s+m \end{bmatrix} \times (i+1)} - \Gamma_{2s+m}^{\begin{bmatrix} s+m \\ s+m \end{bmatrix} \times i} \right) = 9 \cdot 2^{2s+m-2} \cdot \sum_{i=2s+m+1}^k 2^{2i} \\ & + \left(21 \cdot 2^{4s+3m-5} + 735 \cdot 2^{5s+2m-7} - 477 \cdot 2^{4s+2m-6} \right) \cdot \sum_{i=2s+m+1}^k 2^i \\ & \Updownarrow \\ & \Gamma_{2s+m}^{\begin{bmatrix} s+m \\ s+m \end{bmatrix} \times (k+1)} - \Gamma_{2s+m}^{\begin{bmatrix} s+m \\ s+m \end{bmatrix} \times (2s+m+1)} = 9 \cdot 2^{2s+m-2} \cdot 2^{4s+2m+2} \cdot \left(\frac{(2^2)^{k-2s-m} - 1}{2^2 - 1} \right) \\ & + \left(21 \cdot 2^{4s+3m-5} + 735 \cdot 2^{5s+2m-7} - 477 \cdot 2^{4s+2m-6} \right) \cdot 2^{2s+m+1} \cdot (2^{k-2s-m} - 1) \end{aligned}$$

□

12.4. Computation of $\Gamma_{s+j}^{\begin{bmatrix} s+m \\ s+m \end{bmatrix} \times k}$, for $m \geq 2$, $0 \leq j \leq m$, $k \geq s+m$ and $j = m+1$, $k = s+m+1$.

Lemma 12.3. *We have for $m \geq 2$:*

$$(12.7) \quad \Gamma_{s+j} \begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix} \times k = \begin{cases} 2^{6s+2m-3} - 7 \cdot 2^{4s-6} + 3 \cdot 2^{3s-5} & \text{if } j = 0, k = s \\ 2^{k+s-1} - 2^{2s} + 21 \cdot (5 \cdot 2^{4s-6} - 2^{3s-5}) & \text{if } j = 0, k \geq s+1 \\ 2^{6s+2m} - 7 \cdot 2^{4s-2} + 3 \cdot 2^{3s-2} & \text{if } j = 1, k = s+1 \\ 9 \cdot 2^{k+s-1} + 21 \cdot (5 \cdot 2^{4s-2} - 2^{3s-2} - 2^{2s}) & \text{if } j = 1, k \geq s+2 \\ 2^{6s+2m+3} - 7 \cdot 2^{4s+2} + 3 \cdot 2^{3s+1} + 2^{2s+1} & \text{if } j = 2, k = s+2 \\ 39 \cdot 2^{k+s} + 21 \cdot (5 \cdot 2^{4s+2} - 2^{3s+1} - 9 \cdot 2^{2s+1}) & \text{if } j = 2, k \geq s+3 \\ 2^{6s+2m+6} - 7 \cdot 2^{4s+6} + 3 \cdot 2^{3s+4} + 3 \cdot 2^{2s+4} & \text{if } j = 3, k = s+3 \\ 81 \cdot 2^{k+s+2} + 21 \cdot (5 \cdot 2^{4s+6} - 2^{3s+4} - 19 \cdot 2^{2s+4}) & \text{if } j = 3, k \geq s+4 \\ 2^{6s+2m+9} - 7 \cdot 2^{4s+10} + 3 \cdot 2^{3s+7} + 7 \cdot 2^{2s+7} & \text{if } j = 4, k = s+4 \\ 165 \cdot 2^{k+s+4} + 21 \cdot (5 \cdot 2^{4s+10} - 2^{3s+7} - 39 \cdot 2^{2s+7}) & \text{if } j = 4, k \geq s+5 \\ 2^{6s+2m+12} - 7 \cdot 2^{4s+14} + 3 \cdot 2^{3s+10} + 15 \cdot 2^{2s+10} & \text{if } j = 5, k = s+5 \\ (21 \cdot 2^{j-1} - 3) \cdot 2^{k+s+2j-4} & \\ + 21 \cdot (5 \cdot 2^{4s+4j-6} - 2^{3s+3j-5} - 5 \cdot 2^{2s+4j-6} + 2^{2s+3j-5}) & \text{if } 1 \leq j \leq m-1, k \geq s+j+1 \\ 2^{6s+2m+3j-3} - 7 \cdot 2^{4s+4j-6} + 3 \cdot 2^{3s+3j-5} & \\ + (2^{j-1} - 1) \cdot 2^{2s+3j-5} & \text{if } 1 \leq j \leq m+1, k = s+j \\ (21 \cdot 2^{s+3m-5} + 45 \cdot 2^{s+2m-4}) \cdot (2^k - 2^{s+m+1}) & \\ + 105 \cdot 2^{4s+4m-6} - 21 \cdot 2^{3s+3m-5} - 21 \cdot 2^{2s+4m-6} + 9 \cdot 2^{2s+3m-5} & \text{if } j = m, k \geq s+m+1 \end{cases}$$

Let (H_{q-1}) denote the following statement :

$$\Gamma_{s+j} \begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix} \times k = (21 \cdot 2^{j-1} - 3) \cdot 2^{k+s+2j-4} + 21 \cdot (5 \cdot 2^{4s+4j-6} - 2^{3s+3j-5} - 5 \cdot 2^{2s+4j-6} + 2^{2s+3j-5})$$

for $1 \leq j \leq q-1 \leq m-2, k \geq s+j+1$

Proof. Proof by strong induction

We start by prove that (H_{q-1}) holds for $q = 2, 3, 4, 5$

Next we assume that (H_{q-1}) holds. then we prove that this can be used to show that (H_q) holds, that is :

$$\Gamma_{s+q} \begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix} \times k = (21 \cdot 2^{q-1} - 3) \cdot 2^{k+s+2q-4} + 21 \cdot (5 \cdot 2^{4s+4q-6} - 2^{3s+3q-5} - 5 \cdot 2^{2s+4q-6} + 2^{2s+3q-5})$$

for $2 \leq q \leq m-1, k \geq s+q+1$

The case $j = 0, k = s$

Follows from (6.41) with $l = 0, i = s$

The case $j = 0, k \geq s+1$

Follows from (12.1) with $j = 0$ using (6.42) with $l = 0$

The case $j = 1, k = s+1$

Follows from (6.43) with $l = 0$

The case $j = 1, k \geq s+2$ and the case $j = 2, k = s+2$

From (6.1) with $k = s+2, l = 0$ we obtain

$$\sum_{i=0}^{s+2} \Gamma_i \begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix} \times (s+2) = 2^{6s+2m+3}$$

$\Updownarrow$

$$\sum_{i=0}^{s-1} \Gamma_i \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+2) + \Gamma_s \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+2) + \Gamma_{s+1} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+2) + \Gamma_{s+2} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+2) = 2^{6s+2m+3}$$

$\Downarrow$

using (6.39) and (12.7) in the case $j = 0$, $k \geq s+1$ with $k = s+2$ we obtain

$$(7 \cdot 2^{4s-6} - 3 \cdot 2^{3s-5}) + (2^{2s} + 21 \cdot (5 \cdot 2^{4s-6} - 2^{3s-5})) + \Gamma_{s+1} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+2) + \Gamma_{s+2} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+2) = 2^{6s+2m+3}$$

$\Downarrow$

$$\Gamma_{s+1} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+2) + \Gamma_{s+2} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+2) = 2^{6s+2m+3} - 7 \cdot 2^{4s-2} + 3 \cdot 2^{3s-2} - 2^{2s}$$

From (6.2) with $k = s+2$, $l = 0$ we get

$$\sum_{i=0}^{s+2} \Gamma_i \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+2) \cdot 2^{-i} = 2^{5s+2m+1} + 2^{3s+3} - 2^{2s+1}$$

$\Downarrow$

$$\begin{aligned} & \sum_{i=0}^{s-1} \Gamma_i \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+2) \cdot 2^{-i} + \Gamma_s \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+2) \cdot 2^{-s} + \Gamma_{s+1} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+2) \cdot 2^{-(s+1)} + \Gamma_{s+2} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+2) \cdot 2^{-(s+2)} \\ &= 2^{5s+2m+1} + 2^{3s+3} - 2^{2s+1} \end{aligned}$$

$\Downarrow$ using (6.40) and (12.7) in the case $j = 0$, $k \geq s+1$ with $k = s+2$

$$\begin{aligned} & (15 \cdot 2^{3s-6} - 7 \cdot 2^{2s-5}) + (2^s + 21 \cdot (5 \cdot 2^{3s-6} - 2^{2s-5})) \\ &+ \Gamma_{s+1} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+2) \cdot 2^{-(s+1)} + \Gamma_{s+2} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+2) \cdot 2^{-(s+2)} = 2^{5s+2m+1} + 2^{3s+3} - 2^{2s+1} \end{aligned}$$

$\Downarrow$

$$2 \cdot \Gamma_{s+1} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+2) + \Gamma_{s+2} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+2) = 2^{6s+2m+3} + 49 \cdot 2^{4s-1} - 9 \cdot 2^{3s-1} - 2^{2s+2}$$

hence we have (see above)

$$\Gamma_{s+1} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+2) = 105 \cdot 2^{4s-2} - 21 \cdot 2^{3s-2} - 3 \cdot 2^{2s}$$

$$\Gamma_{s+2} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+2) = 2^{6s+2m+3} - 7 \cdot 2^{4s+2} + 3 \cdot 2^{3s+1} + 2^{2s+1}$$

From (12.1) with $j = 1$ we get

$$\Gamma_{s+1} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k - \Gamma_{s+1} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+2) = (21 \cdot 2^{s-2} - 3 \cdot 2^{s-2}) \cdot (2^k - 2^{s+2})$$

$\Downarrow$

$$\Gamma_{s+1} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k = 105 \cdot 2^{4s-2} - 21 \cdot 2^{3s-2} - 3 \cdot 2^{2s} + (21 \cdot 2^{s-2} - 3 \cdot 2^{s-2}) \cdot (2^k - 2^{s+2})$$

$\Downarrow$

$$\Gamma_{s+1} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k = 9 \cdot 2^{k+s-1} + 21 \cdot (5 \cdot 2^{4s-2} - 2^{3s-2} - 2^{2s})$$

The case $j = 2$, $k \geq s+3$ and the case $j = 3$, $k = s+3$

From (6.1) with $k = s + 3$, $l = 0$ we obtain

$$\begin{aligned} \sum_{i=0}^{s+3} \Gamma_i \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) &= 2^{6s+2m+6} \\ \Downarrow \\ \sum_{i=0}^{s-1} \Gamma_i \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) &+ \Gamma_s \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) + \Gamma_{s+1} \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) + \Gamma_{s+2} \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) + \Gamma_{s+3} \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) = 2^{6s+2m+6} \end{aligned}$$

Using (6.39) and (12.7) in the cases $j = 0$, $k = s + 3$ and $j = 1$, $k = s + 3$ we have

$$\begin{aligned} (7 \cdot 2^{4s-6} - 3 \cdot 2^{3s-5}) + (3 \cdot 2^{2s} + 21 \cdot (5 \cdot 2^{4s-6} - 2^{3s-5})) + (15 \cdot 2^{2s} + 21 \cdot (5 \cdot 2^{4s-2} - 2^{3s-2})) \\ + \Gamma_{s+2} \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) + \Gamma_{s+3} \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) &= 2^{6s+2m+6} \\ \Downarrow \\ \Gamma_{s+2} \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) + \Gamma_{s+3} \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) &= 2^{6s+2m+6} - 7 \cdot 2^{4s+2} + 3 \cdot 2^{3s+1} - 9 \cdot 2^{2s+1} \end{aligned}$$

From (6.2) with $k = s + 3$, $l = 0$ we obtain

$$\begin{aligned} \sum_{i=0}^{s+3} \Gamma_i \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+2) \cdot 2^{-i} &= 2^{5s+2m+3} + 2^{3s+6} - 2^{2s+3} \\ \Downarrow \\ \sum_{i=0}^{s-1} \Gamma_i \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) \cdot 2^{-i} &+ \Gamma_s \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) \cdot 2^{-s} + \Gamma_{s+1} \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) \cdot 2^{-(s+1)} + \Gamma_{s+2} \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) \cdot 2^{-(s+2)} \\ &+ \Gamma_{s+3} \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) \cdot 2^{-(s+3)} = 2^{5s+2m+3} + 2^{3s+6} - 2^{2s+3} \end{aligned}$$

Using (6.40) and (12.7) in the cases $j = 0$, $k = s + 3$ and $j = 1$, $k = s + 3$ we have

$$\begin{aligned} (15 \cdot 2^{3s-6} - 7 \cdot 2^{2s-5}) + (3 \cdot 2^s + 21 \cdot (5 \cdot 2^{3s-6} - 2^{2s-5})) + (15 \cdot 2^{s-1} + 21 \cdot (5 \cdot 2^{3s-3} - 2^{2s-3})) \\ + \Gamma_{s+2} \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) \cdot 2^{-(s+2)} + \Gamma_{s+3} \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) \cdot 2^{-(s+3)} &= 2^{5s+2m+3} + 2^{3s+6} - 2^{2s+3} \\ \Downarrow \\ 2 \cdot \Gamma_{s+2} \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) + \Gamma_{s+3} \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) &= 2^{6s+2m+6} + 49 \cdot 2^{4s+3} - 9 \cdot 2^{3s+2} - 21 \cdot 2^{2s+2}, \end{aligned}$$

Hence we have (see above)

$$\begin{aligned} \Gamma_{s+2} \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) &= 105 \cdot 2^{4s+2} - 21 \cdot 2^{3s+1} - 33 \cdot 2^{2s+1} \\ \Gamma_{s+3} \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) &= 2^{6s+2m+6} - 7 \cdot 2^{4s+6} + 3 \cdot 2^{3s+4} + 3 \cdot 2^{2s+4} \end{aligned}$$

From (12.1) with $j = 2$ we get

$$\Gamma_{s+2} \begin{bmatrix} s \\ s+m \end{bmatrix} \times k - \Gamma_{s+2} \begin{bmatrix} s \\ s+m \end{bmatrix} \times (s+3) = (21 \cdot 2^{s+1} - 3 \cdot 2^s) \cdot (2^k - 2^{s+3})$$

$$\begin{aligned}
& \Downarrow \\
& \Gamma_{s+2}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k} = 105 \cdot 2^{4s+2} - 21 \cdot 2^{3s+1} - 33 \cdot 2^{2s+1} + 39 \cdot 2^s \cdot (2^k - 2^{s+3}) \\
& \Downarrow \\
& \Gamma_{s+2}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k} = 39 \cdot 2^{k+s} + 21 \cdot (5 \cdot 2^{4s+2} - 2^{3s+1} - 9 \cdot 2^{2s+1})
\end{aligned}$$

From the above results we now deduce that (H_{q-1}) holds for $q = 2, 3, 4, 5$.

Besides we get the formulas for $\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+j)}$ for $j = 0, 1, 2, 3, 4, 5$

Assume that (H_{q-1}) holds.

From (6.1) with $k = s + q + 1$, $l = 0$ we obtain

$$\begin{aligned}
& \sum_{i=0}^{s+q+1} \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} = 2^{6s+2m+3q} \\
& \Downarrow \\
& \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} + \Gamma_s^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} + \sum_{j=1}^{q-1} \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} \\
& + \Gamma_{s+q}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} + \Gamma_{s+q+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} = 2^{6s+2m+3q} \\
& \Downarrow \\
& (7 \cdot 2^{4s-6} - 3 \cdot 2^{3s-5}) + (2^{2s+q} - 2^{2s} + 21 \cdot (5 \cdot 2^{4s-6} - 2^{3s-5})) \\
& + \sum_{j=1}^{q-1} (21 \cdot 2^{j-1} - 3) \cdot 2^{2s+q+2j-3} + 21 \cdot (5 \cdot 2^{4s+4j-6} - 2^{3s+3j-5} - 5 \cdot 2^{2s+4j-6} + 2^{2s+3j-5}) \\
& + \Gamma_{s+q}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} + \Gamma_{s+q+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} = 2^{6s+2m+3q} \\
& \Downarrow \\
& \Gamma_{s+q}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} + \Gamma_{s+q+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} \\
& = 2^{6s+2m+3q} - 7 \cdot 2^{4s+4q-6} + 3 \cdot 2^{3s+3q-5} - 5 \cdot 2^{2s+4q-6} + 2^{2s+3q-5}
\end{aligned}$$

Using (6.2) with $k = s + q + 1$, $l = 0$ we obtain

$$\begin{aligned}
& \sum_{i=0}^{s+q+1} \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} \cdot 2^{-i} = 2^{5s+2m+2q-1} + 2^{3s+3q} - 2^{2s+2q-1} \\
& \Downarrow \\
& \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} \cdot 2^{-i} + \Gamma_s^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} \cdot 2^{-s} + \sum_{j=1}^{q-1} \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} \cdot 2^{-(s+j)} \\
& + \Gamma_{s+q}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} \cdot 2^{-(s+q)} + \Gamma_{s+q+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} \cdot 2^{-(s+q+1)} = 2^{5s+2m+2q-1} + 2^{3s+3q} - 2^{2s+2q-1} \\
& \Downarrow \\
& (15 \cdot 2^{3s-6} - 7 \cdot 2^{2s-5}) + (2^{s+q} - 2^s + 21 \cdot (5 \cdot 2^{3s-6} - 2^{2s-5}))
\end{aligned}$$

$$\begin{aligned}
& + \sum_{j=1}^{q-1} (21 \cdot 2^{j-1} - 3) \cdot 2^{s+q+j-3} + 21 \cdot (5 \cdot 2^{3s+3j-6} - 2^{2s+2j-5} - 5 \cdot 2^{s+3j-6} + 2^{s+2j-5}) \\
& + \Gamma_{s+q}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} \cdot 2^{-(s+q)} + \Gamma_{s+q+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} \cdot 2^{-(s+q+1)} = 2^{5s+2m+2q-1} + 2^{3s+3q} - 2^{2s+2q-1} \\
& \Downarrow \\
& 2 \cdot \Gamma_{s+q}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} + \Gamma_{s+q+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} \\
& = 2^{6s+2m+3q} + 49 \cdot 2^{4s+4q-5} - 9 \cdot 2^{3s+3q-4} + 5 \cdot 2^{2s+3q-4} - 13 \cdot 2^{2s+4q-5}
\end{aligned}$$

Hence we have (see above)

$$\begin{aligned}
\Gamma_{s+q}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} & = 105 \cdot 2^{4s+4q-6} - 21 \cdot 2^{3s+3q-5} - 21 \cdot 2^{2s+4q-6} + 9 \cdot 2^{2s+3q-5} \\
\Gamma_{s+q+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} & = 2^{6s+2m+3q} - 7 \cdot 2^{4s+4q-2} + 3 \cdot 2^{3s+3q-2} + 2^{2s+4q-2} - 2^{2s+3q-2}
\end{aligned}$$

From (12.1) with $j = q$ we get

$$\begin{aligned}
\Gamma_{s+q}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k} - \Gamma_{s+q}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+q+1)} & = (21 \cdot 2^{s+3q-5} - 3 \cdot 2^{s+2q-4}) \cdot (2^k - 2^{s+q+1}) \\
\Downarrow \\
\Gamma_{s+q}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k} & = 105 \cdot 2^{4s+4q-6} - 21 \cdot 2^{3s+3q-5} - 21 \cdot 2^{2s+4q-6} + 9 \cdot 2^{2s+3q-5} \\
& + (21 \cdot 2^{s+3q-5} - 3 \cdot 2^{s+2q-4}) \cdot (2^k - 2^{s+q+1}) \\
\Downarrow \\
\Gamma_{s+q}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k} & = (21 \cdot 2^{q-1} - 3) \cdot 2^{k+s+2q-4} + 21 \cdot (5 \cdot 2^{4s+4q-6} - 2^{3s+3q-5} - 5 \cdot 2^{2s+4q-6} + 2^{2s+3q-5})
\end{aligned}$$

The case $j = m$, $k \geq s + m + 1$

We obtain from the proof by induction with $q = m$ (see above)

$$\begin{aligned}
\Gamma_{s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+m+1)} & = 105 \cdot 2^{4s+4m-6} - 21 \cdot 2^{3s+3m-5} - 21 \cdot 2^{2s+4m-6} + 9 \cdot 2^{2s+3m-5} \\
\Gamma_{s+m+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+m+1)} & = 2^{6s+5m} - 7 \cdot 2^{4s+4m-2} + 3 \cdot 2^{3s+3m-2} + 2^{2s+4m-2} - 2^{2s+3m-2}
\end{aligned}$$

From (12.1) with $j = m$ we get

$$\begin{aligned}
\Gamma_{s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k} - \Gamma_{s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (s+m+1)} & = (21 \cdot 2^{s+3m-5} + 45 \cdot 2^{s+2m-4}) \cdot (2^k - 2^{s+m+1}) \\
\Downarrow \\
\Gamma_{s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k} & = 105 \cdot 2^{4s+4m-6} - 21 \cdot 2^{3s+3m-5} - 21 \cdot 2^{2s+4m-6} + 9 \cdot 2^{2s+3m-5} \\
& + (21 \cdot 2^{s+3m-5} + 45 \cdot 2^{s+2m-4}) \cdot (2^k - 2^{s+m+1}) \\
\Downarrow \\
\Gamma_{s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k} & = (21 \cdot 2^{s+3m-5} + 45 \cdot 2^{s+2m-4}) \cdot (2^k - 2^{s+m+1})
\end{aligned}$$

$$+ 105 \cdot 2^{4s+4m-6} - 21 \cdot 2^{3s+3m-5} - 21 \cdot 2^{2s+4m-6} + 9 \cdot 2^{2s+3m-5}$$

□

12.5. **Computation of** $\Gamma_{s+m+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k}$, for $1 \leq j \leq s+2$, $k \geq s+m+j$, $m \geq 2$, $j = s+3$, $k = 2s+m+3$.

Lemma 12.4. We have for $m \geq 2$, $k \geq s+m$:

$$(12.8) \quad \sum_{j=1}^{m-1} \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k} \\ = 3 \cdot 2^{k+s+3m-5} - 3 \cdot 2^{3s+3m-5} + 3 \cdot 2^{2s+3m-5} - 2^{k+s-1} \\ + 3 \cdot 2^{3s-2} + 2^{2s} + 7 \cdot 2^{4s+4m-6} - 7 \cdot 2^{2s+4m-6} - 7 \cdot 2^{4s-2} - 2^{k+s+2m-4}$$

$$(12.9) \quad \sum_{j=1}^{m-1} \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k} \cdot 2^{-(s+j)} \\ = 7 \cdot 2^{k+2m-5} - 7 \cdot 2^{2s+2m-5} + 7 \cdot 2^{s+2m-5} - 2^{k-1} \\ + 7 \cdot 2^{2s-3} + 2^s + 15 \cdot 2^{3s+3m-6} - 15 \cdot 2^{s+3m-6} - 15 \cdot 2^{3s-3} - 3 \cdot 2^{k+m-4}$$

Proof. **Proof of (12.8)**

From Lemma 12.3 in the case $1 \leq j \leq m-1$, $k \geq s+j+1$ we obtain :

$$\begin{aligned} & \sum_{j=1}^{m-1} \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k} \\ &= (21 \cdot 2^{k+s-5} - 21 \cdot 2^{3s-5} + 21 \cdot 2^{2s-5}) \cdot \sum_{j=1}^{m-1} 2^{3j} \\ &+ (105 \cdot 2^{4s-6} - 105 \cdot 2^{2s-6}) \cdot \sum_{j=1}^{m-1} 2^{4j} - (3 \cdot 2^{k+s-4}) \cdot \sum_{j=1}^{m-1} 2^{2j} \\ &= (3 \cdot 2^{k+s-5} - 3 \cdot 2^{3s-5} + 3 \cdot 2^{2s-5}) \cdot (2^{3m} - 2^3) \\ &+ (7 \cdot 2^{4s-6} - 7 \cdot 2^{2s-6}) \cdot (2^{4m} - 2^4) - (2^{k+s-4}) \cdot (2^{2m} - 2^2) \end{aligned}$$

Proof of (12.9)

Similar to the proof of (12.8) :

$$\begin{aligned} & \sum_{j=1}^{m-1} \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k} \cdot 2^{-(s+j)} \\ &= (21 \cdot 2^{k-5} - 21 \cdot 2^{2s-5} + 21 \cdot 2^{s-5}) \cdot \sum_{j=1}^{m-1} 2^{2j} \\ &+ (105 \cdot 2^{3s-6} - 105 \cdot 2^{s-6}) \cdot \sum_{j=1}^{m-1} 2^{3j} - (3 \cdot 2^{k-4}) \cdot \sum_{j=1}^{m-1} 2^j \\ &= (7 \cdot 2^{k-5} - 7 \cdot 2^{2s-5} + 7 \cdot 2^{s-5}) \cdot (2^{2m} - 2^2) \end{aligned}$$

$$+ (15 \cdot 2^{3s-6} - 15 \cdot 2^{s-6}) \cdot (2^{3m} - 2^3) - (3 \cdot 2^{k-4}) \cdot (2^m - 2)$$

□

Lemma 12.5. *We have for $m \geq 2$, $1 \leq j \leq s-1$:*

$$(12.10) \quad \Gamma_{s+m+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} \times k = \begin{cases} 105 \cdot 2^{4s+4m-2} - 21 \cdot 2^{3s+3m-2} - 21 \cdot 2^{2s+4m-2} - 63 \cdot 2^{2s+3m-2} & \text{if } j=1, k=s+m+2 \\ 2^{6s+5m+3} - 7 \cdot 2^{4s+4m+2} + 3 \cdot 2^{3s+3m+1} + 2^{2s+4m+2} + 5 \cdot 2^{2s+3m+1} & \text{if } j=2, k=s+m+2 \\ 21 \cdot [2^{k+s+3m-2} + 13 \cdot 2^{k+s+2m-2} + 5 \cdot 2^{4s+4m-2} & \\ - 2^{3s+3m-2} - 5 \cdot 2^{2s+4m-2} - 55 \cdot 2^{2s+3m-2}] & \text{if } j=1, k \geq s+m+2 \\ \\ 105 \cdot 2^{4s+4m+2} - 21 \cdot 2^{3s+3m+1} - 21 \cdot 2^{2s+4m+2} - 441 \cdot 2^{2s+3m+1} & \text{if } j=2, k=s+m+3 \\ 2^{6s+5m+6} - 7 \cdot 2^{4s+4m+6} + 3 \cdot 2^{3s+3m+4} + 2^{2s+4m+6} + 19 \cdot 2^{2s+3m+5} & \text{if } j=3, k=s+m+3 \\ 21 \cdot [2^{k+s+3m+1} + 61 \cdot 2^{k+s+2m} + 5 \cdot 2^{4s+4m+2} & \\ - 2^{3s+3m+1} - 5 \cdot 2^{2s+4m+2} - 265 \cdot 2^{2s+3m+1}] & \text{if } j=2, k \geq s+m+3 \\ \\ 105 \cdot 2^{4s+4m+6} - 21 \cdot 2^{3s+3m+4} - 21 \cdot 2^{2s+4m+6} - 1071 \cdot 2^{2s+3m+5} & \text{if } j=3, k=s+m+4 \\ 2^{6s+5m+9} - 7 \cdot 2^{4s+4m+10} + 3 \cdot 2^{3s+3m+7} + 2^{2s+4m+10} + 47 \cdot 2^{2s+3m+9} & \text{if } j=4, k=s+m+4 \\ 21 \cdot [2^{k+s+3m+4} + 131 \cdot 2^{k+s+2m+3} + 5 \cdot 2^{4s+4m+6} & \\ - 2^{3s+3m+4} - 5 \cdot 2^{2s+4m+6} - 575 \cdot 2^{2s+3m+5}] & \text{if } j=3, k \geq s+m+4 \\ \\ 105 \cdot 2^{4s+4m+10} - 21 \cdot 2^{3s+3m+7} - 21 \cdot 2^{2s+4m+10} - 2331 \cdot 2^{2s+3m+9} & \text{if } j=4, k=s+m+5 \\ 2^{6s+5m+12} - 7 \cdot 2^{4s+4m+14} + 3 \cdot 2^{3s+3m+10} + 2^{2s+4m+14} + 103 \cdot 2^{2s+3m+13} & \text{if } j=5, k=s+m+5 \\ 21 \cdot [2^{k+s+3m+7} + 271 \cdot 2^{k+s+2m+6} + 5 \cdot 2^{4s+4m+10} & \\ - 2^{3s+3m+7} - 5 \cdot 2^{2s+4m+10} - 1195 \cdot 2^{2s+3m+9}] & \text{if } j=4, k \geq s+m+5 \\ \\ 105 \cdot 2^{4s+4m+4j-6} - 21 \cdot 2^{3s+3m+3j-5} - 21 \cdot 2^{2s+4m+4j-6} & \\ - 315 \cdot 2^{2s+3m+5j-8} + 189 \cdot 2^{2s+3m+4j-7} & \text{if } k=s+m+j+1 \\ 2^{6s+5m+3j-3} - 7 \cdot 2^{4s+4m+4j-6} + 3 \cdot 2^{3s+3m+3j-5} & \\ + 2^{2s+4m+4j-6} + 7 \cdot 2^{2s+3m+5j-8} - 9 \cdot 2^{2s+3m+4j-7} & \text{if } k=s+m+j \\ 21 \cdot [2^{k+s+3m+3j-5} + 35 \cdot 2^{k+s+2m+4j-7} - 9 \cdot 2^{k+s+2m+3j-6} + 5 \cdot 2^{4s+4m+4j-6} & \\ - 2^{3s+3m+3j-5} - 5 \cdot 2^{2s+4m+4j-6} - 155 \cdot 2^{2s+3m+5j-8} + 45 \cdot 2^{2s+3m+4j-7}] & \text{if } k \geq s+m+j+1 \end{cases}$$

Let (H_{q-1}) denote the following statement :

$$\Gamma_{s+m+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} \times k = 21 \cdot [2^{k+s+3m+3j-5} + 35 \cdot 2^{k+s+2m+4j-7} - 9 \cdot 2^{k+s+2m+3j-6} + 5 \cdot 2^{4s+4m+4j-6} \\ - 2^{3s+3m+3j-5} - 5 \cdot 2^{2s+4m+4j-6} - 155 \cdot 2^{2s+3m+5j-8} + 45 \cdot 2^{2s+3m+4j-7}]$$

for $1 \leq j \leq q-1 \leq s-2$, $k \geq s+m+j+1$

Proof. Similar to the proof of Lemma 12.3.

We start by computing successively $\Gamma_{s+m+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix}} \times k$ for $j=1, 2, 3, 4$ and $k \geq s+m+j+1$.

The statement (H_{q-1}) is now obtained by observing their structures.

□

Lemma 12.6. *We have for $m \geq 2$, $k \geq 2s+m$:*

$$\begin{aligned}
(12.11) \quad & \sum_{j=1}^{s-1} \Gamma_{s+m+j}^{\left[\begin{smallmatrix} s \\ s+m \\ s+m \end{smallmatrix} \right] \times k} \\
&= 3 \cdot 2^{k+4s+3m-5} - 27 \cdot 2^{k+4s+2m-6} - 3 \cdot 2^{k+s+3m-2} - 11 \cdot 2^{k+s+2m-2} \\
&+ 49 \cdot 2^{k+5s+2m-7} + 51 \cdot 2^{6s+3m-7} + 3 \cdot 2^{3s+3m-2} + 7 \cdot 2^{8s+4m-6} - 7 \cdot 2^{6s+4m-6} \\
&- 7 \cdot 2^{4s+4m-2} + 7 \cdot 2^{2s+4m-2} - 105 \cdot 2^{7s+3m-8} + 21 \cdot 2^{2s+3m-2}
\end{aligned}$$

Setting $k = 2s + m$ in (12.11) we get

$$\begin{aligned}
(12.12) \quad & \sum_{j=1}^{s-1} \Gamma_{s+m+j}^{\left[\begin{smallmatrix} s \\ s+m \\ s+m \end{smallmatrix} \right] \times (2s+m)} \\
&= 7 \cdot 2^{8s+4m-6} + 7 \cdot 2^{2s+4m-2} + 21 \cdot 2^{2s+3m-2} - 7 \cdot 2^{4s+4m-2} \\
&- 7 \cdot 2^{7s+3m-8} - 3 \cdot 2^{3s+4m-2} - 3 \cdot 2^{6s+3m-7} - 2^{6s+4m-6} - 2^{3s+3m+1}
\end{aligned}$$

$$\begin{aligned}
(12.13) \quad & \sum_{j=1}^{s-1} \Gamma_{s+m+j}^{\left[\begin{smallmatrix} s \\ s+m \\ s+m \end{smallmatrix} \right] \times k} \cdot 2^{-(s+m+j)} \\
&= 7 \cdot 2^{k+2s+2m-5} - 63 \cdot 2^{k+2s+m-6} - 7 \cdot 2^{k+2m-3} - 21 \cdot 2^{k+m-3} \\
&+ 107 \cdot 2^{4s+2m-7} + 7 \cdot 2^{2s+2m-3} + 105 \cdot 2^{k+3s+m-7} + 15 \cdot 2^{6s+3m-6} \\
&- 15 \cdot 2^{4s+3m-6} - 15 \cdot 2^{3s+3m-3} + 15 \cdot 2^{s+3m-3} - 217 \cdot 2^{5s+2m-8} + 41 \cdot 2^{s+2m-3}
\end{aligned}$$

Setting $k = 2s + m$ in (12.13) we get

$$\begin{aligned}
(12.14) \quad & \sum_{j=1}^{s-1} \Gamma_{s+m+j}^{\left[\begin{smallmatrix} s \\ s+m \\ s+m \end{smallmatrix} \right] \times (2s+m)} \cdot 2^{-(s+m+j)} \\
&= -2^{4s+3m-6} - 19 \cdot 2^{4s+2m-7} - 7 \cdot 2^{2s+3m-3} - 7 \cdot 2^{2s+2m-2} \\
&- 7 \cdot 2^{5s+2m-8} - 15 \cdot 2^{3s+3m-3} + 15 \cdot 2^{6s+3m-6} + 15 \cdot 2^{s+3m-3} + 41 \cdot 2^{s+2m-3}
\end{aligned}$$

Proof. The assertions follow from Lemma 12.5

□

Lemma 12.7. *We have for $m \geq 2$, $0 \leq j \leq m$:*

$$\begin{aligned}
 & \Gamma_{2s+m+j} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k \\
 (12.15) \quad & = \begin{cases} 2^{9s+5m-3} - 7 \cdot 2^{8s+4m-6} + 7 \cdot 2^{7s+3m-8} + 3 \cdot 2^{6s+3m-7} + 2^{6s+4m-6} & \text{if } j = 0, k = 2s + m \\ 21 \cdot [5 \cdot 2^{8s+4m-6} + 5 \cdot 2^{6s+3m-7} - 2^{6s+4m-6} - 15 \cdot 2^{7s+3m-8}] & \text{if } j = 0, k = 2s + m + 1 \\ 2^{9s+5m} - 7 \cdot 2^{8s+4m-2} + 7 \cdot 2^{7s+3m-3} - 3 \cdot 2^{6s+3m-3} + 2^{6s+4m-2} & \text{if } j = 1, k = 2s + m + 1 \\ 3 \cdot 2^{2k+2s+m-2} + 21 \cdot 2^{k+4s+3m-5} + 735 \cdot 2^{k+5s+2m-7} - 477 \cdot 2^{k+4s+2m-6} & \\ + 105 \cdot 2^{8s+4m-6} - 105 \cdot 2^{6s+4m-6} - 3255 \cdot 2^{7s+3m-8} + 1629 \cdot 2^{6s+3m-7} & \text{if } j = 0, k \geq 2s + m + 1 \\ 21 \cdot [5 \cdot 2^{8s+4m-2} + 7 \cdot 2^{6s+3m-3} - 2^{6s+4m-2} - 15 \cdot 2^{7s+3m-3}] & \text{if } j = 1, k = 2s + m + 2 \\ 2^{9s+5m+3} - 7 \cdot 2^{8s+4m+2} + 7 \cdot 2^{7s+3m+2} - 3 \cdot 2^{6s+3m+2} + 2^{6s+4m+2} & \text{if } j = 2, k = 2s + m + 2 \\ 21 \cdot 2^{2k+2s+m-2} + 21 \cdot 2^{k+4s+3m-2} + 735 \cdot 2^{k+5s+2m-3} - 945 \cdot 2^{k+4s+2m-3} & \\ + 105 \cdot 2^{8s+4m-2} - 105 \cdot 2^{6s+4m-2} - 3255 \cdot 2^{7s+3m-3} + 3255 \cdot 2^{6s+3m-3} & \text{if } j = 1, k \geq 2s + m + 2 \\ 21 \cdot [5 \cdot 2^{8s+4m+2} + 7 \cdot 2^{6s+3m+2} - 2^{6s+4m+2} - 15 \cdot 2^{7s+3m+2}] & \text{if } j = 2, k = 2s + m + 3 \\ 2^{9s+6m+6} - 7 \cdot 2^{8s+4m+6} + 7 \cdot 2^{7s+3m+7} - 3 \cdot 2^{6s+3m+7} + 2^{6s+4m+6} & \text{if } j = 3, k = 2s + m + 3 \\ 21 \cdot 2^{2k+2s+m+1} + 21 \cdot 2^{k+4s+3m+1} + 735 \cdot 2^{k+5s+2m+1} - 945 \cdot 2^{k+4s+2m+1} & \\ + 105 \cdot 2^{8s+4m+2} - 105 \cdot 2^{6s+4m+2} - 3255 \cdot 2^{7s+3m+2} + 3255 \cdot 2^{6s+3m+2} & \text{if } j = 2, k \geq 2s + m + 3 \end{cases}
 \end{aligned}$$

Proof. **The case $j = 0$, $k = 2s + m$**

From (6.1) with $k = 2s + m$, $l = 0$ we obtain using (12.7),(12.8) and (12.12)

$$\begin{aligned}
 & \sum_{i=0}^{2s+m} \Gamma_i \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m) = 2^{9s+5m-3} \\
 & \Downarrow \\
 & \sum_{i=0}^{s-1} \Gamma_i \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m) + \Gamma_s \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m) + \sum_{j=1}^{m-1} \Gamma_{s+j} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m) \\
 & + \Gamma_{s+m} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m) + \sum_{j=1}^{s-1} \Gamma_{s+m+j} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m) + \Gamma_{2s+m} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m) = 2^{9s+6m-3} \\
 & \Downarrow \\
 & (7 \cdot 2^{4s-6} - 3 \cdot 2^{3s-5}) + (2^{3s+m-1} - 2^{2s} + 21 \cdot (5 \cdot 2^{4s-6} - 2^{3s-5})) \\
 & + (3 \cdot 2^{3s+4m-2} - 3 \cdot 2^{3s+3m-5} + 3 \cdot 2^{2s+3m-5} - 2^{3s+m-1} \\
 & + 3 \cdot 2^{3s-2} + 2^{2s} + 7 \cdot 2^{4s+4m-6} - 7 \cdot 2^{2s+4m-6} - 7 \cdot 2^{4s-2} - 2^{3s+3m-4}) \\
 & + (21 \cdot 2^{3s+4m-5} + 45 \cdot 2^{3s+3m-4} - 21 \cdot 2^{2s+4m-4} - 45 \cdot 2^{2s+3m-3} \\
 & + 105 \cdot 2^{4s+4m-6} - 21 \cdot 2^{3s+3m-5} - 21 \cdot 2^{2s+4m-6} + 9 \cdot 2^{2s+3m-5}) \\
 & + (7 \cdot 2^{8s+4m-6} + 7 \cdot 2^{2s+4m-2} + 21 \cdot 2^{2s+3m-2} - 7 \cdot 2^{4s+4m-2} \\
 & - 7 \cdot 2^{7s+3m-8} - 3 \cdot 2^{3s+4m-2} - 3 \cdot 2^{6s+3m-7} - 2^{6s+4m-6} - 2^{3s+3m+1}) \\
 & + \Gamma_{2s+m} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m) = 2^{9s+5m-3} \\
 & \Downarrow \\
 & \Gamma_{2s+m} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m) = 2^{9s+5m-3} - 7 \cdot 2^{8s+4m-6} + 3 \cdot 2^{6s+3m-7} + 7 \cdot 2^{7s+3m-8} + 2^{6s+4m-6}
 \end{aligned}$$

The case $j = 0$, $k = 2s + m + 1$ and the case $j = 1$, $k = 2s + m + 1$.

From (6.1) with $k = 2s + m + 1$, $l = 0$ we obtain using (12.7) and (12.8),(12.11)

$$\begin{aligned}
& \sum_{i=0}^{2s+m+1} \Gamma_i \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1) = 2^{9s+5m} \\
& \Updownarrow \\
& \sum_{i=0}^{s-1} \Gamma_i \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1) + \Gamma_s \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1) + \sum_{j=1}^{m-1} \Gamma_{s+j} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1) \\
& + \Gamma_{s+m} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1) + \sum_{j=1}^{s-1} \Gamma_{s+m+j} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1) + \Gamma_{2s+m} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1) + \Gamma_{2s+m+1} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1) = 2^{9s+5m} \\
& \Updownarrow \\
& (7 \cdot 2^{4s-6} - 3 \cdot 2^{3s-5}) + (2^{3s+m} - 2^{2s} + 21 \cdot (5 \cdot 2^{4s-6} - 2^{3s-5})) \\
& + (3 \cdot 2^{3s+4m-4} - 3 \cdot 2^{3s+3m-5} + 3 \cdot 2^{2s+3m-5} - 2^{3s+m} \\
& + 3 \cdot 2^{3s-2} + 2^{2s} + 7 \cdot 2^{4s+4m-6} - 7 \cdot 2^{2s+4m-6} - 7 \cdot 2^{4s-2} - 2^{3s+3m-3}) \\
& + (21 \cdot 2^{3s+4m-4} + 45 \cdot 2^{3s+3m-3} - 21 \cdot 2^{2s+4m-4} - 45 \cdot 2^{2s+3m-3} \\
& + 105 \cdot 2^{4s+4m-6} - 21 \cdot 2^{3s+3m-5} - 21 \cdot 2^{2s+4m-6} + 9 \cdot 2^{2s+3m-5}) \\
& + (3 \cdot 2^{6s+4m-4} - 27 \cdot 2^{6s+3m-5} - 3 \cdot 2^{3s+4m-1} - 11 \cdot 2^{3s+3m-1} \\
& + 49 \cdot 2^{7s+3m-6} + 51 \cdot 2^{6s+3m-7} + 3 \cdot 2^{3s+3m-2} + 7 \cdot 2^{8s+4m-6} - 7 \cdot 2^{6s+4m-6} \\
& - 7 \cdot 2^{4s+4m-2} + 7 \cdot 2^{2s+4m-2} - 105 \cdot 2^{7s+3m-8} + 21 \cdot 2^{2s+3m-2}) \\
& + \Gamma_{2s+m} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1) + \Gamma_{2s+m+1} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1) = 2^{9s+5m} \\
& \Updownarrow \\
& \Gamma_{2s+m} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1) + \Gamma_{2s+m+1} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1) \\
& = 2^{9s+5m} - 7 \cdot 2^{8s+4m-6} - 5 \cdot 2^{6s+4m-6} + 57 \cdot 2^{6s+3m-7} - 91 \cdot 2^{7s+3m-8}
\end{aligned}$$

From (6.2) with $k = 2s + m + 1$, $l = 0$ we obtain using (12.7) and (12.9),(12.13)

$$\begin{aligned}
& \sum_{i=0}^{2s+m+1} \Gamma_i \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1) \cdot 2^{-i} = 2^{7s+4m-1} + 2^{6s+3m} - 2^{4s+2m-1} \\
& \Updownarrow \\
& \sum_{i=0}^{s-1} \Gamma_i \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1) \cdot 2^{-i} + \Gamma_s \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1) \cdot 2^{-s} + \sum_{j=1}^{m-1} \Gamma_{s+j} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1) \cdot 2^{-(s+j)} \\
& + \Gamma_{s+m} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1) \cdot 2^{-(s+m)} + \sum_{j=1}^{s-1} \Gamma_{s+m+j} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1) \cdot 2^{-(s+m+j)} \\
& + \Gamma_{2s+m} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1) \cdot 2^{-(2s+m)} + \Gamma_{2s+m+1} \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1) \cdot 2^{-(2s+m+1)} = 2^{7s+4m-1} + 2^{6s+3m} - 2^{4s+2m-1} \\
& \Updownarrow \\
& (15 \cdot 2^{3s-6} - 7 \cdot 2^{2s-5}) + (2^{2s+m} - 2^s + 21 \cdot (5 \cdot 2^{3s-6} - 2^{2s-5})) \\
& = (7 \cdot 2^{2s+3m-4} - 7 \cdot 2^{2s+2m-5} + 7 \cdot 2^{s+2m-5} - 2^{2s+m}
\end{aligned}$$

$$\begin{aligned}
& + 7 \cdot 2^{2s-3} + 2^s + 15 \cdot 2^{3s+3m-6} - 15 \cdot 2^{s+3m-6} - 15 \cdot 2^{3s-3} - 3 \cdot 2^{2s+2m-3}) \\
& + (21 \cdot 2^{2s+3m-4} + 45 \cdot 2^{2s+2m-3} - 21 \cdot 2^{s+3m-4} - 45 \cdot 2^{s+2m-3} \\
& + 105 \cdot 2^{3s+3m-6} - 21 \cdot 2^{2s+2m-5} - 21 \cdot 2^{s+3m-6} + 9 \cdot 2^{s+2m-5}) \\
& + (7 \cdot 2^{4s+3m-4} - 63 \cdot 2^{4s+2m-5} - 7 \cdot 2^{2s+3m-2} - 21 \cdot 2^{2s+2m-2} \\
& + 107 \cdot 2^{4s+2m-7} + 7 \cdot 2^{2s+2m-3} + 105 \cdot 2^{5s+2m-6} + 15 \cdot 2^{6s+3m-6} \\
& - 15 \cdot 2^{4s+3m-6} - 15 \cdot 2^{3s+3m-3} + 15 \cdot 2^{s+3m-3} - 217 \cdot 2^{5s+2m-8} + 41 \cdot 2^{s+2m-3}) \\
& + \Gamma_{2s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1)} \cdot 2^{-(2s+m)} + \Gamma_{2s+m+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1)} \cdot 2^{-(2s+m+1)} = 2^{7s+4m-1} + 2^{6s+3m} - 2^{4s+2m-1} \\
& \Downarrow \\
& 2 \cdot \Gamma_{2s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1)} + \Gamma_{2s+m+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1)} \\
& = 2^{9s+5m} + 49 \cdot 2^{8s+4m-5} + 81 \cdot 2^{6s+3m-6} - 13 \cdot 2^{6s+4m-5} - 203 \cdot 2^{7s+3m-7}
\end{aligned}$$

The case $j = 0$, $k \geq 2s + m + 1$

From (12.2) with $j = s$ we get

$$\begin{aligned}
& \Gamma_{2s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k} - \Gamma_{2s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+1)} \\
& = 3 \cdot 2^{2s+m-2} \cdot (2^{2k} - 2^{4s+2m+2}) + (2^k - 2^{2s+m+1}) \cdot (21 \cdot 2^{4s+3m-5} + 735 \cdot 2^{5s+2m-7} - 477 \cdot 2^{4s+2m-6}) \\
& \Downarrow \\
& \Gamma_{2s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k} = 21 \cdot [5 \cdot 2^{8s+4m-6} + 5 \cdot 2^{6s+3m-7} - 2^{6s+4m-6} - 15 \cdot 2^{7s+3m-8}] \\
& + 3 \cdot 2^{2s+m-2} \cdot (2^{2k} - 2^{4s+2m+2}) + (2^k - 2^{2s+m+1}) \cdot (21 \cdot 2^{4s+3m-5} + 735 \cdot 2^{5s+2m-7} - 477 \cdot 2^{4s+2m-6}) \\
& \Downarrow \\
& \Gamma_{2s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k} = 3 \cdot 2^{2k+2s+m-2} + 21 \cdot 2^{k+4s+3m-5} + 735 \cdot 2^{k+5s+2m-7} - 477 \cdot 2^{k+4s+2m-6} \\
& + 105 \cdot 2^{8s+4m-6} - 105 \cdot 2^{6s+4m-6} - 3255 \cdot 2^{7s+3m-8} + 1629 \cdot 2^{6s+3m-7}
\end{aligned}$$

The case $j = 1$, $k = 2s + m + 2$ and the case $j = 2$, $k = 2s + m + 1$

From (6.1) with $k = 2s + m + 2$, $l = 0$ we obtain using (12.7) and (12.8),(12.11)

$$\begin{aligned}
& \sum_{i=0}^{2s+m+2} \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} = 2^{9s+5m+3} \\
& \Downarrow \\
& \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} + \Gamma_s^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} + \sum_{j=1}^{m-1} \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} \\
& + \Gamma_{s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} + \sum_{j=1}^{s-1} \Gamma_{s+m+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} + \Gamma_{2s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} \\
& + \Gamma_{2s+m+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} + \Gamma_{2s+m+2}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} = 2^{9s+5m+3} \\
& \Downarrow \\
& (7 \cdot 2^{4s-6} - 3 \cdot 2^{3s-5}) + (2^{3s+m} - 2^{2s} + 21 \cdot (5 \cdot 2^{4s-6} - 2^{3s-5}))
\end{aligned}$$

$$\begin{aligned}
& + (3 \cdot 2^{3s+4m-4} - 3 \cdot 2^{3s+3m-5} + 3 \cdot 2^{2s+3m-5} - 2^{3s+m} \\
& + 3 \cdot 2^{3s-2} + 2^{2s} + 7 \cdot 2^{4s+4m-6} - 7 \cdot 2^{2s+4m-6} - 7 \cdot 2^{4s-2} - 2^{3s+3m-3}) \\
& + (21 \cdot 2^{3s+4m-3} + 45 \cdot 2^{3s+3m-2} - 21 \cdot 2^{2s+4m-4} - 45 \cdot 2^{2s+3m-3} \\
& + 105 \cdot 2^{4s+4m-6} - 21 \cdot 2^{3s+3m-5} - 21 \cdot 2^{2s+4m-6} + 9 \cdot 2^{2s+3m-5}) \\
& (3 \cdot 2^{6s+4m-3} - 27 \cdot 2^{6s+3m-4} - 3 \cdot 2^{3s+4m} - 11 \cdot 2^{3s+3m} \\
& + 49 \cdot 2^{7s+3m-5} + 51 \cdot 2^{6s+3m-7} + 3 \cdot 2^{3s+3m-2} + 7 \cdot 2^{8s+4m-6} - 7 \cdot 2^{6s+4m-6} \\
& - 7 \cdot 2^{4s+4m-2} + 7 \cdot 2^{2s+4m-2} - 105 \cdot 2^{7s+3m-8} + 21 \cdot 2^{2s+3m-2}) \\
& + 3 \cdot 2^{6s+3m+2} + 21 \cdot 2^{6s+4m-3} + 735 \cdot 2^{7s+3m-5} - 477 \cdot 2^{6s+3m-4} \\
& + 105 \cdot 2^{8s+4m-6} - 105 \cdot 2^{6s+4m-6} - 3255 \cdot 2^{7s+3m-8} + 1629 \cdot 2^{6s+3m-7} \\
& + \Gamma_{2s+m+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} + \Gamma_{2s+m+2}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} = 2^{9s+5m+3} \\
& \Downarrow \\
& \Gamma_{2s+m+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} + \Gamma_{2s+m+2}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} \\
& = 2^{9s+5m+3} - 7 \cdot 2^{8s+4m-2} - 5 \cdot 2^{6s+4m-2} + 51 \cdot 2^{6s+3m-3} - 91 \cdot 2^{7s+3m-3}
\end{aligned}$$

From (6.2) with $k = 2s + m + 2$, $l = 0$ we obtain using (12.7) and (12.9),(12.13)

$$\begin{aligned}
& \sum_{i=0}^{2s+m+2} \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} \cdot 2^{-i} = 2^{7s+4m+1} + 2^{6s+3m+3} - 2^{4s+2m+1} \\
& \Downarrow \\
& \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} \cdot 2^{-i} + \Gamma_s^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} \cdot 2^{-s} + \sum_{j=1}^{m-1} \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} \cdot 2^{-(s+j)} \\
& + \Gamma_{s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} \cdot 2^{-(s+m)} + \sum_{j=1}^{s-1} \Gamma_{s+m+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} \cdot 2^{-(s+m+j)} + \Gamma_{2s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} \cdot 2^{-(2s+m)} \\
& + \Gamma_{2s+m+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} \cdot 2^{-(2s+m+1)} + \Gamma_{2s+m+2}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} \cdot 2^{-(2s+m+2)} = 2^{7s+4m-1} + 2^{6s+3m} - 2^{4s+2m-1} \\
& \Downarrow \\
& (15 \cdot 2^{3s-6} - 7 \cdot 2^{2s-5}) + (2^{2s+m+1} - 2^s + 21 \cdot (5 \cdot 2^{3s-6} - 2^{2s-5})) \\
& = (7 \cdot 2^{2s+3m-3} - 7 \cdot 2^{2s+2m-5} + 7 \cdot 2^{s+2m-5} - 2^{2s+m+1} \\
& + 7 \cdot 2^{2s-3} + 2^s + 15 \cdot 2^{3s+3m-6} - 15 \cdot 2^{s+3m-6} - 15 \cdot 2^{3s-3} - 3 \cdot 2^{2s+2m-2}) \\
& + (21 \cdot 2^{2s+3m-3} + 45 \cdot 2^{2s+2m-2} - 21 \cdot 2^{s+3m-4} - 45 \cdot 2^{s+2m-3} \\
& + 105 \cdot 2^{3s+3m-6} - 21 \cdot 2^{2s+2m-5} - 21 \cdot 2^{s+3m-6} + 9 \cdot 2^{s+2m-5}) \\
& + (7 \cdot 2^{4s+3m-3} - 63 \cdot 2^{4s+2m-4} - 7 \cdot 2^{2s+3m-1} - 21 \cdot 2^{2s+2m-1} \\
& + 107 \cdot 2^{4s+2m-7} + 7 \cdot 2^{2s+2m-3} + 105 \cdot 2^{5s+2m-5} + 15 \cdot 2^{6s+3m-6} \\
& - 15 \cdot 2^{4s+3m-6} - 15 \cdot 2^{3s+3m-3} + 15 \cdot 2^{s+3m-3} - 217 \cdot 2^{5s+2m-8} + 41 \cdot 2^{s+2m-3}) \\
& + 3 \cdot 2^{4s+2m+2} + 21 \cdot 2^{4s+3m-3} + 735 \cdot 2^{5s+2m-5} - 477 \cdot 2^{4s+2m-4} \\
& + 105 \cdot 2^{6s+3m-6} - 105 \cdot 2^{4s+3m-6} - 3255 \cdot 2^{5s+2m-8} + 1629 \cdot 2^{4s+2m-7}
\end{aligned}$$

$$\begin{aligned}
& + \Gamma_{2s+m+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} \cdot 2^{-(2s+m+1)} + \Gamma_{2s+m+2}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} \cdot 2^{-(2s+m+2)} = 2^{7s+4m+1} + 2^{6s+3m+3} - 2^{4s+2m+1} \\
& \Downarrow \\
& 2 \cdot \Gamma_{2s+m+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} + \Gamma_{2s+m+2}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} \\
& = 2^{9s+5m+3} + 49 \cdot 2^{8s+4m-1} + 99 \cdot 2^{6s+3m-2} - 13 \cdot 2^{6s+4m-1} - 203 \cdot 2^{7s+3m-2},
\end{aligned}$$

hence we have (see above)

$$\begin{aligned}
\Gamma_{2s+m+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} & = 21 \cdot [5 \cdot 2^{8s+4m-2} + 7 \cdot 2^{6s+3m-3} - 2^{6s+4m-2} - 15 \cdot 2^{7s+3m-3}] \\
\Gamma_{2s+m+2}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} & = 2^{9s+6m} - 7 \cdot 2^{8s+4m+2} + 7 \cdot 2^{7s+3m+2} - 3 \cdot 2^{6s+3m+2} + 2^{6s+4m+2}
\end{aligned}$$

The case $j = 1$, $k \geq 2s + m + 2$

From (12.3) with $j = 0$ we get

$$\begin{aligned}
& \Gamma_{2s+m+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k} - \Gamma_{2s+m+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} \\
& = 21 \cdot 2^{2s+m-2} \cdot (2^{2k} - 2^{4s+2m+4}) + (2^k - 2^{2s+m+2}) \cdot (21 \cdot 2^{4s+3m-2} + 735 \cdot 2^{5s+2m-3} - 945 \cdot 2^{4s+2m-3}) \\
& \Downarrow \\
& \Gamma_{2s+m+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k} = 21 \cdot [5 \cdot 2^{8s+4m-2} + 7 \cdot 2^{6s+3m-3} - 2^{6s+4m-2} - 15 \cdot 2^{7s+3m-3}] \\
& + 21 \cdot 2^{2s+m-2} \cdot (2^{2k} - 2^{4s+2m+4}) + (2^k - 2^{2s+m+2}) \cdot (21 \cdot 2^{4s+3m-2} + 735 \cdot 2^{5s+2m-3} - 945 \cdot 2^{4s+2m-3}) \\
& \Downarrow \\
& \Gamma_{2s+m+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k} = 21 \cdot 2^{2k+2s+m-2} + 21 \cdot 2^{k+4s+3m-2} + 735 \cdot 2^{k+5s+2m-3} - 945 \cdot 2^{k+4s+2m-3} \\
& + 105 \cdot 2^{8s+4m-2} - 105 \cdot 2^{6s+4m-2} - 3255 \cdot 2^{7s+3m-3} + 3255 \cdot 2^{6s+3m-3}
\end{aligned}$$

The case $j = 2$, $k = 2s + m + 3$ and the case $j = 3$, $k = 2s + m + 3$

From (6.1) with $k = 2s + m + 3$, $l = 0$ we obtain using (12.7) and (12.8),(12.11)

$$\begin{aligned}
& \sum_{i=0}^{2s+m+3} \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} = 2^{9s+5m+6} \\
& \Downarrow \\
& \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} + \Gamma_s^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} + \sum_{j=1}^{m-1} \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} \\
& + \Gamma_{s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} + \sum_{j=1}^{s-1} \Gamma_{s+m+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} + \Gamma_{2s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} + \Gamma_{2s+m+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} \\
& + \Gamma_{2s+m+2}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} + \Gamma_{2s+m+3}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} = 2^{9s+5m+6} \\
& \Downarrow \\
& (7 \cdot 2^{4s-6} - 3 \cdot 2^{3s-5}) + (2^{3s+m+2} - 2^{2s} + 21 \cdot (5 \cdot 2^{4s-6} - 2^{3s-5})) \\
& + (3 \cdot 2^{3s+4m-2} - 3 \cdot 2^{3s+3m-5} + 3 \cdot 2^{2s+3m-5} - 2^{3s+m+2})
\end{aligned}$$

$$\begin{aligned}
& + 3 \cdot 2^{3s-2} + 2^{2s} + 7 \cdot 2^{4s+4m-6} - 7 \cdot 2^{2s+4m-6} - 7 \cdot 2^{4s-2} - 2^{3s+3m-1}) \\
& + (21 \cdot 2^{3s+4m-2} + 45 \cdot 2^{3s+3m-1} - 21 \cdot 2^{2s+4m-4} - 45 \cdot 2^{2s+3m-3} \\
& + 105 \cdot 2^{4s+4m-6} - 21 \cdot 2^{3s+3m-5} - 21 \cdot 2^{2s+4m-6} + 9 \cdot 2^{2s+3m-5}) \\
& + (3 \cdot 2^{6s+4m-2} - 27 \cdot 2^{6s+3m-3} - 3 \cdot 2^{3s+4m+1} - 11 \cdot 2^{3s+3m+1} \\
& + 49 \cdot 2^{7s+3m-4} + 51 \cdot 2^{6s+3m-7} + 3 \cdot 2^{3s+3m-2} + 7 \cdot 2^{8s+4m-6} - 7 \cdot 2^{6s+4m-6} \\
& - 7 \cdot 2^{4s+4m-2} + 7 \cdot 2^{2s+4m-2} - 105 \cdot 2^{7s+3m-8} + 21 \cdot 2^{2s+3m-2}) \\
& + 3 \cdot 2^{6s+3m+4} + 21 \cdot 2^{6s+4m-2} + 735 \cdot 2^{7s+3m-4} - 477 \cdot 2^{6s+3m-3} \\
& + 105 \cdot 2^{8s+4m-6} - 105 \cdot 2^{6s+4m-6} - 3255 \cdot 2^{7s+3m-8} + 1629 \cdot 2^{6s+3m-7} \\
& + 21 \cdot 2^{6s+3m+4} + 21 \cdot 2^{6m+4m+1} + 735 \cdot 2^{7s+3m} - 945 \cdot 2^{6s+3m} \\
& + 105 \cdot 2^{8s+4m-2} - 105 \cdot 2^{6s+4m-2} - 3255 \cdot 2^{7s+3m-3} + 3255 \cdot 2^{6s+3m-3} \\
& + \Gamma_{2s+m+1}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} + \Gamma_{2s+m+2}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} = 2^{9s+5m+3}
\end{aligned}$$

$\Downarrow$

$$\begin{aligned}
& \Gamma_{2s+m+1}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} + \Gamma_{2s+m+2}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+2)} \\
& = 2^{9s+5m+6} - 7 \cdot 2^{8s+4m+2} - 5 \cdot 2^{6s+4m+2} + 51 \cdot 2^{6s+3m+2} - 91 \cdot 2^{7s+3m+2}
\end{aligned}$$

From (6.2) with $k = 2s + m + 3$, $l = 0$ we obtain using (12.7) and (12.9),(12.13)

$$\begin{aligned}
& \sum_{i=0}^{2s+m+3} \Gamma_i^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} \cdot 2^{-i} = 2^{7s+4m+3} + 2^{6s+3m+6} - 2^{4s+2m+3} \\
& \Downarrow \\
& \sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} \cdot 2^{-i} + \Gamma_s^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} \cdot 2^{-s} + \sum_{j=1}^{m-1} \Gamma_{s+j}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} \cdot 2^{-(s+j)} \\
& + \Gamma_{s+m}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} \cdot 2^{-(s+m)} + \sum_{j=1}^{s-1} \Gamma_{s+m+j}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} \cdot 2^{-(s+m+j)} + \Gamma_{2s+m}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} \cdot 2^{-(2s+m)} \\
& + \Gamma_{2s+m+1}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} \cdot 2^{-(2s+m+1)} + \Gamma_{2s+m+2}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} \cdot 2^{-(2s+m+2)} \\
& + \Gamma_{2s+m+3}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} \cdot 2^{-(2s+m+3)} = 2^{7s+4m+3} + 2^{6s+3m+6} - 2^{4s+2m+3} \\
& \Downarrow \\
& (15 \cdot 2^{3s-6} - 7 \cdot 2^{2s-5}) + (2^{2s+m+2} - 2^s + 21 \cdot (5 \cdot 2^{3s-6} - 2^{2s-5})) \\
& + (7 \cdot 2^{2s+3m-2} - 7 \cdot 2^{2s+2m-5} + 7 \cdot 2^{s+2m-5} - 2^{2s+m+2} \\
& + 7 \cdot 2^{2s-3} + 2^s + 15 \cdot 2^{3s+3m-6} - 15 \cdot 2^{s+3m-6} - 15 \cdot 2^{3s-3} - 3 \cdot 2^{2s+2m-1}) \\
& + (21 \cdot 2^{2s+3m-2} + 45 \cdot 2^{2s+2m-1} - 21 \cdot 2^{s+3m-4} - 45 \cdot 2^{s+2m-3} \\
& + 105 \cdot 2^{3s+3m-6} - 21 \cdot 2^{2s+2m-5} - 21 \cdot 2^{s+3m-6} + 9 \cdot 2^{s+2m-5}) \\
& + (7 \cdot 2^{4s+3m-2} - 63 \cdot 2^{4s+2m-3} - 7 \cdot 2^{2s+3m} - 21 \cdot 2^{2s+2m} \\
& + 107 \cdot 2^{4s+2m-7} + 7 \cdot 2^{2s+2m-3} + 105 \cdot 2^{5s+2m-4} + 15 \cdot 2^{6s+3m-6} \\
& - 15 \cdot 2^{4s+3m-6} - 15 \cdot 2^{3s+3m-3} + 15 \cdot 2^{s+3m-3} - 217 \cdot 2^{5s+2m-8} + 41 \cdot 2^{s+2m-3}) \\
& + 3 \cdot 2^{4s+2m+4} + 21 \cdot 2^{4s+3m-2} + 735 \cdot 2^{5s+2m-4} - 477 \cdot 2^{4s+2m-3}
\end{aligned}$$

$$\begin{aligned}
& + 105 \cdot 2^{6s+3m-6} - 105 \cdot 2^{4s+3m-6} - 3255 \cdot 2^{5s+2m-8} + 1629 \cdot 2^{4s+2m-7} \\
& + 21 \cdot 2^{4s+2m+3} + 21 \cdot 2^{4s+3m} + 735 \cdot 2^{5s+2m-1} - 945 \cdot 2^{4s+2m-1} \\
& + 105 \cdot 2^{6s+3m-3} - 105 \cdot 2^{4s+3m-3} - 3255 \cdot 2^{5s+2m-4} + 3255 \cdot 2^{4s+2m-4} \\
& + \Gamma_{2s+m+2}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} \cdot 2^{-(2s+m+2)} + \Gamma_{2s+m+3}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} \cdot 2^{-(2s+m+3)} = 2^{7s+4m+3} + 2^{6s+3m+6} - 2^{4s+2m+3} \\
& \Downarrow \\
& 2 \cdot \Gamma_{2s+m+2}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} + \Gamma_{2s+m+3}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} \\
& = 2^{9s+5m+6} + 49 \cdot 2^{8s+4m+3} + 99 \cdot 2^{6s+3m+3} - 13 \cdot 2^{6s+4m+3} - 203 \cdot 2^{7s+3m+3},
\end{aligned}$$

hence we have (see above)

$$\begin{aligned}
\Gamma_{2s+m+2}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} & = 21 \cdot [5 \cdot 2^{8s+4m+2} + 7 \cdot 2^{6s+3m+2} - 2^{6s+4m+2} - 15 \cdot 2^{7s+3m+2}] \\
\Gamma_{2s+m+3}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} & = 2^{9s+6m+6} - 7 \cdot 2^{8s+4m+6} + 7 \cdot 2^{7s+3m+7} - 3 \cdot 2^{6s+3m+7} + 2^{6s+4m+6}
\end{aligned}$$

The case $j = 2$, $k \geq 2s + m + 3$

From (12.3) with $j = 1$ we get

$$\begin{aligned}
& \Gamma_{2s+m+2}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times k} - \Gamma_{2s+m+2}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+3)} \\
& = 21 \cdot 2^{2s+m+1} \cdot (2^{2k} - 2^{4s+2m+6}) + (2^k - 2^{2s+m+3}) \cdot (21 \cdot 2^{4s+3m+1} + 735 \cdot 2^{5s+2m+1} - 945 \cdot 2^{4s+2m+1}) \\
& \Downarrow \\
& \Gamma_{2s+m+2}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times k} = 21 \cdot [5 \cdot 2^{8s+4m+2} + 7 \cdot 2^{6s+3m+2} - 2^{6s+4m+2} - 15 \cdot 2^{7s+3m+2}] \\
& + 21 \cdot 2^{2s+m+1} \cdot (2^{2k} - 2^{4s+2m+6}) + (2^k - 2^{2s+m+3}) \cdot (21 \cdot 2^{4s+3m+1} + 735 \cdot 2^{5s+2m+1} - 945 \cdot 2^{4s+2m+1}) \\
& \Downarrow \\
& \Gamma_{2s+m+2}^{\left[\begin{smallmatrix} s+m \\ s+m \end{smallmatrix} \right] \times k} = 21 \cdot 2^{2k+2s+m+1} + 21 \cdot 2^{k+4s+3m+1} + 735 \cdot 2^{k+5s+2m+1} - 945 \cdot 2^{k+4s+2m+1} \\
& + 105 \cdot 2^{8s+4m+2} - 105 \cdot 2^{6s+4m+2} - 3255 \cdot 2^{7s+3m+2} + 3255 \cdot 2^{6s+3m+2}
\end{aligned}$$

□

12.6. An explicit formula for $\Gamma_i^{\left[\begin{smallmatrix} s+1 \\ s+1 \end{smallmatrix} \right] \times k}$, for $s \leq i \leq 3s+2$, $k \geq i$. We shall use the following results obtained in Lemma 12.2

Lemma 12.8. *We have for $m = 1$, $k \geq s + j + 1$*

(12.16)

$$\begin{aligned}
& \Gamma_{s+j}^{\left[\begin{smallmatrix} s+1 \\ s+1 \end{smallmatrix} \right] \times k} - \Gamma_{s+j}^{\left[\begin{smallmatrix} s+1 \\ s+1 \end{smallmatrix} \right] \times (s+j+1)} \\
& = \begin{cases} 2^{s-1} \cdot (2^k - 2^{s+1}) & \text{if } j = 0 \\ 33 \cdot 2^{s-1} \cdot (2^k - 2^{s+2}) & \text{if } j = 1 \\ (735 \cdot 2^{s+4j-9} - 105 \cdot 2^{s+3j-7}) \cdot (2^k - 2^{s+j+1}) & \text{if } 2 \leq j \leq s \\ 3 \cdot 2^{2s-1} \cdot (2^{2k} - 2^{4s+4}) + (735 \cdot 2^{5s-5} - 393 \cdot 2^{4s-4}) \cdot (2^k - 2^{2s+2}) & \text{if } j = s+1, \\ 53 \cdot 2^{2s-1} \cdot (2^{2k} - 2^{4s+6}) + (735 \cdot 2^{5s-1} - 1629 \cdot 2^{4s-1}) \cdot (2^k - 2^{2s+3}) & \text{if } j = s+2, \end{cases}
\end{aligned}$$

We have for $m = 1$, $k \geq 2s + 2 + j$

(12.17)

$$\begin{aligned} & \Gamma_{2s+1+j}^{\left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix} \right] \times k} - \Gamma_{2s+1+j}^{\left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix} \right] \times (2s+2+j)} \\ &= \begin{cases} 105 \cdot 2^{2s+4j-6} \cdot (2^{2k} - 2^{4s+2j+4}) + (735 \cdot 2^{5s+4j-5} - 3255 \cdot 2^{4s+5j-7}) \cdot (2^k - 2^{2s+j+2}) & \text{if } 2 \leq j \leq s, \\ 2^{3s-1} \cdot (2^{3k} - 2^{9s+6}) - 7 \cdot 2^{6s-2} \cdot (2^{2k} - 2^{6s+6}) + 7 \cdot 2^{9s-2} \cdot (2^k - 2^{3s+3}) & \text{if } j = s+1 \end{cases} \end{aligned}$$

Theorem 12.9. *We have*

(12.18)

$$\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix} \right] \times (s+j+1)} = \begin{cases} 105 \cdot 2^{4s-6} - 21 \cdot 2^{3s-5} & \text{if } j = 0, \\ 105 \cdot 2^{4s-2} - 21 \cdot 2^{3s-2} - 3 \cdot 2^{2s} & \text{if } j = 1, \\ 21 \cdot [5 \cdot 2^{4s+2} - 2^{3s+1} - 5 \cdot 2^{2s+1}] & \text{if } j = 2, \\ 21 \cdot [5 \cdot 2^{4s+6} - 2^{3s+4} - 25 \cdot 2^{2s+4}] & \text{if } j = 3, \\ 21 \cdot [5 \cdot 2^{4s+10} - 2^{3s+7} - 55 \cdot 2^{2s+8}] & \text{if } j = 4, \\ 21 \cdot [5 \cdot 2^{4s+4j-6} - 2^{3s+3j-5} - 15 \cdot 2^{2s+5j-10} + 5 \cdot 2^{2s+4j-8}] & \text{if } 2 \leq j \leq s+2, \\ 21 \cdot [5 \cdot 2^{8s-2} + 2^{6s-4} - 15 \cdot 2^{7s-5}] & \text{if } j = s+1, \\ 21 \cdot [5 \cdot 2^{8s+2} + 3 \cdot 2^{6s} - 15 \cdot 2^{7s}] & \text{if } j = s+2 \end{cases}$$

(12.19)

$$\Gamma_{2s+3+j}^{\left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix} \right] \times (2s+4+j)} = \begin{cases} 21 \cdot [5 \cdot 2^{8s+4j+6} + 5 \cdot 2^{6s+6j+5} - 15 \cdot 2^{7s+5j+5}] & \text{if } 0 \leq j \leq s-2, \\ 315 \cdot 2^{12s-1} & \text{if } j = s-1 \end{cases}$$

We have

(12.20)

$$\Gamma_i^{\left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix} \right] \times k} = \begin{cases} 1 & \text{if } i = 0, \\ 105 \cdot 2^{4i-6} - 21 \cdot 2^{3i-5} & \text{if } 1 \leq i \leq s-1, k \geq i+1, \\ 2^{k+s-1} - 2^{2s} + 105 \cdot 2^{4s-6} - 21 \cdot 2^{3s-5} & \text{if } i = s, k \geq s+1, s \geq 1, \\ 33 \cdot 2^{k+s-1} + [105 \cdot 2^{4s-2} - 21 \cdot 2^{3s-2} - 69 \cdot 2^{2s}] & \text{if } i = s+1, k \geq s+2, \\ 630 \cdot 2^{k+s-1} + 21 \cdot [5 \cdot 2^{4s+2} - 2^{3s+1} - 65 \cdot 2^{2s+1}] & \text{if } i = s+2, k \geq s+3, \\ 1365 \cdot 2^{k+s+2} + 21 \cdot [5 \cdot 2^{4s+6} - 2^{3s+4} - 285 \cdot 2^{2s+4}] & \text{if } i = s+3, k \geq s+4, \\ 2835 \cdot 2^{k+s+5} + 21 \cdot [5 \cdot 2^{4s+10} - 2^{3s+7} - 595 \cdot 2^{2s+8}] & \text{if } i = s+4, k \geq s+5, \\ 105 \cdot (7 \cdot 2^{j-2} - 1) \cdot 2^{k+s+3j-7} & \text{if } i = s+j, \\ + 21 \cdot [5 \cdot 2^{4s+4j-6} - 2^{3s+3j-5} - 155 \cdot 2^{2s+5j-10} + 25 \cdot 2^{2s+4j-8}] & 2 \leq j \leq s, k \geq s+j+1, \\ 3 \cdot 2^{2s-1} \cdot (2^{2k} - 2^{4s+4}) + (735 \cdot 2^{5s-5} - 393 \cdot 2^{4s-4}) \cdot (2^k - 2^{2s+2}) & \text{if } i = 2s+1, k \geq 2s+2, \\ + 21 \cdot [5 \cdot 2^{8s-2} + 2^{6s-4} - 15 \cdot 2^{7s-5}] & \text{if } i = 2s+2, k \geq 2s+3, \\ 53 \cdot 2^{2s-1} \cdot (2^{2k} - 2^{4s+6}) + (735 \cdot 2^{5s-1} - 1629 \cdot 2^{4s-1}) \cdot (2^k - 2^{2s+3}) & \text{if } i = 2s+3, k \geq 2s+4, \\ + 21 \cdot [5 \cdot 2^{8s+2} + 3 \cdot 2^{6s} - 15 \cdot 2^{7s}] & \text{if } i = 2s+4, k \geq 2s+5, \\ 105 \cdot (2^{2k+2s+4j+2} + 7 \cdot 2^{k+5s+4j+3} - 31 \cdot 2^{k+4s+5j+3}) & \text{if } i = 2s+5, k \geq 2s+6, \\ + 105 \cdot (2^{8s+4j+6} - 31 \cdot 2^{7s+5j+5} + 93 \cdot 2^{6s+6j+5}) & \text{if } i = 2s+6, k \geq 2s+7, \\ 2^{3k+3s-1} - 7 \cdot 2^{2k+6s-2} + 7 \cdot 2^{k+9s-2} - 2^{12s-1} & 0 \leq j \leq s-2, \\ & \text{if } i = 3s+2, k \geq 3s+2 \end{cases}$$

$$(12.21) \quad \Gamma_i \begin{bmatrix} s+1 \\ s+1 \end{bmatrix}^{\times i} = \begin{cases} 2^{3s+3i-1} - 7 \cdot 2^{4i-6} + 3 \cdot 2^{3i-5} & \text{if } 1 \leq i \leq s+1, \\ 2^{6s+3j-1} - 7 \cdot 2^{4s+4j-6} + 3 \cdot 2^{3s+3j-5} \\ + 7 \cdot 2^{2s+5j-10} - 5 \cdot 2^{2s+4j-8} & \text{if } i = s+j, 2 \leq j \leq s+3, \\ 2^{9s+2} + 7 \cdot 2^{7s-5} - 7 \cdot 2^{8s-2} + 7 \cdot 2^{6s-4} & \text{if } i = 2s+1, \\ 2^{9s+5} + 7 \cdot 2^{7s} - 7 \cdot 2^{8s+2} + 2^{6s} & \text{if } i = 2s+2, \\ 2^{9s+8} + 7 \cdot 2^{7s+5} - 7 \cdot 2^{8s+6} - 2^{6s+5} & \text{if } i = 2s+3, \\ 2^{9s+3j+8} - 7 \cdot 2^{8s+4j+6} + 7 \cdot 2^{7s+5j+5} - 2^{6s+6j+5} & \text{if } i = 2s+3+j, 0 \leq j \leq s-1 \end{cases}$$

We have the following reduction formulas

$$(12.22) \quad \Gamma_{2s+2+j} \begin{bmatrix} s+1 \\ s+1 \end{bmatrix}^{\times k} = 16^{2j} \Gamma_{2s+2-j} \begin{bmatrix} s+(1-j) \\ s+(1-j) \end{bmatrix}^{\times (k-2j)} \quad \text{if } 0 \leq j \leq 1$$

$$(12.23) \quad \Gamma_{2s+3+j} \begin{bmatrix} s+1 \\ s+1 \end{bmatrix}^{\times k} = 16^{2+3j} \Gamma_{2(s-j)+1} \begin{bmatrix} s-j \\ s-j \end{bmatrix}^{\times (k-2-3j)} \quad \text{if } 0 \leq j \leq s-1$$

Example. We have for $s = 3$:

$$\Gamma_i \begin{bmatrix} 3+1 \\ 3+1 \end{bmatrix}^{\times k} = \begin{cases} 1 & \text{if } i = 0, k \geq 1, \\ 21 & \text{if } i = 1, k \geq 2, \\ 378 & \text{if } i = 2, k \geq 3, \\ 2^{k+2} + 6320 & \text{if } i = 3, k \geq 4, \\ 33 \cdot 2^{k+2} + 100416 & \text{if } i = 4, k \geq 5, \\ 630 \cdot 2^{k+2} + 1524096 & \text{if } i = 5, k \geq 6, \\ 1365 \cdot 2^{k+5} + 21224448 & \text{if } i = 6, k \geq 7, \\ 96 \cdot 2^{2k} + 163008 \cdot 2^{k+2} + 1029 \cdot 2^{18} & \text{if } i = 7, k \geq 8, \\ 1696 \cdot 2^{2k} + 2176512 \cdot 2^{k+2} + 5723 \cdot 2^{18} & \text{if } i = 8, k \geq 9, \\ 105 \cdot 2^{2k+8} + 2625 \cdot 2^{k+15} - 90720 \cdot 2^{18} & \text{if } i = 9, k \geq 10, \\ 105 \cdot 2^{2k+12} - 315 \cdot 2^{k+20} + 215040 \cdot 2^{18} & \text{if } i = 10, k \geq 11, \\ 2^{3k+8} - 7 \cdot 2^{2k+16} + 7 \cdot 2^{k+25} - 2^{35} & \text{if } i = 11, k \geq 11. \end{cases}$$

Proof. To prove Theorem 12.9 we proceed as in Theorem 10.9. □

12.7. **Computation of** $\Gamma_i \begin{bmatrix} s+m \\ s+m \end{bmatrix}^{\times k}$ *for* $2s+m+1 \leq i \leq 3s+2m$, $k \geq i$, $m \geq 2$.

Lemma 12.10. *We have for* $m \geq 2$, $0 \leq j \leq m-1$:

$$(12.24) \quad \Gamma_{2s+m+1+j} \begin{bmatrix} s+m \\ s+m \end{bmatrix}^{\times k} = \begin{cases} 21 \cdot 2^{2k+2s+m-2} + 21 \cdot 2^{k+4s+3m-2} + 735 \cdot 2^{k+5s+2m-3} - 945 \cdot 2^{k+4s+2m-3} \\ + 105 \cdot 2^{8s+4m-2} - 105 \cdot 2^{6s+4m-2} - 3255 \cdot 2^{7s+3m-3} + 3255 \cdot 2^{6s+3m-3} & \text{if } j = 0, \\ & k \geq 2s+m+2 \\ 21 \cdot 2^{2k+2s+m+3j-2} + 21 \cdot 2^{k+4s+3m+3j-2} + 735 \cdot 2^{k+5s+2m+4j-3} - 945 \cdot 2^{k+4s+2m+4j-3} \\ + 105 \cdot 2^{8s+4m+4j-2} - 105 \cdot 2^{6s+4m+4j-2} - 3255 \cdot 2^{7s+3m+5j-3} + 3255 \cdot 2^{6s+3m+5j-3} & \text{if } 0 \leq j \leq m-2, \\ & k \geq 2s+m+2+j \\ 53 \cdot 2^{2s-1} \cdot (2^{2k+4m-4} - 2^{4s+8m-2}) + (735 \cdot 2^{5s-1} - 1629 \cdot 2^{4s-1}) \cdot (2^{k+6m-6} - 2^{2s+8m-5}) \\ + 21 \cdot [5 \cdot 2^{8s+8m-6} + 3 \cdot 2^{6s+8m-8} - 15 \cdot 2^{7s+8m-8}] & \text{if } j = m-1, \\ & k \geq 2s+2m+1 \end{cases}$$

$$\begin{aligned}
& \Gamma_{2s+m+1+j} \left[\begin{smallmatrix} s \\ s+m \\ s+m \end{smallmatrix} \right] \times (2s+m+1+j) \\
(12.25) \quad & = \begin{cases} 2^{9s+6m} - 7 \cdot 2^{8s+4m-2} + 7 \cdot 2^{7s+3m-3} - 3 \cdot 2^{6s+3m-3} + 2^{6s+4m-2} & \text{if } j = 0 \\ 2^{9s+6m+2j} - 7 \cdot 2^{8s+4m+4j-2} + 7 \cdot 2^{7s+3m+5j-3} - 3 \cdot 2^{6s+3m+5j-3} + 2^{6s+4m+4j-2} & \text{if } 0 \leq j \leq m-2 \\ 2^{9s+8m-3} - 7 \cdot 2^{8s+8m-6} + 7 \cdot 2^{7s+8m-8} - 2^{6s+8m-8} & \text{if } j = m-1 \end{cases}
\end{aligned}$$

We have for $m \geq 2$, $0 \leq j \leq s-1$:

$$\begin{aligned}
& \Gamma_{2s+2m+1+j} \left[\begin{smallmatrix} s \\ s+m \\ s+m \end{smallmatrix} \right] \times k \\
(12.26) \quad & = \begin{cases} 105 \cdot (2^{2k+2s+4m+4j-2} + 7 \cdot 2^{k+5s+6m+4j-3} - 31 \cdot 2^{k+4s+6m+5j-3}) \\ + 105 \cdot (2^{8s+8m+4j-2} - 31 \cdot 2^{7s+8m+5j-3} + 93 \cdot 2^{6s+8m+6j-3}) & \text{if } 0 \leq j \leq s-2, k \geq 2s+2m+2+j \\ 2^{3k+2m+3s-3} - 7 \cdot 2^{2k+4m+6s-6} + 7 \cdot 2^{k+6m+9s-8} - 2^{8m+12s-9} & \text{if } j = s-1, k \geq 3s+2m \end{cases}
\end{aligned}$$

$$\begin{aligned}
& \Gamma_{2s+2m+1+j} \left[\begin{smallmatrix} s \\ s+m \\ s+m \end{smallmatrix} \right] \times (2s+2m+1+j) \\
(12.27) \quad & = \begin{cases} 2^{9s+6m} - 7 \cdot 2^{8s+8m-2} + 7 \cdot 2^{7s+8m-3} - 2^{6s+8m-3} & \text{if } j = 0 \\ 2^{9s+8m+3j} - 7 \cdot 2^{8s+8m+4j-2} + 7 \cdot 2^{7s+8m+5j-3} - 2^{6s+8m+6j-3} & \text{if } 0 \leq j \leq s-2 \\ 21 \cdot 2^{8m+12s-9} & \text{if } j = s-1 \end{cases}
\end{aligned}$$

We have the following reduction formulas

$$(12.28) \quad \Gamma_{2s+m+1+j} \left[\begin{smallmatrix} s \\ s+m \\ s+m \end{smallmatrix} \right] \times k = 16^{2j} \Gamma_{2s+1+(m-j)} \left[\begin{smallmatrix} s \\ s+(m-j) \\ s+(m-j) \end{smallmatrix} \right] \times (k-2j) \quad \text{if } 0 \leq j \leq m-1$$

$$(12.29) \quad \Gamma_{2s+2m+1+j} \left[\begin{smallmatrix} s \\ s+m \\ s+m \end{smallmatrix} \right] \times k = 16^{2m+3j} \Gamma_{2(s-j)+1} \left[\begin{smallmatrix} s-j \\ s-j \\ s-j \end{smallmatrix} \right] \times (k-2m-3j) \quad \text{if } 0 \leq j \leq s-1$$

Proof.

□

Example. $s = m = 2$

$$\begin{aligned}
& \Gamma_i \left[\begin{smallmatrix} 2 \\ 2+2 \\ 2+2 \end{smallmatrix} \right] \times k \\
& = \begin{cases} 1 & \text{if } i = 0, k \geq 1 \\ 21 & \text{if } i = 1, k \geq 2 \\ 2^{k+1} + 362 & \text{if } i = 2, k \geq 3 \\ 9 \cdot 2^{k+1} + 6048 & \text{if } i = 3, k \geq 4 \\ 348 \cdot 2^k + 92640 & \text{if } i = 4, k \geq 5 \\ 5712 \cdot 2^k + 1295616 & \text{if } i = 5, k \geq 6 \\ 3 \cdot 2^{2k+4} + 1161 \cdot 2^{k+6} + 15808512 & \text{if } i = 6, k \geq 7 \\ 21 \cdot 2^{2k+4} + 2163 \cdot 2^{k+9} + 92897280 & \text{if } i = 7, k \geq 8 \\ 424 \cdot 2^{2k+4} + 167808 \cdot 2^{k+6} - 1485832192 & \text{if } i = 8, k \geq 9 \\ 105 \cdot 2^{2k+10} - 315 \cdot 2^{k+17} + 3523215360 & \text{if } i = 9, k \geq 10 \\ 2^{3k+7} - 7 \cdot 2^{2k+14} + 7 \cdot 2^{k+22} - 2^{31} & \text{if } i = 10, k \geq 10 \end{cases}
\end{aligned}$$

12.8. An explicit formula for $\Gamma_i \begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix} \times k$, for $1 \leq i \leq 3s+2m$, $k \geq i$.

Theorem 12.11. *We have*

(12.30)

$$\Gamma_i \begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix} \times k = \begin{cases} 1 & \text{if } i = 0, k \geq 1 \\ 105 \cdot 2^{4i-6} - 21 \cdot 2^{3i-5} & \text{if } 1 \leq i \leq s-1, k \geq i+1, \\ 2^{k+s-1} - 2^{2s} + 21 \cdot (5 \cdot 2^{4s-6} - 2^{3s-5}) & \text{if } i = s, k \geq s+1 \\ \\ (21 \cdot 2^{j-1} - 3) \cdot 2^{k+s+2j-4} & \\ + 21 \cdot (5 \cdot 2^{4s+4j-6} - 2^{3s+3j-5} - 5 \cdot 2^{2s+4j-6} + 2^{2s+3j-5}) & \text{if } i = s+j, 1 \leq j \leq m-1, \\ & k \geq s+j+1 \\ \\ (21 \cdot 2^{s+3m-5} + 45 \cdot 2^{s+2m-4}) \cdot (2^k - 2^{s+m+1}) & \\ + 105 \cdot 2^{4s+4m-6} - 21 \cdot 2^{3s+3m-5} - 21 \cdot 2^{2s+4m-6} + 9 \cdot 2^{2s+3m-5} & \text{if } i = s+m, k \geq s+m+1 \\ \\ 21 \cdot [2^{k+s+3m+3j-5} + 35 \cdot 2^{k+s+2m+4j-7} - 9 \cdot 2^{k+s+2m+3j-6} & \\ + 5 \cdot 2^{4s+4m+4j-6} - 2^{3s+3m+3j-5} - 5 \cdot 2^{2s+4m+4j-6} & \\ - 155 \cdot 2^{2s+3m+5j-8} + 45 \cdot 2^{2s+3m+4j-7}] & \text{if } i = s+m+j, 1 \leq j \leq s-1, \\ & k \geq s+m+j+1 \\ \\ 3 \cdot 2^{2k+2s+m-2} + 21 \cdot 2^{k+4s+3m-5} + 735 \cdot 2^{k+5s+2m-7} - 477 \cdot 2^{k+4s+2m-6} & \\ + 105 \cdot 2^{8s+4m-6} - 105 \cdot 2^{6s+4m-6} - 3255 \cdot 2^{7s+3m-8} + 1629 \cdot 2^{6s+3m-7} & \text{if } i = 2s+m, k \geq 2s+m+1 \\ \\ 21 \cdot 2^{2k+2s+m+3j-2} + 21 \cdot 2^{k+4s+3m+3j-2} + 735 \cdot 2^{k+5s+2m+4j-3} & \\ - 945 \cdot 2^{k+4s+2m+4j-3} + 105 \cdot 2^{8s+4m+4j-2} - 105 \cdot 2^{6s+4m+4j-2} & \\ - 3255 \cdot 2^{7s+3m+5j-3} + 3255 \cdot 2^{6s+3m+5j-3} & \text{if } i = 2s+m+1+j, \\ & 0 \leq j \leq m-2, k \geq 2s+m+2+j \\ \\ 53 \cdot 2^{2s-1} \cdot (2^{2k+4m-4} - 2^{4s+8m-2}) & \\ + (735 \cdot 2^{5s-1} - 1629 \cdot 2^{4s-1}) \cdot (2^{k+6m-6} - 2^{2s+8m-5}) & \\ + 21 \cdot [5 \cdot 2^{8s+8m-6} + 3 \cdot 2^{6s+8m-8} - 15 \cdot 2^{7s+8m-8}] & \text{if } i = 2s+2m, k \geq 2s+2m+1 \\ \\ 105 \cdot (2^{2k+2s+4m+4j-2} + 7 \cdot 2^{k+5s+6m+4j-3} - 31 \cdot 2^{k+4s+6m+5j-3}) & \\ + 105 \cdot (2^{8s+8m+4j-2} - 31 \cdot 2^{7s+8m+5j-3} + 93 \cdot 2^{6s+8m+6j-3}) & \text{if } i = 2s+2m+1+j, \\ & 0 \leq j \leq s-2, k \geq 2s+2m+2+j \\ \\ 2^{2k+2m+3s-3} - 7 \cdot 2^{2k+4m+6s-6} + 7 \cdot 2^{k+6m+9s-8} - 2^{8m+12s-9} & \text{if } i = 3s+2m, k \geq 3s+2m \end{cases}$$

(12.31)

$$\Gamma_i \begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix} \times i = \begin{cases} 2^{3s+2m+3i-3} - 7 \cdot 2^{4i-6} + 3 \cdot 2^{3i-5} & \text{if } 1 \leq i \leq s+1 \\ \\ 2^{6s+2m+3j-3} - 7 \cdot 2^{4s+4j-6} + 3 \cdot 2^{3s+3j-5} \\ + (2^{j-1} - 1) \cdot 2^{2s+3j-5} & \text{if } i = s+j, 1 \leq j \leq m+1 \\ \\ 2^{6s+5m+3j-3} - 7 \cdot 2^{4s+4m+4j-6} + 3 \cdot 2^{3s+3m+3j-5} \\ + 2^{2s+4m+4j-6} + 7 \cdot 2^{2s+3m+5j-8} - 9 \cdot 2^{2s+3m+4j-7} & \text{if } i = s+m+j, 1 \leq j \leq s-1 \\ \\ 2^{9s+5m-3} - 7 \cdot 2^{8s+4m-6} + 7 \cdot 2^{7s+3m-8} + 3 \cdot 2^{6s+3m-7} + 2^{6s+4m-6} & \text{if } i = 2s+m \\ \\ 2^{9s+5m+2j} - 7 \cdot 2^{8s+4m+4j-2} + 7 \cdot 2^{7s+3m+5j-3} - 3 \cdot 2^{6s+3m+5j-3} + 2^{6s+4m+4j-2} & \text{if } i = 2s+m+1+j, \\ & 0 \leq j \leq m-2 \\ \\ 2^{9s+8m-3} - 7 \cdot 2^{8s+8m-6} + 7 \cdot 2^{7s+8m-8} - 2^{6s+8m-8} & \text{if } i = 2s+2m \\ \\ 2^{9s+8m+3j} - 7 \cdot 2^{8s+8m+4j-2} + 7 \cdot 2^{7s+8m+5j-3} - 2^{6s+8m+6j-3} & \text{if } i = 2s+2m+1+j, \\ & 0 \leq j \leq s-2 \\ \\ 21 \cdot 2^{8m+12s-9} & \text{if } i = 3s+2m \end{cases}$$

We have for $0 \leq j \leq m-2$, $k \geq 2s+m+2+j$

$$\begin{aligned} (12.32) \quad & \Gamma_{2s+m+1+j} \begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix} \times k = 16^{2j} \Gamma_{2s+1+(m-j)} \begin{bmatrix} s \\ s+(m-j) \\ s+(m-j) \end{bmatrix} \times (k-2j) \\ & = 2^{8j} \cdot [21 \cdot 2^{2(k-2j)+2s+(m-j)-2} + 21 \cdot 2^{(k-2j)+4s+3(m-j)-2} + 735 \cdot 2^{k-2j+5s+2(m-j)-3} \\ & \quad - 945 \cdot 2^{k-2j+4s+2(m-j)-3} + 105 \cdot 2^{8s+4(m-j)-2} - 105 \cdot 2^{6s+4(m-j)-2} \\ & \quad - 3255 \cdot 2^{7s+3(m-j)-3} + 3255 \cdot 2^{6s+3(m-j)-3}] \\ & = 21 \cdot 2^{2k+2s+m+3j-2} + 21 \cdot 2^{k+4s+3m+3j-2} + 735 \cdot 2^{k+5s+2m+4j-3} \\ & \quad - 945 \cdot 2^{k+4s+2m+4j-3} + 105 \cdot 2^{8s+4m+4j-2} - 105 \cdot 2^{6s+4m+4j-2} \\ & \quad - 3255 \cdot 2^{7s+3m+5j-3} + 3255 \cdot 2^{6s+3m+5j-3} \end{aligned}$$

We have for $j = m-1$, $k \geq 2s+2m+1$

$$\begin{aligned} (12.33) \quad & \Gamma_{2s+2m} \begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix} \times k = 16^{2m-2} \Gamma_{2s+2} \begin{bmatrix} s \\ s+1 \\ s+1 \end{bmatrix} \times (k-2(m-1)) \\ & = 2^{8m-8} \cdot [53 \cdot 2^{2s-1} \cdot (2^{2(k-2m+2)} - 2^{4s+6}) + (735 \cdot 2^{5s-1} - 1629 \cdot 2^{4s-1}) \cdot (2^{k-2m+2} - 2^{2s+3}) \\ & \quad + 21 \cdot [5 \cdot 2^{8s+2} + 3 \cdot 2^{6s} - 15 \cdot 2^{7s}]] \\ & = 53 \cdot 2^{2s-1} \cdot (2^{2k+4m-4} - 2^{4s+8m-2}) + (735 \cdot 2^{5s-1} - 1629 \cdot 2^{4s-1}) \cdot (2^{k+6m-6} - 2^{2s+8m-5}) \\ & \quad + 21 \cdot [5 \cdot 2^{8s+8m-6} + 3 \cdot 2^{6s+8m-8} - 15 \cdot 2^{7s+8m-8}] \end{aligned}$$

We have for $0 \leq j \leq s-2$, $k \geq 2s+2m+2+j$

$$\begin{aligned}
 (12.34) \quad \Gamma_{2s+2m+1+j}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix} \times k} &= 16^{2m+3j} \Gamma_{2(s-j)+1}^{\begin{bmatrix} s-j \\ s-j \\ s-j \end{bmatrix} \times (k-2m-3j)} \\
 &= 2^{8m+12j} \cdot 105 \cdot (2^{2(k-2m-3j)+2(s-j)-2} + 7 \cdot 2^{k-2m-3j+5(s-j)-3} - 31 \cdot 2^{k-2m-3j+4(s-j)-3} \\
 &\quad + 2^{8(s-j)-2} - 31 \cdot 2^{7(s-j)-3} + 93 \cdot 2^{6(s-j)-3}) \\
 &= 105 \cdot (2^{2k+2s+4m+4j-2} + 7 \cdot 2^{k+5s+6m+4j-3} - 31 \cdot 2^{k+4s+6m+5j-3}) \\
 &\quad + 105 \cdot (2^{8s+8m+4j-2} - 31 \cdot 2^{7s+8m+5j-3} + 93 \cdot 2^{6s+8m+6j-3})
 \end{aligned}$$

We have for $j = s-1$, $k \geq 3s+2m$

$$\begin{aligned}
 (12.35) \quad \Gamma_{3s+2m}^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix} \times k} &= 16^{2m+3s-3} \Gamma_3^{\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \times (k-2m-3s+3)} \\
 &= 2^{8m+12s-12} \cdot [2^{3(k-2m-3s+3)} - 7 \cdot 2^{2(k-2m-3s+3)} + 7 \cdot 2^{k-2m-3s+3+1} - 2^3] \\
 &= 2^{3k+2m+3s-3} - 7 \cdot 2^{2k+4m+6s-6} + 7 \cdot 2^{k+6m+9s-8} - 2^{8m+12s-9}
 \end{aligned}$$

(12.36)

Let q be a rational integer ≥ 1 , then

$$\begin{aligned}
 g_{k,s,m}(t, \eta, \xi) &= g(t, \eta, \xi) = \sum_{\deg Y \leq k-1} \sum_{\deg Z \leq s-1} E(tYZ) \sum_{\deg U \leq s+m-1} E(\eta YU) \sum_{\deg V \leq s+m-1} E(\eta YV) \\
 &= 2^{3s+2m+k-r(D \begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix} \times k)}(t, \eta, \xi) \\
 &\int_{\mathbb{P}^3} g^q(t, \eta, \xi) dt d\eta d\xi = \\
 &= \sum_{\substack{(t, \eta, \xi) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \\ r(D \begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix} \times k) \\ (t, \eta, \xi) = i}} 2^{\left(k+3s+2m-r(D \begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix} \times k)(t, \eta, \xi)\right)q} \int_{\mathbb{P}_{k+s-1}} dt \int_{\mathbb{P}_{k+s+m-1}} d\eta \int_{\mathbb{P}_{k+s+m-1}} d\xi \\
 &= \sum_{i=0}^{\inf(k, 3s+2m)} \sum_{(t, \eta, \xi) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1}} 2^{(k+3s+2m-i)q} \int_{\mathbb{P}_{k+s-1}} dt \int_{\mathbb{P}_{k+s+m-1}} d\eta \int_{\mathbb{P}_{k+s+m-1}} d\xi \\
 &= 2^{(k+3s+2m)q - (3k+3s+2m-3)} \sum_{i=0}^{\inf(k, 3s+2m)} \Gamma_i^{\begin{bmatrix} s \\ s+m \\ s+m \end{bmatrix} \times k} \cdot 2^{-iq}
 \end{aligned}$$

We denote by $R_q(k, s, m)$ the number of solutions $(Y_1, Z_1, U_1, V_1, \dots, Y_q, Z_q, U_q, V_q)$ of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0, \\ Y_1 U_1 + Y_2 U_2 + \dots + Y_q U_q = 0, \\ Y_1 V_1 + Y_2 V_2 + \dots + Y_q V_q = 0, \end{cases}$$

satisfying the degree conditions

$$\deg Y_i \leq k-1, \quad \deg Z_i \leq s-1, \quad \deg U_i \leq s+m-1, \quad \deg V_i \leq s+m-1 \quad \text{for } 1 \leq i \leq q.$$

Then

(12.37)

$$R_q(k, s, m) = \int_{\mathbb{P} \times \mathbb{P} \times \mathbb{P}} g_{k,s,m}^q(t, \eta, \xi) dt d\eta d\xi = 2^{(k+3s+2m)q - (3k+3s+2m-3)} \sum_{i=0}^{\inf(k, 3s+2m)} \Gamma_i \left[\begin{smallmatrix} s \\ s+m \end{smallmatrix} \right] \times k \cdot 2^{-iq}$$

Example. We have for $s = 3$, $m = 4$, $k = 10$:

$$\Gamma_i \left[\begin{smallmatrix} 3 \\ 3+4 \\ 3+4 \end{smallmatrix} \right] \times 10 = \begin{cases} 1 & \text{if } i = 0 \\ 21 & \text{if } i = 1 \\ 378 & \text{if } i = 2 \\ 10416 & \text{if } i = 3 \\ 140352 & \text{if } i = 4, \\ 1994112 & \text{if } i = 5, \\ 29598720 & \text{if } i = 6 \\ 458661888 & \text{if } i = 7, \\ 109389 \cdot 2^{16} & \text{if } i = 8, \\ 213759 \cdot 2^{19} & \text{if } i = 9, \\ 2^{44} - 14273 \cdot 2^{23} & \text{if } i = 10 \end{cases}$$

Example. $s = 3$, $m=4$, $k = 7$, $q = 3$

The number $\Gamma_i \left[\begin{smallmatrix} 3 \\ 3+4 \\ 3+4 \end{smallmatrix} \right] \times 7$ of rank i matrices of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 & \alpha_7 \\ \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 & \alpha_7 & \alpha_8 \\ \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 & \alpha_7 & \alpha_8 & \alpha_9 \\ \beta_1 & \beta_2 & \beta_3 & \beta_4 & \beta_5 & \beta_6 & \beta_7 \\ \beta_2 & \beta_3 & \beta_4 & \beta_5 & \beta_6 & \beta_7 & \beta_8 \\ \beta_3 & \beta_4 & \beta_5 & \beta_6 & \beta_7 & \beta_8 & \beta_9 \\ \beta_4 & \beta_5 & \beta_6 & \beta_7 & \beta_8 & \beta_9 & \beta_{10} \\ \beta_5 & \beta_6 & \beta_7 & \beta_8 & \beta_9 & \beta_{10} & \beta_{11} \\ \beta_6 & \beta_7 & \beta_8 & \beta_9 & \beta_{10} & \beta_{11} & \beta_{12} \\ \beta_7 & \beta_8 & \beta_9 & \beta_{10} & \beta_{11} & \beta_{12} & \beta_{13} \\ \gamma_1 & \gamma_2 & \gamma_3 & \gamma_4 & \gamma_5 & \gamma_6 & \gamma_7 \\ \gamma_2 & \gamma_3 & \gamma_4 & \gamma_5 & \gamma_6 & \gamma_7 & \gamma_8 \\ \gamma_3 & \gamma_4 & \gamma_5 & \gamma_6 & \gamma_7 & \gamma_8 & \gamma_9 \\ \gamma_4 & \gamma_5 & \gamma_6 & \gamma_7 & \gamma_8 & \gamma_9 & \gamma_{10} \\ \gamma_5 & \gamma_6 & \gamma_7 & \gamma_8 & \gamma_9 & \gamma_{10} & \gamma_{11} \\ \gamma_6 & \gamma_7 & \gamma_8 & \gamma_9 & \gamma_{10} & \gamma_{11} & \gamma_{12} \\ \gamma_7 & \gamma_8 & \gamma_9 & \gamma_{10} & \gamma_{11} & \gamma_{12} & \gamma_{13} \end{pmatrix}$$

is equal to

$$\begin{cases} 1 & \text{if } i = 0 \\ 21 & \text{if } i = 1 \\ 378 & \text{if } i = 2 \\ 6832 & \text{if } i = 3 \\ 108096 & \text{if } i = 4, \\ 1714560 & \text{if } i = 5, \\ 27276288 & \text{if } i = 6 \\ 2^{35} - 3553 \cdot 2^{13} & \text{if } i = 7, \end{cases}$$

The number of solutions
($Y_1, Z_1, U_1, V_1, Y_2, Z_2, U_2, V_2, Y_3, Z_3, U_3, V_3$) of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + Y_3 Z_3 = 0, \\ Y_1 U_1 + Y_2 U_2 + Y_3 U_3 = 0, \\ Y_1 V_1 + Y_2 V_2 + Y_3 V_3 = 0, \end{cases}$$

satisfying the degree conditions

$$\deg Y_i \leq 6, \quad \deg Z_i \leq 2, \quad \deg U_i \leq 6, \quad \deg V_i \leq 6 \quad \text{for } 1 \leq i \leq 3.$$

is equal to

$$\begin{aligned} R_3(7, 3, 4) &= \int_{\mathbb{P} \times \mathbb{P} \times \mathbb{P}} g_{7,3,4}^3(t, \eta, \xi) dt d\eta d\xi = 2^{37} \cdot \sum_{i=0}^7 \Gamma_i \left[\begin{smallmatrix} 3 \\ 3+4 \\ 3+4 \end{smallmatrix} \right]_{\times 7} \cdot 2^{-3i} \\ &= 2^{37} \cdot (1 + 21 \cdot 2^{-3} + 378 \cdot 2^{-6} + 6832 \cdot 2^{-9} + 108096 \cdot 2^{-12} + 1714560 \cdot 2^{-15} \\ &\quad + 27276288 \cdot 2^{-18} + (2^{35} - 3553 \cdot 2^{13}) \cdot 2^{-21}) = 4243395 \cdot 2^{29} \end{aligned}$$

Example. The fraction of square triple persymmetric $\begin{bmatrix} s & m \\ s+m \end{bmatrix} \times (3s+2m)$ matrices which are

$$\text{invertible is equal to } \frac{\Gamma_{3s+2m} \left[\begin{smallmatrix} s & m \\ s+m \end{smallmatrix} \right]_{\times (3s+2m)}}{\sum_{i=0}^{3s+2m} \Gamma_i \left[\begin{smallmatrix} s & m \\ s+m \end{smallmatrix} \right]_{\times (3s+2m)}} = \frac{21}{64}.$$

REFERENCES

- [1] Cherly, Jorgen.
Exponential sums and rank of persymmetric matrices over $\mathbf{F}_2$
arXiv : 0711.1306, 46 pp
- [2] Cherly, Jorgen.
Exponential sums and rank of double persymmetric matrices over $\mathbf{F}_2$
arXiv : 0711.1937, 160 pp

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE BREST, 29238 BREST CEDEX 3, FRANCE
E-mail address: Jorgen.Cherly@univ-brest.fr